



Journal of Bioscience and Applied Research

JBAAR

WWW.JBAAR.ORG



Effect of Sulforaphane and Methotrexate combined treatment on Histone Deacetylase Activity in Solid Ehrlich Carcinoma

Mai M. El-Keey¹; Mohamad A. El Ghonamy^{*1}; Tarek M. Ali¹; Wafaa M. Ibrahim²; Ehab Tousson³.

¹ Biochemistry department, Faculty of Science, Tanta University.² Medical Biochemistry department, Faculty of Medicine, Tanta University.³ Zoology department, Faculty of Science, Tanta University.

mh_7akeem@hotmail.com

Abstract

Histone acetylation is one of the posttranslational modification that plays a role in the regulation of gene expression, also modulation of acetylation with histone deacetylase inhibitors (HDACi) had been considered a novel strategy for cancer chemoprevention. This study was aimed to investigate the activity of histone deacetylase enzyme (HDAC) in tumor cell model Solid Ehrlich carcinoma under the effect of the antitumor drug methotrexate in combination with sulforaphane (SFN) which is a natural compound found in cruciferous vegetables. In the present study sulforaphane was extracted from cabbage outer leaves. The HDAC activity significantly increased in tumor bearing mice, non-significantly decreased after treatment with methotrexate and significantly decreased after treatment with methotrexate with oral supplement by sulforaphane. However, The DNA damage decreased significantly in tumor tissue of tumor bearing mice after treatment with methotrexate and increased significantly in tumor tissue of tumor bearing mice treated with oral sulforaphane in combination with methotrexate treatment after carcinogenesis. In conclusion the sulforaphane have antitumor effect via its pro-oxidant activity and affect the HDAC activity via its metabolite's in an inhibitory manner that enhanced the effect of methotrexate.

Key words: histone deacetylase (HDAC), sulforaphane (SFN), DNA damage.

1. Introduction

According to recent estimates, cancer continues to remain the second leading cause of death and is becoming the leading one in old age (Donnellan et al., 2017). It is projected that by

2030 the number of new cancer cases will increase by 70% worldwide due to demographic changes alone (Bao et al., 2017). Experimental tumors have great importance for the purposes of modeling, and one of the commonest is the Ehrlich ascitis tumor, which is derived from a

spontaneous murine mammary adenocarcinoma (Ceylan et al., 2017). Mutation studies have suggested that chronic oxidative stress, particularly from chronic inflammation, is associated with carcinogenesis (Choudhury et al., 2017). In addition to inducing DNA, lipid, and protein damage, oxidative damage to protein-coding or -noncoding RNA may potentially cause errors in protein synthesis or dysregulation of gene expression. This has been proposed as an underlying mechanism of several human diseases (Nunomura, 2016). When the pro-oxidant/anti-oxidant equilibrium is lost, oxidative stress is generated, altering and damaging many intracellular molecules, including DNA, RNA, lipids and proteins (Margaritelis et al., 2016). Upon exposure to oxidants or oxidative stress-inducing agents, a common adaptive response induced in mammalian cells is the upregulation of stress-response genes, many of which encode antioxidant defense enzymes. While high levels of ROS production may lead to the induction of apoptosis or necrosis, increasing evidence demonstrates that low or transient ROS exposure increases cell proliferation, likely through altered expression of growth factors and proto-oncogenes (Tharmalingam et al., 2017).

There are two protein families with HDAC activity; Sirtuins and the classical HDAC family (Hsieh et al., 2016, Serrano-Gomez et al., 2016). HDACi function by blocking access to the active site (reversible or irreversible) of HDAC (Lee et al., 2017), later it was discovered that HDACi have potent proliferation-inhibitory properties with cancer cells (Shao et al., 2017).

Isothiocyanates (ITCs) are synthesized from glucosinolates (GSs) stored in plants, upon catalytic breakdown by myrosinase, a thioglucoside glucohydrolase present mainly in the crucifers and released during stress or damage and to some extent in the microflora of the intestinal tract (Li et al., 2017); ITCs rapidly accumulate in the cells, and such accumulation levels may reach 100- to 200-fold over extracellular levels (Sosa et al., 2013). Sulforaphane (SFN) was first isolated from broccoli in the early 1990s as an inducer of

phase 2 enzymes (Xenobiotic metabolism) and since then numerous studies have proposed various anti neoplastic pharmacological aspects of SFN, thereby suggesting its potential as a promising candidate in cancer chemoprevention (Zhang et al., 2007).

This study was aimed to investigate the activity of histone deacetylase enzyme (HDAC) in tumor cell model Solid Ehrlich carcinoma comparing normal tissue and tumor tissue, under treatment with the anti-tumor drug methotrexate and the naturally isothiocyanate sulforaphane alone and in combination with methotrexate.

2. Materials and Methods

2.1. Extraction of Sulforaphane:

Crucifer aqueous extracts were prepared according to the method of Bertelli et al and Dandan Han et al (Bertelli et al., 1998, Han et al., 2011) with few modifications. Briefly, plant material was ground to a fine homogenous powder then 0.5 gram of powdered material was left to autolyse in 7ml HCl pH 3.0 (0.001 N) at 37 C° over night to allow complete hydrolysis of the sulphur glycosides. The supernatant was filtrated through a piece of gauze then extracted with 5ml Dichloromethane three times then the dichloromethane fraction was salted with sodium sulphate anhydrous and dehydrated using a rotary evaporator at 30°C. The residue was dissolved in 2ml of 20% v/v acetonitrile and was filtered and injected to solid-phase extraction silica column "10 cm × 2.5 cm silica" that was previously equilibrated with dichloromethane, the solid-phase extraction silica column then washed with ethyl acetate as a washing solvent that will not remove sulforaphane but the other unnecessary compounds, then dichloromethane was used as the elution solvent in subsequent steps where the largest relative amount of sulforaphane was washed out. The concentration of sulforaphane was determined using UV1102 spectrophotometer. The maximum wave length (λ_{max}) which is unique for sulforaphane is 240 nm and its molar attenuation coefficient ϵ which is $950 \text{ M}^{-1} \text{ cm}^{-1}$ (Tao et al.). Sulforaphane was detected by GC-MS using Perkin Elmer Clarus

580 Gas chromatograph according to the method described by Matusheski et al (Matusheski et al., 2001), followed by checking the mass spectrum of the extracted sulforaphane in MAINLIB library. More quantities were then extracted, dehydrated by rotary evaporation and dissolved in saline for further animal oral treatment.

2.2. Animals

In this study, 60 albino female mice weighing about 20–25 g were housed in wire mesh cages and were fed standard rat chew and allowed free access to water. They were kept under constant environmental conditions [temperature (23 ± 2 °C), relative humidity ($80 \pm 5\%$) and light (12 h light/dark cycles)].

All experiments were carried out according to the guidelines of the ethical committee of Faculty of Science, Tanta University.

2.3. Solid Ehrlich Carcinoma (SEC) Tumor Model

A model of SEC was used, where 1×10^6 of the Ehrlich carcinoma cells (ECC) obtained from the Pharmacology and Experimental Oncology Unit of the NCI, Cairo University, Egypt were implanted subcutaneously into the right thigh of the hind limb of mice. A palpable solid tumor mass (about 100mm³) was developed within 12 days.

MST (Median survival time) was monitored by recording the mortality daily for 4 weeks and % ILS (increase in life span percent) was calculated (Tanongkankit et al., 2013).

2.4. Experimental design:

Mice were randomly equally divided into the following groups:

- Group I (Negative control group): This group included 10 untreated mice.

The remaining mice were subjected to Ehrlich tumor cells which were implanted

subcutaneously into the right thigh of the hind limb of mice and divided as follow:

- Group II: Tumor bearing group “Positive control group”.
- Group III: Methotrexate “MTX” provided from Sanofi Aventis was intraperitoneal injected once each 3 days of a dose 1.25 mg/Kg for one month.
- Group IV: 1.5 mg of SFN dissolved in 0.5 ml saline “50 mg/Kg”, delivered by gastric tube once daily one day before Ehrlich cells implantation and continued for one month.
- Group V: 1.5 mg of SFN dissolved in saline, delivered by gastric tube once daily one day after Ehrlich cells implantation and continued for one month.
- Group VI (Methotrexate and sulforaphane treated group): 1.5 mg of SFN dissolved in 0.5 ml saline, delivered by gastric tube once daily with intraperitoneal MTX once each 3 days of a dose 1.25 mg/Kg were given one day after subcutaneous implantation of Ehrlich cells and continued for one month.

2.5. Tissue Sampling:

After scarification, Tumors were dissected carefully and washed three times with ice cold saline to remove extraneous materials, the tumor was chilled on ice and divided into four pieces that were wrapped in aluminum foil and stored at -80 °C till used for preparation of tissue homogenates and nuclear extracts (Chastre et al., 1984).

Muscles from the right thigh of normal group were dissected, and washed three times with ice cold saline, the tissue was chilled on ice and divided into four pieces that were wrapped in aluminum foil and stored at -80 °C till used for preparation of tissue homogenates and nuclear extracts (Chastre et al., 1984).

2.6. Preparation of Tumor Nuclear Protein Extracts:

for estimation of total protein content and HDAC activity by using membrane, nuclear and cytoplasmic protein extraction kit supplied by Biobasic INC. Canada(Ouyang et al., 2009).

2.7. Biochemical assays:

Estimation of protein content in the nuclear extract of tumor tissue according to the method of lowry et al. (1951) (Lowry et al., 1951); Histone deacetylase activity was determined using colorimetric HDAC activity assay kit supplied by BioVision USA, Inc.; assessment of DNA damage using the diphenylamine method

(Zhu et al., 1998); determination of Lipid Peroxide “Malondialdehyde (MDA)” using Lipid peroxide colorimetric assay kit supplied by BioDiagnostic Egypt and determination of total antioxidant capacity using total antioxidant capacity colorimetric assay kit supplied by BioDiagnostic Egypt.

2.8. Statistical analysis

One-way analysis of variance (ANOVA) was used to assess significant differences among groups and the Tuckey test was used to compare all groups with each other and showed the significant effect of treatment.

3. Results and Discussion




Figure 1: HDAC expressed as μM deacetylated lysine per μg extracted nuclear protein, DNA damage expressed as Percent, Tumor MDA expressed as $\text{nmol}/10\text{g}$ tissue, Tissue TAC expressed as mM/g tissue




Figure 2: Forward linear correlation between the HDAC activities and fragmented DNA and MDA.

In the present work the naturally isothiocyanate sulforaphane was extracted from outer leaves of green cabbage (*Brassica oleracea*) in a concentration of 83.3 µg/g fresh weight (Figure 3). Farag and Motaal extracted SFN from cabbage at a concentration of 540 µg/g (Farag et al., 2010). The differences in sulforaphane content are associated with the variability of plant cultivars and multiple environmental factors, e.g. daily temperatures, soil fertility, pathogen challenge, wounding and plant growth regulators (Fernández-León et al., 2017, Francisco et al., 2017).

HDACs were found to be overexpressed in numerous types of cancer (Spehalski et al., 2017). In the current study all mice bearing Ehrlich tumors showed significant increase in HDAC activity in tumor tissue, comparing normal mice muscle tissue. A significant decrease in HDAC activity was observed after treatment with sulforaphane alone or in combination with the antitumor drug methotrexate.

In agreement with the present data Ho et al. demonstrated that sulforaphane inhibits HDAC activity in human colorectal and prostate cancer cells causing enhanced histone acetylation, derepression of P21 and Bax, and induction of cell cycle arrest/apoptosis, leading to cancer prevention (Ho et al., 2009, Johnson et al., 2017).

In the present work there was a significant increase in DNA damage in tumor tissue comparing normal tissue, but there was a significant decrease in DNA damage in tumor tissue of mice treated only with methotrexate while the percent of DNA damage increased non-significantly after treating the mice bearing tumors with sulforaphane alone or in combination with methotrexate treatment.

Ferreira de Oliveira et al (2014) also reported that the MG-63 osteosarcoma cell line was exposed to SFN at a dietary concentration (5 µM) which resulted in genomic instability as confirmed by increased number of DNA breaks (Ferreira de Oliveira et al., 2014, Coutinho et al., 2017).

Malondialdehyde (MDA) is the principal and most studied product of polyunsaturated fatty acid peroxidation. This aldehyde is a highly toxic molecule and should be considered as more than just a marker of lipid peroxidation. Its interaction with DNA and proteins has often been referred to as potentially mutagenic and atherogenic (Carrillo et al., 2017).

In the present work the MDA level increased significantly in all tumor tissue and non-significantly increased in tumor tissue treated with both methotrexate and sulforaphane.

Likewise De Oliveira et al observed that the increase in intracellular ROS was not gradual

between the 5 and 10 mM SFN exposure, and a sharp increase was observed particularly for the 48-h exposure period. Moreover, for 10 mM SFN, longer exposure times resulted in larger accumulation of intracellular ROS (Ferreira de Oliveira et al., 2014, Coutinho et al., 2017).

The present work recorded a positive significant correlation between HDAC activity and fragmented DNA percentage. In agreement with the data reported by Shan et al. that class I HDACs, HDAC2 expression was dramatically correlated with expression of a set of DNA damage repair (DDR) genes, including RAD51, in breast cancer. Treatment with the HDAC inhibitor mocetinostat in two basal-like breast cancer cell lines, SUM149 and HCC1937, decreased expression of RAD51 at both mRNA and protein levels, indicating a possible mechanism by which HDAC2 regulates RAD51 expression in breast cancer (Shan et al., 2017). Robert and Rehman also previously reported that HDACi can inhibit DNA repair responses in a few cell lines, which might increase the sensitivity of tumour cells to chemotherapy and radiotherapy by leading to increased DNA damage by these treatments (Robert et al., 2016, Rehman et al., 2016).

The present study recorded a positive significant correlation between HDAC activity and tumor MDA level. Consistently Portakal et al. have found that MDA content in cancerous tissues of the breast was higher than in the corresponding noncancerous tissues (Portakal et al., 2000).

4. Conclusion

SFN is a naturally ITC that found in cabbage outer leaves and showed an inhibitory effect to HDAC in Ehrlich solid tumor tissue when given orally. SFN also showed a pro-oxidant activity in tumor tissue that enhanced the tumor treatment with MTX.

5. Acknowledgements

Thanks to Dr Marwa Mohammed Atef Khalid for supplying chemicals used in DNA damaged assay.

6. References

- E. Donnellan and A. A. Khorana, (2017): Cancer and Venous Thromboembolic Disease: A Review. *The Oncologist*. 22: 199-207.
- J. Bao and L. Qiao, (2017): New developments on targeted cancer therapy. *Cancer letters*. 387: 1-2.
- D. Ceylan, A. Aksoy, T. Ertekin, A. H. Yay, M. Nisari, G. Ş. Karatoprak and H. Ülger, (2017): The effects of gilaburu (*Viburnum opulus*) juice on experimentally induced Ehrlich ascites tumor in mice.
- G. Choudhury and W. MacNee, (2017): Role of Inflammation and Oxidative Stress in the Pathology of Ageing in COPD: Potential Therapeutic Interventions. *COPD: Journal of Chronic Obstructive Pulmonary Disease*. 14: 122-135.
- A. Nunomura, (2016): Idiopathic basal ganglia calcification (Fahr's disease) and dementia. *Psychiatry Clin Neurosci*. 70: 129-30.
- N. V. Margaritelis, J. N. Cobley, V. Paschalis, A. S. Veskokoukis, A. A. Theodorou, A. Kyparos and M. G. Nikolaidis, (2016): Going retro: Oxidative stress biomarkers in modern redox biology. *Free Radic Biol Med*. 10.1016/j.freeradbiomed.2016.02.005
- S. Tharmalingam, S. Sreetharan, A. V. Kulesza, D. R. Boreham and T. Tai, (2017): Low-Dose Ionizing Radiation Exposure, Oxidative Stress and Epigenetic Programming of Health and Disease. *Radiation Research*.
- C.-L. Hsieh, H.-P. Ma, C.-M. Su, Y.-J. Chang, W.-Y. Hung, Y.-S. Ho, W.-J. Huang and R.-K. Lin, (2016): Alterations in histone deacetylase 8 lead to cell migration and poor prognosis in breast cancer. *Life sciences*. 151: 7-14.
- S. J. Serrano-Gomez, M. Maziveyi and S. K. Alahari, (2016): Regulation of epithelial-

- mesenchymal transition through epigenetic and post-translational modifications. *Molecular cancer*. 15: 1.
- C.-H. Lee, J.-C. Lin, C.-L. Ho, M. Sun, W.-T. Yen and C. Lin, (2017): Efficacy and safety of micafungin versus extensive azoles in the prevention and treatment of invasive fungal infections for neutropenia patients with hematological malignancies: A meta-analysis of randomized controlled trials. *PloS one*. 12: e0180050.
- G. Shao, W. Lai, X. Wan, J. Xue, Y. Wei, J. Jin, L. Zhang, Q. Lin, Q. Shao and S. Zou, (2017): Inactivation of EGFR/AKT signaling enhances TSA-induced ovarian cancer cell differentiation. *Oncology Reports*. 37: 2891-2896.
- R. Li, D. Song, F. Vriesekoop, L. Cheng, Q. Yuan and H. Liang, (2017): Glucoraphenin, sulforaphane, and antiproliferative capacity of radish sprouts in germinating and thermal processes. *European Food Research and Technology*. 243: 547-554.
- V. Sosa, T. Moliné, R. Somoza, R. Paciucci, H. Kondoh and M. E. LLeonart, (2013): Oxidative stress and cancer: an overview. *Ageing research reviews*. 12: 376-390.
- Y. Zhang and L. Tang, (2007): Discovery and development of sulforaphane as a cancer chemopreventive phytochemical. *Acta pharmacologica Sinica*. 28: 1343-1354.
- D. Bertelli, M. Plessi, D. Braghiroli and A. Monzani, (1998): Separation by solid phase extraction and quantification by reverse phase HPLC of sulforaphane in broccoli. *Food Chemistry*. 63: 417-421.
- D. Han and K. H. Row, (2011): Separation and purification of sulforaphane from broccoli by solid phase extraction. *International journal of molecular sciences*. 12: 1854-1861.
- S. Tao, P. Liu, G. Luo, M. R. de la Vega, H. Chen, T. Wu, J. Tillotson, E. Chapman and D. D. Zhang, p97 Negatively Regulates NRF2 by Extracting Ubiquitylated NRF2 from the KEAP1-CUL3 E3 Complex. *Molecular and Cellular Biology*. MCB. 00660-16.
- N. V. Matusheski, M. A. Wallig, J. A. Juvik, B. P. Klein, M. M. Kushad and E. H. Jeffery, (2001): Preparative HPLC method for the purification of sulforaphane and sulforaphane nitrile from Brassica oleracea. *Journal of agricultural and food chemistry*. 49: 1867-1872.
- Y. Tanongkankit, S. S. Sablani, N. Chiewchan and S. Devahastin, (2013): Microwave-assisted extraction of sulforaphane from white cabbages: Effects of extraction condition, solvent and sample pretreatment. *Journal of Food Engineering*. 117: 151-157.
- J. Chastre, F. o. Viau, P. Brun, J. Pierre, M.-C. Dauge, A. Bouchama, A. Akesbi and C. Gibert, (1984): Prospective Evaluation of the Protected Specimen Brush for the Diagnosis of Pulmonary Infections in Ventilated Patients 1-3. *American Review of Respiratory Disease*. 130: 924-929.
- W. Ouyang, D. Zhang, J. Li, U. N. Verma, M. Costa and C. Huang, (2009): Soluble and insoluble nickel compounds exert a differential inhibitory effect on cell growth through IKK α -dependent cyclin D1 down-regulation. *Journal of cellular physiology*. 218: 205-214.
- O. H. Lowry, N. J. Rosebrough, A. L. Farr and R. J. Randall, (1951): Protein measurement with the Folin phenol reagent. *J biol Chem*. 193: 265-275.
- W.-H. Zhu, A. Majluf-Cruz and G. A. Omburo, (1998): Cyclic AMP-specific phosphodiesterase inhibitor rolipram and RO-20-1724 promoted apoptosis in HL60 promyelocytic leukemic cells via cyclic AMP-independent mechanism. *Life sciences*. 63: 265-274.

- M. A. Farag and A. A. A. Motaal, (2010): Sulforaphane composition, cytotoxic and antioxidant activity of crucifer vegetables. *Journal of Advanced Research*. 1: 65-70.
- A. M. Fernández-León, M. F. Fernández-León, D. González-Gómez, M. C. Ayuso and M. J. Bernalte, (2017): Quantification and bioaccessibility of intact glucosinolates in broccoli 'Parthenon' and Savoy cabbage 'Dama'. *Journal of Food Composition and Analysis*. 61: 40-46.
- M. Francisco, M. Tortosa, M. Martínez-Ballesta, P. Velasco, C. García-Viguera and D. Moreno, (2017): Nutritional and phytochemical value of Brassica crops from the agri-food perspective. *Annals of Applied Biology*. 170: 273-285.
- E. Ho, J. D. Clarke and R. H. Dashwood, (2009): Dietary sulforaphane, a histone deacetylase inhibitor for cancer prevention. *The Journal of nutrition*. 139: 2393-2396.
- G. S. Johnson, J. Li, L. M. Beaver, W. M. Dashwood, D. Sun, P. Rajendran, D. E. Williams, E. Ho and R. H. Dashwood, (2017): A functional pseudogene, NMRAL2P, is regulated by Nrf2 and serves as a coactivator of NQO1 in sulforaphane-treated colon cancer cells. *Molecular nutrition & food research*. 61: 1600769.
- J. M. P. Ferreira de Oliveira, C. Remédios, H. Oliveira, P. Pinto, F. Pinho, S. Pinho, M. Costa and C. Santos, (2014): Sulforaphane induces DNA damage and mitotic abnormalities in human osteosarcoma MG-63 cells: correlation with cell cycle arrest and apoptosis. *Nutrition and cancer*. 66: 325-334.
- L. Coutinho, H. Oliveira, A. R. Pacheco, L. Almeida, F. Pimentel, C. Santos and J. M. P. F. de Oliveira, (2017): Hesperetin-etoposide combinations induce cytotoxicity in U2OS cells: Implications on therapeutic developments for osteosarcoma. *DNA repair*. 50: 36-42.
- W. Carrillo, X. Guzmán and E. Vilcacundo, (2017): Native and Heated Hydrolysates of Milk Proteins and Their Capacity to Inhibit Lipid Peroxidation in the Zebrafish Larvae Model. *Foods*. 6: 81.
- W. Shan, Y. Jiang, H. Yu, Q. Huang, L. Liu, X. Guo, L. Li, Q. Mi, K. Zhang and Z. Yang, (2017): HDAC2 overexpression correlates with aggressive clinicopathological features and DNA-damage response pathway of breast cancer. *American journal of cancer research*. 7: 1213-1226.
- C. Robert, P. K. Nagaria, N. Pawar, A. Adewuyi, I. Gojo, D. J. Meyers, P. A. Cole and F. V. Rassool, (2016): Histone deacetylase inhibitors decrease NHEJ both by acetylation of repair factors and trapping of PARP1 at DNA double-strand breaks in chromatin. *Leukemia research*. 45: 14-23.
- M. U. Rehman, P. Jawaid, Q. L. Zhao, P. Li, K. Narita, T. Katoh, T. Shimizu and T. Kondo, (2016): Low-dose spiruchostatin-B, a potent histone deacetylase inhibitor enhances radiation-induced apoptosis in human lymphoma U937 cells via modulation of redox signaling. *Free radical research*. 50: 596-610.
- O. Portakal, Ö. Özkaya, M. Erden inal, B. Bozan, M. Koşan and I. Sayek, (2000): Coenzyme Q10 concentrations and antioxidant status in tissues of breast cancer patients. *Clinical Biochemistry*. 33: 279-284.