



BioBacta



Journal of Bioscience and Applied Research

www.jbaar.org



Marine pollution by some heavy metals and physiological response of *Ruditapes decussatus*

G.M El Khodary, E.H Radwan, M.M Elghazaly and D. El Bahnasawy

Damanhour University, Faculty of Science, Zoology department, Egypt

Corresponding author: dr_eman_hashem@yahoo.com

Abstract

Bivalves can accumulate toxicants such as heavy metals in their tissues, for this reason they are considered as a good bio-indicators for water quality. The risk is increased due to eating these clams raw or lightly cooked. The aim of this study is to determine the concentration of some selected metals (Cu, Zn, Mn, Cd, and Pb) in the soft tissue of *Ruditapes decussatus* collected in the summer of 2017, from three locations at Alexandrian coasts, Egypt, Abo Quir, (loc.#1), Sedi Beshr (loc.#2) and (loc.#3), El-Max) and to find out whether pollution alters the clam physiological functions or not. The present data showed that the highest mean value of salinity was reported in water samples collected from Abo Quir (loc.#1) and the highest level of dissolved oxygen was reported in water samples collected from Sedi Beshr (loc.#2). The present results showed that the studied heavy metals concentrations are highly significant in samples of water and soft tissue of *Ruditapes decussatus* collocated from Abo Quir bay. From the above cited results it is concluded that loc.#1(Abo Quir bay) represent the most polluted site in the present study. Statistical analysis showed significant increase of MDA and significant decrease of SOD and GPx in the soft tissue of *Ruditapes decussatus* collected from Abo-Quir Bay (loc.#1). There was high significant difference between the concentration of MDA, SOD and GPx in the tissue of clams collected from the three locations ($p < 0.001$). The correlation coefficient of heavy metals in tissue, heavy metals in water and oxidative stress biomarkers in showed that the concentration of MDA in the tissue of calm collected from Abo Quir Bay (loc.#1), Sedi Beshr (loc.#2) and El-Max (loc.# 3) were positively correlated with the mean activity level of SOD and the concentration of SOD in the clams tissue collected from Abo Quir Bay (loc.#1) and Sedi Beshr (loc.#2) were negatively correlated with the activity level of GPx. The present work also showed that the correlations between different heavy metals in the tissues of *R. decussatus* collected from three locations indicated that different rates and mechanisms of metal accumulation were taking place..

Key words: Environment, Bivalves, Heavy metals, Clam, Protein, Oxidative stress.

Introduction

Sea food is an important nutritious food source worldwide and appears in all kinds of popular dishes. Since bivalves are filter feeders and have a tendency to bio accumulate heavy metals at higher concentration more than those of the surrounding sea water (Cosson, 2000; Fang *et al.*, 2003). The risk is enhanced by the fact that these shellfish are eaten raw or relatively lightly cooked (Crocì *et al.*, 2002; Formiga-Cruz *et al.*, 2003). The importance of oxidative stress response as potential biomarkers of environmental pollution has been addressed by different experimental approaches (Orbea *et al.*, 2002; Rodríguez-Ariza *et al.*, 2003, Ferreira *et al.*, 2005 and Regoli *et al.*, 2014). Metal accumulation can cause an increase in Reactive Oxygen Species (ROS) like hydrogen peroxide, superoxide anion radical, hydroxyl radical, leading to oxidative stress (Livingstone, 2001). Antioxidant defense systems that prevent the formation of ROS include the antioxidant enzymes such as superoxide dismutase (SOD) and catalase (CAT) that have been extensively used as biomarkers of oxidative stress (Ferreira *et al.*, 2005, 2007 and Vrankovic, 2015).

In Egypt, *Ruditapes decussatus* is an abundant benthic bivalve that in direct contact with heavy metals both of natural and anthropogenic origin. This clam is an example of a filter-feeding marine organism satisfying many criteria required for being a biological indicator (Bebiano, 2003). Marine coast are characterized by high fluctuations in chemical and physical parameters and in most cases there exist a significant occurrence of human activities including industrial, domestic and agricultural waste discharges (Meherm, 2002). The present work was designed to determine the concentration of some selected metals (Cu, Zn, Mn, Cd, and Pb) in sea water and the soft tissue of *Ruditapes decussatus* collected from Abo- Quir, Sedi Beshr and EL Max in

Alexandria sea water, Egypt. In addition the oxidative stress biomarkers (lipid peroxidation (MDA); and antioxidant enzymes; superoxide dismutase (SOD) and Glutathione peroxidase (GPx) were measured to evaluate the physiological response of *Ruditapes decussatus*.

Material and methods

The study areas and sample collection: The present investigation was carried out at three locations in the Mediterranean Sea Alexandria, Egypt). The first location is Abo-Quir Bay (loc.#1) which receives different pollutants contributing to various waste source categories discharged through three main openings namely; El-Tabia pumping station, outlet of Lake Edku (Boughaz El-Maddya) and Rosetta mouth of the Nile River (Shreadah and Tayel, 1992). The second selected area is Sedi Beshr (loc. #2) which not have any industrial activity like the other two locations Abo-Quir Bay and El-Max (Fig. 1). Water sample is fairly clean. Due to existence of resorts it turns to relatively few polluted water (Amin and Galal, 2000).

The third selected area is Max (loc.#3) which affected by a mixed agricultural runoff and industrial wastes from a chloro-alkali plant and receives airborne particles from the fumes of adjacent industrial plants, including an oil refinery and cement factory. *Ruditapes decussatus* clams were collected from the three locations; Abo Quir (loc.#1), Sedi Beshr (loc.#2) and El Max (loc.#3) during Summer 2017. Dead or damaged specimens were eliminated and a standardized shell size (3.5-3.8mm) was used. Also, Water samples were collected from a precise depth corresponding to the clam settlements from each location.

Physicochemical analysis of sea water: Determination of physicochemical parameters (salinity, dissolved oxygen and pH) was

measured in sea water samples. Salinity was determined using Inductive Salinometer (Beckman mode) according to (Grasshoff, 1976). The pH-value of water sample was measured using Bench type (JENWAY, 3410 Electrochemistry Analyzer pH-meter).

dissolved oxygen determination, was done according to a modified Winkler’s method (Grasshoff, 1976).



Fig.1. location map of the study sites.

Analysis of heavy metals: The concentrations of heavy metals: Copper (Cu), Zinc (Zn), Manganese (Mn), Cadmium (Cd), and lead (Pb), in water samples and soft tissues of *Ruditapes decussatus* were measured using the atomic absorption spectrophotometer by the method described by UNEP/FAO/IAEA/IOC, (1984); El-Sikaily *et al.* (2004). The results of metal concentrations in water samples and soft tissues were expressed as ($\mu\text{g/l}$) and ($\mu\text{g/g}$) of dry weight respectively.

Physiological and biochemical analysis: The determination of biochemical markers of oxidative stress in the soft tissue of *Ruditapes decussatus* occurred by the measurement of lipid peroxidation (MDA); according to the method

of Yoshioka *et al.* (1979), and antioxidant enzymes; superoxide dismutase (SOD) and Glutathione peroxidase (GPx); were determined according to the method of Beauchamp and Fridovich, (1971) and Rao (1996); respectively. Also, total protein content (T.p.) was measured according to method described by Lowry *et al.* (1951).

Statistical analysis: Data were fed to the computer and analyzed using IBM SPSS software package version 20.0. (Armonk, NY: IBM Corp).The Kolmogorov- Smirnov, Shapiro and D'agstino tests were used to verify the normality of distribution of variables, F - test was used to compare two groups for normally distributed quantitative variables.

Pearson coefficient was used to correlate between quantitative variables. Significance of the obtained results was judged at the 5% level. The minimum level of statistical significance was set at $P < 0.05$.

Results

The physico-chemical parameters obtained from analysis of sea water samples collected

from the three studied locations; Abo Quir Bay (loc.#1), Sedi Beshr (loc.#2) and El-Max (loc.#3) collected in Summer 2017 are presented in Table 1 and Figs. 2 & 3. The present data showed that the highest mean value of salinity was reported in loc.#1 as 36.74% while the highest mean value of dissolved oxygen were; 10.0 ± 0.19 mg/l in (loc.#1). The temperature ranged between (34.6- 37 .5) in the studied location.

Table (1): The mean and standard deviation (Mean±S.D.) of Physico-chemical parameters of water samples collected from; Abo Quir Bay (loc.#1), Sedi Beshr (loc.#2) and El-Max (loc.#3) .

Physico-chemical parameters	Abo Quir	Sedi Beshr	El Max	F	p
Salinity‰	36.74 ^a ±0.98	34.01 ^b ±0.79	34.37 ^b ±0.72	18.841*	<0.001*
pH	7.84 ± 0.10	7.75 ± 0.07	7.81 ± 0.09	1.544	0.246
Dissolved Oxygen (mg/L)	9.02 ^c ± 0.24	10.0 ^a ± 0.19	9.56 ^b ± 0.19	33.476*	<0.001*
Temperature	37.50 ^a ±1.52	34.67 ^b ±1.21	35.83 ^b ±4.02	1.831	0.194

F, p: F and p values for ANOVA test, significant between groups was done using Post Hoc Test (LSD).

*: Statistically significant at $p \leq 0.05$. Data was expressed using Mean ± SD.

F ratio: Frequency, P Value: Probability.

Means with Common letters are not significant (Means with Different letters are significant).

The heavy metal concentrations of water collected from the studied locations are presented in Table (2) and fig. (4) during summer 2017. The results showed that the heavy metals concentrations of Zn, Mn, Pb, Cu, and Cd is highly significant in samples of water collocated from Abo Quir bay; its concentrations were 5.22 ± 0.37 , 4.50 ± 0.49 , 5.43 ± 0.55 , 3.48 ± 0.43 and 3.59 ± 0.41 (µg/l)

respectively. The pattern of metals occurrence in the water samples of selected locations can be summarized as follows in descending order Abo Quir Bay (loc.#1) > El-Max (loc.#3) > Sedi Beshr (loc.#2). From the above cited results, it is indicated that; loc.#1 (Abo Quir Bay) represents the most polluted site in the present study.

Table (2): Mean and standard deviation (Mean±S.D.) of heavy metals concentration in water (µg/l) collected from Abo Quir Bay (loc.#1), Sedi Beshr (loc.#2) and El-Max (loc.#3) .

Heavy metals in water (µg/l)	Abo Quir(loc.#1)	Sedi Beshr(loc.#2)	El Max(loc.#3)	F	p
Zn	5.22 ^a ± 0.37	2.70 ^c ± 0.63	3.67 ^b ± 0.53	35.891*	<0.001*
Mn	4.50 ^a ± 0.49	2.53 ^b ± 0.40	2.92 ^b ± 0.41	34.669*	<0.001*
Pb	5.43 ^a ± 0.55	3.73 ^b ± 0.44	4.13 ^b ± 0.28	24.860*	<0.001*
Cu	3.48 ^a ± 0.43	2.03 ^c ± 0.20	2.87 ^b ± 0.31	29.510*	<0.001*
Cd	3.59 ^a ± 0.41	1.89 ^b ± 0.33	2.04 ^b ± 0.28	45.343*	<0.001*

Means with Common letters are not significant (Means with Different letters are significant)

F, p: F and p values for ANOVA test, significant between groups.*: Statistically significant at $p \leq 0.05$
 Data was expressed using Mean \pm SD. F ratio: Frequency, P Value: Probability.

Table (3) and fig. 5 showed that; the concentrations of Zn, Mn, Pb, Cu, and Cd in the soft tissue of *Ruditapes decussates* were accumulated in the samples collected from Abo Quir Bay (loc.#1) more than the concentration of metals in the samples collected from Sedi Beshr (loc.#2) and El-Max (loc.#3). It showed significant differences between the samples

taken from the three studied locations ($p \leq 0.001$) and its pattern of accumulation was Abo Quir Bay (loc.#1) >El-Max (loc.#3)> Sedi Beshr (loc.#2). The mean concentrations of Mn, Pb and Cd in soft tissue of the studied clam at Abo Quir Bay exceeded the allowable limit of (WHO, 1989).

Table (3): Mean and standard deviation (Mean \pm S.D.) of heavy metals concentration ($\mu\text{g/g}$) in the soft tissue of *Ruditapes decussatus* collected from Abo Quir Bay (loc.#1), Sedi Beshr (loc.#2) and El-Max (loc.#3)

Heavy metals In tissue of the clam ($\mu\text{g/g}$)	Number of replicants	Abo Quir	Sedi Beshr	El Max	WHO (1989) ($\mu\text{g/g}$)	F	p
Zn	5	6.35 ^a \pm 0.59	4.73 ^b \pm 0.48	5.32 ^b \pm 0.56	100	13.597*	<0.001*
Mn	5	5.95 ^a \pm 0.82	2.63 ^c \pm 0.53	4.08 ^b \pm 0.48	5.4	42.338*	<0.001*
Pb	5	5.50 ^a \pm 0.66	2.93 ^c \pm 0.41	4.47 ^b \pm 0.53	2	33.756*	<0.001*
Cu	5	5.03 ^a \pm 0.42	2.92 ^c \pm 0.40	3.92 ^b \pm 0.44	30	38.079*	<0.001*
Cd	5	5.25 ^a \pm 0.61	1.83 ^c \pm 0.33	3.33 ^b \pm 0.68	1	56.099*	<0.001*

Means with Common letters are not significant (Means with Different letters are significant)
 F, p: F and p values for ANOVA test, significant between groups *: Statistically significant at $p \leq 0.05$
 Data was expressed using Mean \pm SD. F ratio: Frequency, P Value: Probability.




Figure: (2) Mean and standard deviation (Mean \pm SD) of salinity and temperature in Summer (mg/l) between the three studied locations. Data in the same column with different superscript letters (a, b, and c) are significantly different, $p < 0.05$ (one-way ANOVA).




Figure (3): Mean and standard deviation (Mean± SD) of pH and dissolved oxygen (mg/l) in Summer between the three studied locations. Data in the same column with different superscript letters (a,b, and c) are significantly different, $p < 0.05$ (one-way ANOVA).




Figure (4): Mean and standard deviation (Mean± SD) of heavy metals in water (µg/l) in the three studied locations in Summer. Data in the same column with different superscript letters (a, b, and c) are significantly different, $p < 0.05$ (one-way ANOVA).




Figure (5): Mean and standard deviation (Mean± SD) of heavy metals in tissue (µg/g) in the three studied locations in summer. Data in the same column with different superscript letters (a, b, and c) are significantly different, $p < 0.05$ (one-way ANOVA).




Figure (6): Mean and standard deviation (Mean± SD) of oxidative stress biomarkers in tissue of the clam (U/g) from the three locations in Summer. Data in the same column with different superscript letters (a, b, and c) are significantly different, p<0.05(one-way ANOVA).




Figure (7): Mean and standard deviation (Mean± SD) of total protein in tissue of the clam (mg/l) from the three locations in summer. Data in the same column with different superscript letters (a, b, and c) are significantly different, p<0.05(one-way ANOVA).

Oxidative stress biomarker in the tissue of *Ruditapes decussatus* was showed significant difference in the three studied locations (Table 4, fig 6 and 7).The mean activity level of MDA at Abo Quir Bay recorded the highest value with value of 17.93±0.98 (U/g). Statistical analysis showed

significant decrease of SOD and GPx in the soft tissue of *Ruditapes decussatus* collected from Abo-Quir Bay (loc.#1). On the other side, total protein showed no significant differences in the three studied locations.

Table (4): Mean and standard deviation (Mean±SD) of oxidative stress biomarkers in tissue of the clam (U/g) and total protein (mg/l) in Summer collected from the three locations

Oxidative Stress Biomarkers (in tissue)	Abo-Quir Bay	Sedi Beshr	El-Max	F	p
MDA (U/g)	17.93 ^a ± 0.98	12.80 ± 1.06	15.50 ^b ± 1.13	35.295*	<0.001*
SOD (U/g)	0.72 ^b ± 0.09	1.07 ^a ± 0.22	0.80 ^b ± 0.10	9.665*	0.002*
GPx (U/g)	8.57 ^c ± 1.12	12.15 ^a ± 0.60	9.85 ^b ± 0.47	32.444*	<0.001*
Total protein (mg/l)	4.21 ± 0.55	4.76 ± 0.50	4.52 ± 0.40	1.957	0.176

Correlation coefficient of concentrations of heavy metals in sea water, heavy metals in tissue and oxidative stress biomarkers of *R. decussatus* collected from the studied locations were summarized in Tables 5, 6 and 7. The concentration of MDA in the tissue of clam collected from Abo Quir Bay (loc.#1), Sedi Beshr (loc.#2) and El-Max (loc.# 3) were positively correlated with the mean activity level of SOD at $r= 0.283$, 0.564 and 0.625 respectively. In contrast, the concentration of SOD in the tissue of *R. decussatus* collected from Abo Quir Bay (loc.#1) and sedi Beshr (loc.#2) were negatively correlated with the activity level of GPx. While, in El-Max (loc.# 3) concentration of SOD in the clam tissue was significant correlation ($p = 0.03$) with GPx ($r= 0.854$), negative significant correlation with the concentrations of Zn and Pb and negative correlation with Mn and Cd.

Table (6) showed high negative significant correlation ($P= 0.001$) between the concentrations of Zn with Pb ($r= -.976$) and Cu ($r= -0.970$) in the tissue of *R. decussatus* collected from Sedi Beshr (loc.#2). Negative significant correlation also demonstrated between the concentration of Mn and Cd ($p= 0.02$). In contrast, positive significant correlation were detected between the concentration of Pb with Cu ($r= 0.996$) and concentration of Mn with Cd ($r= 0.873$) in the tissue of *R. decussatus* collected from Sedi Beshr (loc.#2) and El-Max (loc.# 3) respectively. Moreover, positive significant correlation was showed between Zn in the tissue of clams with Pb ($r= 0.826$) and Cu ($r= 0.947$) in sea water collected from Abo Quir Bay (loc.# 1). The correlation among the concentration of heavy metals in sea water collected from the studied locations showed only a positive significant correlation between the concentrations of Pb with Cu collected from Abo Quir Bay (loc.#1) and El-Max (loc.# 3) ($r= 0.889$ and 0.832 respectively).

Discussion

In the present study, marine bivalves *Ruditapes decussatus* were collected from three areas of Alexandrian coast, Egypt, Abo Quir Bay (loc.#1), Sedi

Beshr (loc.#2) and El-Max (loc.#3). These species was identified in prior reports (Matias *et al.*, 2009). Developmental process in marine bivalves are recognized to be affected by different types of physical and natural factors involving nourishment quality temperature, salinity, photoperiod, sea hydrodynamics and biomechanics (Osman and Whitlatch, 2004; Bates, 2005; Lambert, 2005). Salinity and temperature and are two of the most important factors impacting the distribution of marine living forms (Epelbaum *et al.*, 2009).. In this study, water temperature and pH showed non-significant differences in the three examined locations. Yet, there was a significant difference in the other physico-parameters e.g. dissolved oxygen level and salinity. PH ratio is an important limiting chemical factor for aquatic life that may influence the biochemical reactions of the aquatic organism. The severe alteration in pH of the water may lead to harmful or fatal effect on aquatic organisms and as a result have an effect on the human and animal health. In general, water flows have pH range between 6 to 9 and any alteration in this range in pH can affect life forms in aquatic systems. If the pH value increases higher than this range, minor amounts of ammonia are required to reach a level that is toxic to fish, whereas when pH declines, acidity of the water increases affecting the fish (Murdoch, 1991). In the present study, the pH value of water samples collected from the studied locations ranged between 7.75–7.84.

. The highest salinity value is documented through summer may be describe to the higher degree of evaporation (Damotharan *et al.*, 2010). Higher salinity may probably be due to deposition of organic wastes, and inputs of different pollutant materials in this reach (Francis *et al.*, 2014). Similarly, in present study, the highest mean salinity (S‰) value of water sample collected from the three studied location recorded in Abo-Quir Bay with value of $36.74a \pm 0.98$.

The dissolved oxygen (DO) is one of the more important parameters available in the field of the water pollution control, as it allows the assessment of the aerobic conditions of a water-course, that receives discharge of pollutants (Parvez *et al.*, 2006). Benthic organisms have behavioral and

physiological adaptations for low dissolved oxygen. Organisms respond by decreasing metabolism and thus O₂ consumption (Wu, 2002). This can result in diminished growth and reproduction (Grove and Breitburg, 2005).). Even though it does not operate directly on growth as many toxins do, but limits the extent for aerobic metabolism. The most important natural physical factors affecting oxygen concentration in the aquatic environment are salinity and temperature. Dissolved O₂ solubility is reduced with increasing temperature and salinity (Ospar, 2005). These findings confirmed the present results.

R. decussatus is among the benthic invertebrates which are indirect contact with heavy metals both of anthropogenic and natural origin. The areas under investigation exposed to large amounts of industrial zone, that may reflect the reason of Abo- Quir Bay and El-Max have high concentration of heavy metals in the collected species. Utilization of heavy metals in industry, marine transport, and agriculture waste from El-Mahmoudia canal that flow off in El-Max leads to wide distribution of these components in marine environment. Their toxicity, potential for human exposure and environmental occurrence is dependant on a number of factors such as their concentration and source (Wu and Zhang, 2010, Radwan *et al.*, 2012; Radwan *et al.*, 2014; Radwan, 2016; Radwan *et al.*, 2016; Radwan *et al.*, 2017; Radwan *et al.*, 2018).

The heavy metals were accumulated within the clam soft tissues at high concentration levels amounting to thousands of times those accumulating in water due to filter-feeding of these aquatic organisms (Salman, 2011). The gained results demonstrate the process of bio-magnification across the tropic levels. This phenomenon was detected in other bivalves when put in comparison to water (El-Shenawy, 2002; El-Gamal, 2011). Filter

feeder oyster uptake dissolved metals ions in solution or in particular form, via food by filtering large volumes of water each day (Sajwan *et al.*, 2008). There are multiple ways to preserve homeostasis of essential metals to detoxify non-essential metals in the clam's tissues which include binding metals to low-molecular weight proteins, such as storing them within metal-containing granules or lysosomes, so that they can accumulate metals within their tissues in higher levels than their ambient water (Marigomez, 2002 and Jitar *et al.*, 2015). Consequently, *R. decussatus* was considered as a possible bio-monitor of contamination with heavy metals.

In the present study, a high accumulation of the selected heavy metals during summer 2017 were noticed in the tissues of *R. decussatus* collected from the three studied locations. The metabolism is sensitive to water temperature. Thus, any seasonal changes in the temperature could affect the metabolism with subsequent affection on the detoxification rate and accumulation of toxic materials. The metabolic acceleration owing to heat may lead to acceleration of metal accumulation within bivalve tissues, whereas, the reduction in the metabolic rates of bivalves once the environment becomes colder, may diminish the rate of assimilation of metals (Lannig *et al.*, 2006).

In the present work, positive significant correlation was showed only between Zn in the tissue of clams with Pb ($r= 0.826$) and Cu ($r= 0.947$) in sea water collected from Abo Quir Bay (loc.# 1). Like in other studies (Tessier *et al.*, 1984; Ahn *et al.* 2001), few correlations between metal concentrations in *A. umbonella* digestive glands and the environment were detected. The present work also showed that the correlations between different heavy metals in the tissues of *R. decussates* collected from three

locations indicated that different rates and mechanisms of metal accumulation were taking place. This might be due to differences in toxico - kinetic properties and efficient pathways of detoxification found in bivalves (Yap *et al.*, 2002). Metallothioneins (MTs) are low molecular weight non-enzymatic proteins that are heat stable and rich in cysteine, free of aromatic amino acids. The thiol groups of cysteine residues allow MTs to bind essential and non-essential metals with high affinity (Amiard *et al.*, 2006). MTs play a role in the homeostatic control of essential metals (Cu, Zn) to accomplish metabolic and enzymatic demands (Roesijadi, 1996). They also have an important role in the detoxification of non-essential trace metals such as Ag, Cd and Hg, that protects organisms against oxidative stress by scavenging the intracellular free radicals (Langston *et al.*, 1998 and Li *et al.*, 2013). Recent studies have showed that Cd can accumulate in marine invertebrates, and this can result in an elevation of the intracellular level of Metallothioneins (Amiard *et al.*, 2006). Furthermore, Duffus, (1980) indicated that Zn reduced the Cd concentration as it increases the rates of formation of the metallothionein protein. This may show that the negative significant correlation ($P= 0.001$) between the concentrations of Zn with Pb and Cu and negative correlation between Zn with Cd in the tissue of *R. decussates* collected from Sedi Beshr (loc.#2) and El-Max (loc.#3).

The pollution toxicity in aquatic organisms may be connected to an increased production of 'reactive oxygen species' (ROS) leading to great oxidative damage (Livingstone, 2001). Under normal conditions, cells possess antioxidant defenses that prevent the generation of ROS, and repair or degrade oxidatively modified molecules (Halliwell and Gutteridge, 1999). Alterations in the levels of antioxidants have been considered as biomarkers of contaminant-mediated pro-oxidant challenge in a variety of aquatic organisms, including mussels (Regoli *et al.*, 2004). ROS can be extremely toxic to aquatic organisms such as mussels, often leading to lipids oxidation in

membranes (MDA is an end product of lipid peroxidation LPX), protein oxidation and DNA damage (Almeida *et al.*, 2007). Ferreira *et al.*, 2005 stated that this oxidative damage could occur when detoxifying and antioxidant systems are unable to neutralize the active intermediates produced by xenobiotic and their metabolites. In the current study, the mean activity level of MDA in the tissues of *R. decussates* collected from Abo Quir Bay and El-Max (loc.#3) were significantly higher than those found in the tissues of clams collected from Sedi Beshr (loc.#2). Pampanin *et al.* (2005) showed that lipid peroxidation is a familiar mechanism of cellular injury in vertebrate and invertebrates, and it considered as an indicator of oxidative damage in tissues. Consequently, mean activity of MDA was recorded significant increase at Abo-Quir Bay, in where pollution toxicity increased.

The present study shows the mean level of SOD and GPX in the tissue of *R. decussatus* collected from Sedi Beshr (loc.#2) were higher than those collected from Abo Quir Bay and El-Max (loc.#3). Moreover, in El-Max (loc.# 3) concentration of SOD in the clam tissue shows negative significant correlation with the concentrations of Zn and Pb and negative correlation with the concentration of Mn and Cd. Enzymes with antioxidant properties such as SOD & GPx (Géret *et al.*, 2003) are defense biomarkers. The most important role of the antioxidant defenses is to stop the action of excess oxyradicals produced from exposure to xenobiotics (Valavanidis *et al.*, 2006). The high significant correlation between the mean activity of GPx with SOD in the tissue of *R. decussates* collected from El-Max (loc.#3) and negative correlation in tissues of clams collected from Abo Quir and Sedi Beshr (loc.#2) Bay. This may be due to that GPx activity increases more rapidly than other defenses during aerobic recovery in marine bivalves that is suggestive of their task as first responders to oxyradical detoxification (Pannunzio and Storey, 1998). Finally, it is important to accept that, although the accumulation of metals in *R. decussatus* has been shown to be in response to antioxidant

defense components, it cannot be considered to be due to the effect of heavy metals alone, but also due to other physical or chemical contaminants that may be present in these sites (El-Raey *et al.*, 2006). We can conclude that, the levels of the recorded metals exceed the maximum permissible levels except zinc and copper as mentioned by WHO (1989). This finding showed that *R. decussatus* collected from Abo Quir Bay and El-Max are not safe for human consumption. In addition the presence of heavy metals induces a biological stress and oxidative damage to *R. decussatus*.

References

- Amiard JC, Amiard-Triquet C, Barka S, Pellerin J, Rainbow PS (2006) Metallothioneins in aquatic invertebrates: Their role in metal detoxification and their use as biomarkers. *Aquat Toxicol* 76: 160–202.
- Amin, G. (2000): Whatever Happened to the Egyptians? Changes in Egyptian Society from 1950 to the Present. Cairo: American University in Cairo Press.
- Bates, W.R. (2005): Environmental factors affecting reproduction and development in Ascidians and other Protochordates. *Canadian Journal of Zoology*, 83(1): 51–61.
- Beauchamp, C. and Fridovich, I. (1971): Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry*, 44 (1): 276–87.
- Bebianno, M.J. and Serafim, M.A. (2003): Variation of metal and metallo-thionein concentrations in a natural population of *Ruditapes decussatus*. *Archives of Environmental Contamination and Toxicology*, 44: 53 – 66.
- Cosson, S; Ruxandra,. "Understanding dissociations in dyscalculia: a brain imaging study of the impact of number size on the cerebral networks for exact and approximate calculation." *Brain* 123.11 (2000): 2240-2255.
- Croci, T., Landi, M., Rinaldi-Carmona, M., Maffrand, J. P., Le Fur, G., & Manara, L. (2002). Modulation of gastric emptying and gastrointestinal transit in rats through intestinal cannabinoid CB1 receptors. *European journal of pharmacology*, 450(1), 77-83.
- Damotharan, P.; Perumal, N.V. and Perumal, P. (2010): Seasonal variation of physico-chemical characteristics of Point Calimere coastal waters (South east coast of India): Middle-East. *Science Research*, 6(4): 333-339.
- Duffus, J. (1980). *Environmental Toxicology Resource and Environmental Sciences Series*: Edward Arnold Publishers Ltd., London England.
- Duffus, J.H. (2002): Heavy Metals – A meaningless term. *Pure and Applied Chemistry*, 74: 793-807.
- El Gamal, M. (2011): The effect of depuration on heavy metal, petroleum hydrocarbon and microbial contamination levels in *Paphia undulata* (Bivalvia: Veneridae). *Czech Journal of Animal Science*, 56 (8):345-354.
- El-Raey, M.; Shalaby, E.; Guirguis, S.; Ghatass, Z.; Said, H.; Zahran, A.; Rashad, M. and Sivertsen, B. (2006): Ambient air pollution monitoring network over Alexandria city and Nile delta, Egypt. *Environmental Physics Conference*, 2:18–22.
- El Shenawy, S.N. (2002): The effect of metal bioaccumulation on glutathione and lipid peroxidation as biomarkers of aquatic ecosystem pollution of *Ruditapes decussatus* and *Venerupis pullastra* from Lake Timsah, Ismailia. *Egypt Journal of Zoology*, 39: 475-492.
- El Sikaily, A.; Khaled, A. and El Nemr, A. (2004): Heavy metals monitoring using bivalves from Mediterranean Sea and Red sea. *Environmental Monitoring Assessment*, 98 (1-3), 41-58.
- Epelbaum, L.M.; Herborg, T.W.; Therriault, C.M. and Pearce, C.M. (2009): Temperature and salinity effects on growth, survival, reproduction, and potential distribution of two non-indigenous botryllid ascidians in British Columbia. *Experimental Marine Biology and Ecology*, 369(1): 43–52.
- Fang, Z.; Cheung, Y.H.; and Wong, H. (2003): Heavy metals in oysters, mussels and clams collected from coastal sites along the Pearl River Delta, South China. *Environmental Science* 15(1): 9-24.
- Ferreira, A.G.; Machado, A.L. and Zalmon, I.R. (2005): Temporal and spatial variation on heavy metal concentrations in the oyster *Ostrea equestris* on the northern coast of Rio de Janeiro

state, Brazil. Brazilian Journal of Biology, 65: 67-76.

Ferreira, M.; Moradas, F.P. and Reis, H.M. (2007): The effect of long-term depuration on levels of oxidative stress biomarkers in mullets (*Mugil cephalus*) chronically exposed to contaminants. Marine Environmental Research, 64: 181–90 .

Formiga-Cruz, M., et al. "Evaluation of potential indicators of viral contamination in shellfish and their applicability to diverse geographical areas." Applied and environmental microbiology 69.3 (2003): 1556-1563.

Francis, O.; Arimoro; Robert, B.; Ikomi; and Nwadukwe, Ovie, D. Eruotor, Augustine, O. Edegbene, (2014): Fluctuating salinity levels and an increasing pollution gradient on fish community structure and trophic levels in a small creek in the Niger delta, Nigeria. International Aquatic Research, 7(4): 187-202.

Géret, F.; Serafim, A. and Bebianno, M.J. (2003): Variation of antioxidant enzyme activities, metallothioneins and lipid peroxidation levels in *Ruditapes decussatus*. Ecotoxicology, 12: 415–424.

Grasshoff, K. (1976): Methods of sea water analysis. Verlag. Chemie. Chapter 4; Determination of oxygen .

Grove, M. and Breitburg, D.L. (2005): Growth and reproduction of gelatinous zooplankton exposed to low dissolved oxygen. Marine Ecology Progress Series, 301: 185–198.

Halliwell, B. and Gutteridge, J.M.C. (1999): Free Radicals in Biology and Medicine, 3rd Edition. Oxford University Press, 1-25.

Jitar, O.; Teodosiu, C.; Robu, B.; Strungaru, S.A.; Plavan, G. and Nicoara, M. (2015): Environmental impact and risk assessment of the main pollution sources from the Romanian black sea coast. Environmental Engineering and Management Journal February, 14(2): 331-340.

Lambert, C.C. (2005): Historical introduction, overview, and reproductive biology of the Protochordates .Canadian Journal of Zoology, 83(1): 1–7.

Lannig, G.; Flores, J.F. and Sokolova, I.M. (2006): Temperature-dependent stress response in oysters, *Crassostrea virginica*: Pollution

reduces temperature tolerance in oysters. Aquatic Toxicology, 79: 278-287.

Langston WJ, Bebianno MJ, Burt GR (1998) Metal handling strategies in molluscs. In: Langston WJ, Bebianno MJ (Eds.), Metal Metabolism in Aquatic Environments. Chapman and Hall, London, pp. 219–283.

Li Y, Chai X, Wu H, Jing W, Wang L (2013) The response of metallothionein and malondialdehyde after exclusive and combined Cd / Zn exposure in the crab *Sinopotamon henanense* . PloS ONE 8(11): e80475 doi: 10.1371/journal.pone.0080475 [PMC free article] [PubMed]

Livingstone, D.R. (2001): Contaminant-stimulated reactive oxygen species production and oxidative damage in aquatic organisms. Marine Pollution Bulletin, 42: 656-66.

Lowry, O.H.; Rosebrough, N.J.; Farr, A.L. and Randall, R.J. (1951): Protein Measurement with the Folin Phenol Reagent. Journal of Biological Chemistry, 193: 265–275.

Marigómez, I.; Soto, M.; Cajaraville, M.P.; Angulo, E. and Giamberini, L. (2002): Cellular and subcellular distribution of metals in molluscs. Microscopy Research and Technique, 56 (5):358-92.

Matias, D.; Joaquim, S.; Leitão, A. and Massapina, C. (2009): Effect of geographic origin, temperature and timing of brood stock collection on conditioning, spawning success and larval viability of *Ruditapes decussatus* (Linné, 1758). Aquaculture International, 17(3):257-271 .

Meherm, M. "Assessment of the state of pollution by Antifouling paints in marine Environment of Alexandria coastal Region." PhD in Maritime transport technology (Environmental protection), Arab Academy for Science & Technology and Maritime transport (2002): 1-150.

Murdoch, T. (1991): Stream keeper's field guide: watershed inventory and stream monitoring methods. Adopt-a-Stream Foundation, Lewiston. (5):285-291.

Orbea, A.; Ortiz-Zarragoitia, M.; Solé, M.; Porte, C.; Cajaraville, M.P. (2002): Antioxidant enzymes and peroxisome proliferation in relation to contaminant body burdens of PAHs and PCBs in bivalve molluscs, crabs and fish from the

Urdaibai and Plentzia estuaries (Bay of Biscay). *Aquatic Toxicology*, 58: 75-98.

Osman, R.W. and Whitlatch, R.B. (2004): The control of the development of a marine benthic community by predation on recruits. *Experimental Marine Biology and Ecology*, 311: 117–145.

Ospar, (2005): Revised common procedure for the identification of the eutrophication status of the Oskar Maritime Area. Reference Number 5005-3. Oskar Commission.

Pampanin, D.M.; Camus, L.; Gomiero, A.; Marangon, I.; Volpato, E. and Nasci, C. (2005): Susceptibility to oxidative stress of mussels (*Mytilus galloprovincialis*) in the Venice Lagoon (Italy). *Marine Pollution Bulletin*, 50:1548-1557.

Pannunzio, T.M. and Storey, K. (1998): Antioxidant defenses and lipid peroxidation during anoxia stress and aerobic recovery in the marine gastropod *Littorina littorea*. *Experimental Marine Biology and Ecology*, 221: 277-292.

Parvez, S.; Pandey, S.; Ali, M. and Raisuddin, S. (2006): Biomarkers of oxidative stress in *Wallago attu* (Bl. and Sch.) during and after a fish-kill episode at Panipat, India. *Science total Environmental*, 368: 627–636.

Radwan EH (2016). Determination of total hydrocarbon and its relation to amino acids found in two bivalve edible species from Alexandria and El Ismailia coast, Egypt. *J Advances in biology*. Vol 9, No 5pp 1834-1844.

Radwan EH, Abdel Wahab WM, Radwan KhH (2012). Ecological and physiological studies on *Pinctada radiata* (Leach, 1814) collected from Alexandria coastal water (Mediterranean sea, Egypt. *Egypt J Exp Biol (Zool)*. 8(2) pp 223-231 .

Radwan EH, Fahmy GH, Saber MKh and Saber MKh (2017). The impact of some organic and inorganic pollutants on fresh water (Rashid branch, River Nile), Egypt. *J of advanced in biology*. Vol. 10, No. 2. Pp 2133-2145.

Radwan EH, Hamed ShSh, and Saad GA (2014). Temporal and spatial effects on some physiological parameters of the bivalve *Lithiphaga lithophaga* (Linnaeus, 1758) from coastal regions of Alexandria, Egypt. *Open J of ecology*. 4, 732-743.

Radwan EH, Hassan AA, Fahmy GH, El Shewemi SS, and Salam Sh (2018). Impact of environmental pollutants and parasites on the ultrastructure of the Nile boliti, *Oreochromis auruus*. *J of Biosciences and applied Research*, Vol 4, No 1 pp 58-83.

Radwan EH, Saad GA and Hamed ShSh (2016). Ultrastructural study on the foot and the shell of the oyster *Pinctada radiata* (Leach, 1814), Bivalvia: Petridae). *J Bioscience and applied research*, Vol. 2 No, 4,pp 274-283.

Rao, G.S. (1996): Glutathionyl hydroquinone: a potent pro-oxidant and a possible toxic metabolite of benzene. *Toxicology*, 106(1–3): 49–54 .

Regoli, Francesco, et al. "A multidisciplinary weight of evidence approach for environmental risk assessment at the Costa Concordia wreck: integrative indices from Mussel Watch." *Marine environmental research* 96 (2014): 92-104.

Regoli, F.; Frenzilli, G.; Bochetti, R.; Annarumma, F.; Scarcelli, V.; Fattorini, D. and Nigro, M. (2004): Time-course variations of oxyradical metabolism, DNA integrity and lysosomal stability in mussels, *Mytilus galloprovincialis*, during a field translocation experiment. *Aquatic Toxicology*, 68: 167–178.

Rodri'guez-Ariza, A.; Rodri'guez-Ortega, M.J.; Marengo, J.L.; Amezcua, O.; Alhama, J. and Lo'pez-Barea, J. (2003): Uptake and clearance of PCB congeners in *Chamaelea gallina*: response of oxidative stress biomarkers. *Comparative Biochemistry & Physiology*, 134(C): 57–67.

Roesijadi G (1996) Metallothionein and its role in toxic metal regulation. *Comp Biochem Physiol C* 113: 117–123.

Sajwan, K.; Kumar, K.S.; Paramasivam, S.; Compton, S. and Richardson, J. (2008): Elemental status in sediment and American oyster collected from Savannah marsh/estuarine ecosystem: A preliminary assessment. *Environmental contamination and toxicology*, 54(2):245-58.

Salman, J.M.; Randall, A.H.; Ayad, M.J. and Almamoori, (2002): Seasonal variation of heavy metal in water and two species of Molluscs in Hilla River Iraq. *International Journal of Geology, Earth & Environmental Sciences*, 4 (2): 16-24.

Shreadah, M.; and Tayel, F. (1992): Impacts of industrial, sewage and agricultural effluents on Lake Edku and Abo-Quir Bay. Bull. Fac. Sci., Alex. Univ., 32(A), 103-155.

UNEP/FAO/IAEA/IOC (1984): Sampling of selected marine organisms and sample preparation for trace metal analysis: Reference methods for marine pollution studies. UNEP 1: 11–21.

Valavanidis, A.; Vlahogianni, T.; Dassenakis, M. and Scoullos, M. (2006): Molecular biomarkers of oxidative stress in aquatic organisms in relation to toxic Environmental pollutants. Ecotoxicology Environmental Safety, 64: 178–189.

Vranković, Jelena, and Marija Slavić. "Biomarker responses in *Corbicula fluminea* to the presence of dioxin-like polychlorinated biphenyls and seasonal changes." Ecological indicators 48 (2015): 99-106.

WHO (1989): Heavy metals environmental aspects. Environmental Health Criteria (85).

Wu, S.C. and Zhang, Y. (2010): Active DNA demethylation: many roads lead to Rome. Nature Reviews Molecular Cell Biology, 11(10):750-9.

Wu, S.S.R. (2002): Hypoxia: from molecular responses to ecosystem responses. Marine Environmental Bulletin, 45(1-2): 35–45.

Yap, C.K.; Ismail, A. and Tan, S.G. (2002): Background concentrations of Cd, Cu, Pb and Zn in the green-lipped mussel *Perna viridis* (Linnaeus) from Peninsular Malaysia. Marine pollution Bulletin, 46(8): 1043- 1048.

Yoshioka, T.; Kawada, K.; Shimada, T. (1979): Lipid peroxidation in maternal and cord blood and protective mechanisms against activated oxygen toxicity in the blood. Am J Obstet Gynecol.; 135: 372–376.

Table (5): Correlation coefficient of heavy metals in tissue, heavy metals in water and oxidative Stress biomarkers:

Heavy metals in water										Oxidative Stress Biomarkers			Summer /Abo Quir Bay			
Cd	Cu	Pb	Mn	Heavy Metals in tissue						T. P.	GPx	SOD	r	MDA	Oxidative Stress Biomarkers	
Zn	Cd	Cu	Pb	Mn	Zn	T. P.	GPx	SOD	p	SOD	r	SOD				GPx
-0.293	-0.240	-0.432	0.340	-0.019	0.633	-0.545	0.750	-0.150	-0.017	0.800			-0.373	0.283		
0.573	0.647	0.392	0.510	0.972	0.178	0.264	0.086	0.777	0.974	0.056	0.466	0.586	p			
0.414	0.734	0.353	-0.206	0.310	0.440	-0.357	0.531	-0.311	0.743	0.132	-0.222		r			
0.414	0.097	0.492	0.696	0.550	0.382	0.487	0.278	0.548	0.090	0.803	0.673		p			
-0.168	0.169	0.416	-0.465	-0.821*	-0.265	0.941*	0.100	0.491	-0.049	0.238			r			
0.750	0.749	0.412	0.353	0.045	0.612	0.005	0.850	0.322	0.927	0.650			p			
-0.366	-0.121	-0.144	0.185	-0.557	0.596	0.025	0.864*	0.249	-0.006				r			
0.476	0.820	0.786	0.726	0.251	0.212	0.962	0.026	0.634	0.990				p			
0.193	0.947*	0.826*	0.155	-0.014	0.502	-0.245	0.228	0.324					r			
0.714	0.004	0.043	0.770	0.979	0.310	0.640	0.665	0.531					p			
-0.417	0.264	0.628	0.464	-0.767	0.283	0.328	-0.026						r			
0.411	0.613	0.182	0.354	0.075	0.586	0.525	0.961						p			
0.100	0.155	-0.066	0.056	-0.232	0.721	-0.065							r			
0.850	0.770	0.901	0.916	0.658	0.106	0.902							p			
0.032	0.015	0.255	-0.512	-0.633	-0.416								r			
0.953	0.978	0.626	0.299	0.178	0.411								p			
0.171	0.286	0.149	0.669	-0.040									r			
0.746	0.583	0.778	0.146	0.939									p			
0.586	-0.061	-0.381	0.034										r			
0.221	0.908	0.456	0.949										p			
-0.118	-0.124	-0.015											r			
0.824	0.815	0.978											p			
0.037	0.889*												r			
0.944	0.018												p			
0.296													r			
0.568													p			
													r			
													p			

