



BioBacta



Journal of Bioscience and Applied Research

www.jbaar.org



The protective effects and ameliorative potency of the haemolymph from the Saudi scorpion *Androctonus crassicauda* against the oxidative stress induced by its crude venom: A pharmacological study

Nagwa N. Azzam

Biology Department, Faculty of Sciences and Arts in Tehamah (Girls Section), King Khaled University, KSA

Abstract:

Background: Envenomation from dangerous scorpions remains a horrible threat in many parts of the globe, especially the developed countries, reflecting a reliable cause of a lot of mortalities and morbidities for both children and adolescents; as the annual number of scorpion stings exceeds 1,200,000 resulting in approximately 3250 deaths. In Saudi Arabia, scorpionism constitutes an acute major medical problem with about 15,000 average stung, yearly. Saudi government has given high priority to the development of health care services to monitor scorpionism carefully, to manage its treatment protocols successfully, and to explain the pathophysiological effects of the venom. A wide variety to treat scorpionism was used, either singly or in combination. Mostly, treatment modalities include antivenom immunoglobulin (SAV) and chemical antidotes, with varying degrees of effectiveness and side effects, though the cons and the so expensive wholesale cost associated with SAV treatment. We are in bad need and warranty challenged to obtain safer, more effective and not economically burdensome bioactive antivenins. Fourteen medically important scorpion species belonging to *Buthidae* have been identified in KSA. Although *Androctonus crassicauda* (*A. c.*) is an endemic and highly venomous scorpion in KSA, few studies were dealing with it. **Objective:** To evaluate the possible involvement of cellular oxidative reactions of the crude venom of the Saudi scorpion *A. c.*, as the main cause of multiple organ dysfunctions, and how far scorpion haemolymph (SH) could be effective to protect and / or treat the envenomation-associated metabolic disorders. **Materials and methods:** Twenty four adult male albino mice (25-32 g) were randomly divided into four groups; six in each. The controls; injected subcutaneously (SC) with 0.01 ml NaCl 0.9% / kg BW, envenomed animals; receiving a single dose of crude venom of the Saudi scorpion *A. c.* (0.12 mg/kg BW; SC) and tested two hours post-injection, envenomed animals treated within 5 min. with a SC dose of 0.12 mg of SH / kg BW, and envenomed animals treated with the same dose and route with SH, two hours prior to scorpionism. Animals were dissected and different biochemical parameters and oxidative stress biomarkers were measured in serum, liver and brain tissues. **Results:** Scorpion envenomation was

accompanied with oxidative damage and hyperglycemia; which is causative for generation of additional reactive oxygen species, and its subsequent metabolic disorders. Also, SH was predicted to significantly protect from and reverse all the cytotoxic manifestations following envenomation. **Conclusion:** The cytotoxic effects of crude venom of the Saudi scorpion *A. crassicauda* could be attributed to generation of reactive oxygen species causing oxidative damage. The scorpion haemolymph can be used as a bioactive therapeutic agent to protect and treat the dysfunctions subsequent to envenomation, as it may contain novel molecule (s) to do this.

Key Words: Scorpionism, *Androctonus crassicauda*, Scorpion venom, Scorpion haemolymph, antivenins, Oxidative stress, antioxidants

1. Introduction:

The scorpion (Order: *Scorpiones*) is one of the oldest known creatures existed as the most ancient and species rich terrestrial animals on the earth (Ding et al., 2014 & Al Asmari and Khan, 2016). Order *Scorpiones* is a distinctive group of class *Arachnida*; representing only 1.5 % of its members and including its oldest fossils, within the phylum *Arthropoda*, and is dating back to the Silurian period (almost 433–438 mya) (Dunlop and Selden, 2013; Waddington et al., 2015; Santibáñez-López et al., 2016 & Hosseini et al., 2017). Scorpions are a largely nocturnal secretive group of animals widely distributed worldwide; in tropical, subtropical and desert habitats; generally south of 45° N latitude with increased diversity in tropical regions at latitudes between 23 and 38 degrees (Polis, 1990 & Bowles and Swaby, 2006). Although most scorpions are not aggressive and stinging incidences usually occur accidentally (Bowles and Swaby, 2006), they have developed a negative reputation due to their stings and envenomation, where they pose serious health hazard due to their poisonous venoms (Uawonggul et al., 2006; Al Asmari et al., 2012; 2015; 2016 & 2017) in many underdeveloped regions (Kshirsagar et al., 2012 & Khatony et al., 2015).

One undeniable fact is that all scorpions are venomous and scorpion venom is the cause of

widespread fear in the globe (Frembgen, 2004 & Morrow, 2017), especially in the developing countries. Fortunately, only very few are dangerous to humans (Mansour, 2001 & Alkahlout, et al., 2015), where out of 1500 scorpion species about 50 are medically important. *Buthidae* is not only the largest family among scorpion lineages, but is the one that includes all the medically significant and the most potentially lethal species primarily distributed in Africa and Southeast Asia, as well (Goyffon and El Ayebe, 2002; Soleglad et al., 2003; Prendini and Wheeler, 2005; Bosmans and Tytgat, 2007; Chippaux and Goyffon, 2008; Caliskan et al., 2013; Quintero-Hernández et al., 2013; Sunagar et al., 2013; Hmed et al., 2013 & Al Asmari et al., 2016). Members of this family show extreme potency of their venom and the relatively smaller pincers; particularly pronounced in the fat-tailed scorpions' genera; *Androctonus* and *Parabuthus*, so they heavily rely on their venom arsenal for predation (Sunagar et al., 2013).

Scorpionism is a medical, social and economic problem (Telmisany and Othman, 1993) that represents a real threat in many parts of the world (Jarrar and Al-Rowaily, 2008; Mohamad et al., 2014 & Selmane et al., 2017), including Saudi Arabia (Fatani et al., 2006; 2010 & Al Asmari et al., 2017). Bawaskar and Bawaskar (2012) and Jalalia and Rahim (2014) reported that stings by venomous

animals; especially scorpions, are a concern to health authorities in many Middle East countries, because of the severity, extent, and a wide range of clinical effects. Due to its scattered geographical distribution, the at-risk population exceeds 2.5 billion people (Al Asmari et al., 2008). Scorpion stings (Ss) are reported to occur by an average approximates 1.2 million/year, leading to more than 3250 deaths with an annual mortality rate of about 3%. Scorpions exceed snakes in their toxicity. For every person killed by a poisonous snake, 10 are killed by a poisonous scorpion (Chippaux and Goyffon, 2008; Bawaskar and Bawaskar, 2012; Quintero-Hernández et al., 2013; Yilmaz et al., 2013; Adi-Bessalem et al., 2014; Ding et al., 2014 & Ebrahimi et al., 2017). Thus scorpion sting, and the resulting envenomation, constitutes a real predicament for healthcare professionals (Freire-Maia et al., 1994 & Daoudi et al., 2017).

In Saudi Arabia, scorpions are widely scattered. The Saudi fauna comprises at least 28 species and subspecies belonging to *Buthidae*, *Scorpionidae* and *Hemiscorpiidae* families (Desouky and Al-Shammari, 2011 & Al Asmari et al., 2013). Different fatal species; as *Androctonus*, are well identified (Al Asmari et al., 2007, 2009a & 2009b). Scorpion stings are common in this country due to its climate, socioeconomic structure and social habits. An average exceeding 18,500 annual stings is recorded in the different regions of Saudi Arabia (Desouky and Al-Shammari, 2011). Ss cases are a frequent presentation to emergency departments (EDs) in many areas and encountered on a daily basis, in certain regions (Alkahlout et al., 2015). Young children and elderly people are particularly at high risk (Telmisany and Othman, 1993). The most vulnerable people include the shepherds who tend the livestock in the desert,

farmers, and the people who spend their weekends in the desert for recreation. Scorpion stings cause a wide range of conditions, from severe local skin reactions, as pain, swelling and hypertension; that can be treated with analgesics and/or antihistamines, to neurologic, respiratory, and cardiovascular collapse and other systemic manifestations/supportive care, and coma (Al Sadoon and Jarrar, 2003; Chippaux and Goyffon, 2010; Jalali et al., 2011; Azzam, 2012 & Yilmaz et al., 2013). The synergistic impact of cardiac, respiratory, autonomic and metabolic abnormalities in scorpion sting patients may lead to multisystem failure and death (Murthy et al., 1991; Yugandhar et al., 1999; Uawonggul et al., 2006; Bawaskar and Bawaskar, 2007 & Al-Asmari et al., 2017). The clinical picture and response to scorpion envenomation varies depending on many factors; as the scorpion species, scorpion size, number of stings, bulk of the venom glands, status of the venom canals of the telson, the amount of venom injected, the age of the offender, the site and depth of sting penetration, quantity of the injected venom, and the proportion of the venom reaching the circulatory system, victim's age, weight, general health physiology status, genetics and emotional conditions, the interval between envenomation and hospitalization, the anatomical location of the sting and the season (time of sting) (Krishnan et al., 2007; Boşnak et al., 2009; Karnad, 2009; Uluğ et al., 2012; Azzam, 2012; Kassiri et al., 2014; Khatony et al., 2015 & Ebrahimi et al., 2017). The symptoms of a sting start immediately within few minutes and usually progress to a maximum severity (death of victim) within 5-7 hours, depending on the scorpion species (Mebs, 2002; Zargan et al., 2003; Petricevich, 2010 & Azzam, 2012).

The crude venom of a scorpion, produced by the columnar cells of the venom gland (Al-Asmari et al., 2016), is a water-soluble antigenic and heterogeneous complex multi-component mixture (Bosnak et al., 2009) of mostly small molecular weight peptides and small proteins that have evolved for prey capture and/or defense (Dutertre and Lewis, 2010) and can impair the cell function by interfering with ion channels permeability of the excitable cell membranes (Allen, 1985). Scorpion venoms consist of hundreds of different pharmacologically active components including variable combinations of proteins; usually called toxins, of 1,000–9,000 Da in size (Omran, 2003; Cao et al., 2006; Dutertre and Lewis, 2010 & Zhu et al., 2011). Toxins of a scorpion species are almost similar in structural and functional characteristics (Cao et al., 2014). It includes insoluble, generally non-toxic, fraction and a soluble fraction containing toxic peptides that exhibit activity on ion channels, in addition to mucopolysaccharides, nucleotides, vasoactive amines (serotonin or histamine), protease inhibitors and enzymes (Gazarian et al., 2005; de la Vega et al., 2010). Additionally, scorpion venom constituents include water, lipids, mucosa, ions, low molecular weight peptides, metals, mucoproteins, hyaluronidase, phospholipase, biogenic amines, glycosaminoglycans, phosphodiesterases, cytokine-releasers, and relative low molecular mass molecules, like serotonins, histamines, tryptophans, polynucleotides, protease inhibitors and histamine-releasers, small organic molecules; as polyamines, amino acids, and neurotransmitters, salts, nontoxic proteins, bioactive amines and many unidentified substances (Mahadevan, 2000; Gwee et al., 2002; Dittrich et al., 2002; Inceoglu et al., 2003; Sollod et al., 2005; Gazarian et al., 2005; Olivera and Teichert, 2007; Fox and Serrano, 2008 & Fry et al.

2009), that exert acute toxicological effects in humans (Bawaskar and Bawaskar, 2012; Venancio et al., 2013; Ding et al., 2014 & Al Asmari et al., 2016). In general, the toxic activity of scorpion venom is predominantly attributed to the presence of peptides that disrupt Na⁺, K⁺, Ca²⁺ and Cl⁻ channels in neuronal cells (Possani et al., 2000 & Venancio et al., 2013). Apart from these peptides, several molecules that play a role in scorpion poisoning or that exhibit properties of biotechnological interest are also present in scorpion venoms (Wu et al., 2010; Zeng et al., 2012; Zhao et al., 2011 & Venancio et al., 2013). The main constituents of the venom are the neurotoxins; chemicals affecting the nervous system, ultimately killing or paralyzing their prey (Tan et al., 2006 & Francielle et al., 2015).

Androctonus crassicauda; one of the five most dangerous scorpions in the world (Al-Ramahi and Al-Hasnawi, 2002), inhabits KSA especially Tehamah, Al-Baha, Hail and Al-Riyadh regions (Jarrar and Al-Rowaily, 2008 & Al Asmari et al., 2013). The venom of this species is neurotoxin; which is fast acting and can be absorbed very quickly, because of its small molecular weight proteins, resulting in fatal or paralytic effects (Al-Ramahi and Al-Hasnawi, 2002; Ozkan and Filazi, 2004 & Wikipedia, 2018). They are especially acting on the nerves responsible for respiration, leading ultimately to death by respiratory failure (Wikipedia, 2018). The symptoms of the sting start immediately within few minutes after stinging and usually progress to a maximum severity within 5 hours (Mebs, 2002).

A wide variety of treatments to scorpion envenomation were used either singly or in combination. No one treatment or combination of treatments showed any clear superiority (Curry et

al., 1984 & Azzam, 2012). Historically, the main modes of therapy for scorpion stings were supportive treatment and immunotherapy in the form of antivenom (Rodrigo and Gnanathan, 2017). Scorpion sting treatment employs various strategies including the use of specific medicines such as antiserum, especially for patients with severe symptoms (Carmo et al., 2015). All of the used antivenoms are animal-derived antisera of equine origin. Although effectiveness in neutralizing scorpion venoms, these animal-derived antisera are draw backed due to the heterologous nature of their protein contents, which may elicit both early and late adverse reactions in human recipients (Chippaux, 2012 & Rodríguez, 2015). Additionally, only a subset of the antibody/antibodies fragments present in these antivenoms have a therapeutic value since the presence of non-toxic immunogens in the venoms used for immunization may elicit therapeutically irrelevant antibodies in the immunized animal (Pucca et al., 2011 & Laustsen et al., 2016). Also, as these antivenoms are derived from animal serum, individual differences in the immune responses of the production animals may give rise to batch-to-batch variation (Rodríguez-Rodríguez et al., 2016). Finally, due to the very minute amounts of venom that can be extracted from scorpions, production of antisera against Ss is dependent on a highly laborious venom collection process, where large numbers of scorpions are needed to be milked in order to procure enough venom for immunization (Meadows and Russell, 1970 & Laustsen et al., 2016). Thus making antivenom (AV) is a complicated and so prohibitively expensive process (Danny, 2015). Also, AV is generally in short supply in most parts of Africa and many parts of Asia (William, 2015 & Shrestha, 2017). These challenges warrant

technological innovation, not only to obtain safer, cheaper and more effective AVs with therapeutic impact (Azzam, 2012), but also to establish more sustainable productions processes that are independent of both venoms and animals (Rodríguez, 2015 & Laustsen et al., 2016) with evaluation on their costcompetitiveness (Laustsen et al., 2016).

The present work was to investigate the possible involvement of cellular oxidative reactions in envenomation-induced changes by the crude venom of the Saudi scorpion *Androctonus crassicauda*; which were estimated to be the main concerned metabolic disorders in the brain and liver tissues of the victim, and to evaluate how far the scorpion haemolymph could be effective in protecting and/or ameliorating the cytotoxic changes due to scorpionism in the examined tissues and thus in restoring the envenomation associated metabolic abnormalities to be more or less within the normal values.

2. Materials and Methods:

2.1. The Experimental Animals:

Approval of this study was obtained from the animal use and care committee at the (65cm × 25cm × 15 cm) home cages, with sawdust-covered floors, the standard conditions of controlled humidity (70-80 %), temperature (25 ± 5°C) and light (12 hrs. darkness/ light cycle) and the conventional animal facilities of the lab. Healthy male Swiss Webster albino mice, weighing 20-25g, and, were used. Each cage contains six animals. They had access to stable food (mice chow) and water ad-libitum.

2.2. Scorpions and Venom collection and preparation:

Forty of the *Androctonus crassicauda* (A.c.) scorpions were collected by using ultra violet (UV) lamps during the no-moon nights from the wild in

Tehamah (Assir region; KSA) by professional hunters, specifically recruited for this job. Scorpions were brought to laboratory in tightly closed plastic bottles. The scorpion house was designed so that each scorpion batch was housed in larger plastic containers, which were placed in rows on shelves (Al Asmari et al., 2007; 2009a & 2009b). Avoiding scorpion cannibalism, scorpions were singly kept in a separate plastic boxes (6×6×10 inch) (L×W×H) on top of a soil bed of about 2 inches in depth, above which proper size stones, mimicking their natural habitat. Lids of boxes were designed to provide proper aeration and water-sprayed periodically to maintain humidity. Scorpions were fed on mealworms (Al Asmari et al., 2009a), once weekly (de Roodt et al., 2010); as they do not require food frequently due to their low metabolic rate (Neubauer, 2013 & Yaqoob et al., 2016). Scorpion in the laboratory were stored for one month and water was given to them through small humid cotton rolls, soaked in water and supplied for each animal on a regular basis (Moghadam et al., 2013 & Yaqoob et al., 2016).

Venom was extracted two weeks after the SH collection, not on the day immediately after feeding and also no food is added until three days after the venom extraction (Whittemore et al., 1963 & Candido and Lucas, 2004), by the electrical stimulation method explained by Abdel-Rahman (Abdel-Rahman, 2013); as the major components of venom collected by this method are toxins (Bücherl and Buckley, 1971). The venom was dissolved with sterile double-distilled water and centrifuged at 15,000 rpm for 15 min at 4°C (Özkan and Filazi 2004). The precipitate was discarded and the supernatant was stored at -20 °C until use. When required, the venom was reconstituted by adding 0.9% sodium chloride solution (physiological saline

solution; PSS). It was used for cytotoxic assays at a dose equals 1/5 LD₅₀, subcutaneously (SC); to simulate the natural route of envenomation and all envenomed mice were sacrificed two hours post injection to ensure absorption and common cement of venom's deleterious actions (Fatani et al., 2010; Azzam, 2012 & Azzam et al., 2013).

2.3. Determination of Median Lethal Dose (LD₅₀):

The median lethal dose; MLD, of *A. crassicauda* crude venom was determined according to the method of Meier and Theakston (Meier and Theakston, 1986). Different known concentrations of the venom solution dissolved in a PSS were prepared. Ten weighted mice were treated subcutaneously (SC) with different doses (D) of scorpion venom; as the SC route provides the highest LD₅₀ value and is the most frequent route of accidental scorpion envenomation (Krifi et al., 1999 & Özkan et al., 2007). Accordingly, the approximate SC LD₅₀ of *A. crassicauda* crude venom was calculated to be 0.60 ± 0.05 mg/kg.

2.4. Haemolymph collection and preparation:

To collect the haemolymph, a scorpion was starved five days prior to the collection of haemolymph (Geethabali and Rajashekhar, 1988) and treated according to Azzam (2012). After resting and feeding, the animal can be used again (Geethabal and Rajashekhar, 1988), either for more haemolymph or for venom production. The haemolymph was centrifuged at 3000 r.p.m. for 20 min to avoid coagulation. The supernatant was lyophilized and stored at -20°C until use. The used dose equals 0.05 mg/kg dissolved in a PSS. Immediately before usage 1/5 LD₅₀ of *A. crassicauda* crude venom solution dissolved in saline were incubated 10 min. at 37°C with the SH solution at a ratio of 2:1 (SH:SV, v:v). In treated

animals in all groups, injection was applied 5 min. after envenomation and half an hour before dissection (Azzam, 2012 & Azzam et al., 2013).

2.5. Cytotoxic Assays:

2.5.1. Study Design:

The experimental animals were adapted to the lab. environment for one week, then they were randomly divided into the experimental groups (n=6). 24 male Swiss Webster albino mice, weighing 20-25g, were allocated into four main groups:

- 1- The control (C) group: given a SC injection of the PSS (0.9% NaCl); 0.01 ml / kg B.W.
- 2- Envenomed (E) animals; received a single SC dose of 1/5 LD₅₀ of the *A. crassicauda* crude venom (0.12 mg / kg B. W.) and sacrificed for cytotoxic assays two hours post-injection of crude venom.
- 3- Envenomed animals protected with SH (EHP); treated with SH two one week earlier before injection with a single SC dose of 0.05 mg of the *A. crassicauda* crude venom/ kg B. W. of the experimental animal, and the examined tissues were used for biochemical measurements two hours after envenomation, as discussed earlier.
- 4- Envenomed animals treated with SH (EHT); injected with SH after 5 min. of envenomation and then dissected after two hours for bioassays, as mentioned before.

At the end of each experimental period, animals were sacrificed by decapitation, using sharp razor blade. Blood and tissue samples were treated and prepared for the biochemical assays as explained by Azzam (2012).

The following biochemical parameters were measured in the control and treated groups, in serum, liver and brain tissues:

2.5.2. Serum Glucose (SG) Assay:

SG was estimated by using the kit obtained from Biodiagnostic Company for Laboratory Services, referenced to the method of Trinder (1969).

2.5.3. Oxidative Stress Biomarkers:

Protein carbonyl content (PCOC) was determined according to the method of Reznick and Packer (1994) and Levine et al. (1994) while nitric oxide (NO) concentration was investigated according to Green et al. (1982) method. The Biodiagnostic Company for Laboratory Services kit; applying the techniques of Satoh (1978) and Ohkawa et al. (1979) was used to determine the lipid peroxidation end product; malondialdehyde (MDA).

2.5.4. Enzymatic and Non-Enzymatic Antioxidants Assays:

Assayment of Cu/Zn superoxide dismutase (SOD) activity was as per the procedures of Winterbourne et al. (1974) and Maral et al. (1977). The reduced glutathione (GSH) was estimated using the method adopted by Beutler et al. (1963). According to the techniques used in the kits of Biodiagnostic Company for Laboratory Services all the following parameters were estimated; glutathione peroxidase (GPx) (Paglia and Valentine, 1967), glutathione reductase (GSH-R) (Goldberg and Spooner, 1983), glutathione-s-transferase (GST) (Habig et al., 1974), Liver catalase (CAT) (Aebi, 1974 & 1984) activities, hydrogen peroxide (H₂O₂) content (Fossati et al., 1980 & Aebi, 1984) and finally the total antioxidant capacity (TAC) (Koracevic et al., 2001).

2.6. Statistical Analysis:

Raw data were statistically analyzed by using SPSS software (Statistical Package for Social Science, version 17.01, Illinois, USA) (Dancey and

Reidy, 2002). Tabulation and graphics of data were done using Microsoft Excel 2007. According to the mathematical principles described by Field, all of the data of control and treated groups were expressed as mean value \pm standard error (Field, 2000). One-way analysis of variance (ANOVA) was carried out followed by *post hoc* Duncan's test (Bewick et al., 2004); for comparison between control and treated groups. The probability criterion for significance for each statistical test was $P \leq 0.05$.

3. Results:

The results of this work; presented in tables "1, 2 and 3", indicated that a single SC injection of 0.12 mg of the crude venom of *A.c.*/kg B.W. in the experimental animals causes marked significant changes in the biochemical parameters in the serum and tested organs of the envenomed group, in comparison with the control values, at $P \leq 0.05$. Serum glucose (SG) concentration, malondialdehyde content in brain and liver (bMDA & lMDA), brain and liver protein carbonyl contents (bPCOC & lPCOC), nitric oxide in serum and the examined tissues (sNO, bNO & lNO), glutathione-S-transferase activity in brain and liver tissues (bGST & lGST) were significantly elevated post venom treatment in comparison with the control values. Otherwise, significantly declined brain and liver reduced glutathione (bGSH & lGSH) level, activities of liver catalase (lCAT) and superoxide dismutase activity in serum, brain and liver (sSOD, bSOD & lSOD), brain and liver glutathione peroxidase (bGPx & lGPx) and tissues glutathione reductase (bGSH-R & lGSH-R) hydrogen peroxide content in serum, brain and liver (sH₂O₂, bH₂O₂ & lH₂O₂) were noticed with consequently sharp and significant decremented total antioxidant capacity (TAC) (tables "1"- "3" & figures "1"- "12").

Pre and post-treatment of the scorpion-envenomed animals with the desired dose of the SH as mentioned earlier; has been predicted to prevent and restore all the metabolic disturbances and significantly ameliorate them to be more or less within the normal values; as cleared from the investigated data.

Applying one-way ANOVA revealed that the difference between treated and control groups for the increased SG to be very highly significant ($F_{6,6} = 119.682$, $P \leq 0.05$). This was the same for differences in brain and liver MDA contents ($F_{6,6} = 685.314$ and 270.131 respectively, $P \leq 0.05$), brain and liver PCOC ($F_{6,6} = 71.081$ and 95.126 , respectively, $P \leq 0.05$), serum, brain and liver NO contents ($F_{6,6} = 211.423$, 424.192 and 327.090 , respectively, $P \leq 0.05$), the activity of GST in brain and liver tissues ($F_{6,6} = 281.312$ and 39.011 , respectively, $P \leq 0.05$) and H₂O₂ content in serum, brain and liver of the examined animals ($F_{6,6} = 96.050$, 182.015 and 212.075 , respectively, $P \leq 0.05$). Also, one-way ANOVA test between groups certified very highly significant differences on the decreases for the mean values for the GSH content in sera, brain and liver tissues of the experimental animals ($F_{6,6} = 139.411$, 143.228 and 143.016 , respectively, $P \leq 0.05$), lCAT ($F_{6,6} = 61.24$, $P \leq 0.05$), Cu/Zn SOD ($F_{6,6} = 82.124$, 127.039 and 114.615 for serum, brain and liver brain and liver, respectively, $P \leq 0.05$), GPx activities ($F_{6,6} = 45.307$, 20.276 , respectively, $P \leq 0.05$), GSH-R activity in brain and liver tissues ($F_{6,6} = 37.003$ and 31.954 , respectively, $P \leq 0.05$), and finally gave insurance for very highly significant differences between groups in their venom-evoked decreases in sera, brain and liver TAC concentrations ($F_{6,6} = 75.448$, 187.123 and 182.213 , respectively, $P \leq 0.05$).

Table (1): Effects of the Experimental Conditions on Serum Glucose and Oxidative Stress Biomarkers

Group	SG Mg / 100 ml	MDA mg / g wet tissue		PCOC mM / g wet tissue		NO mM / g wet tissue		
		bMDA	lMDA	bPCC	lPCC	sNO	bNO	INO
C	91.37 ± 6.97	12.40 ± 1.30	67.65 ± 6.68	5.35 ± 0.76	4.56 ± 1.02	8.72 ± 0.98	12.48 ± 0.91	12.40 ± 1.30
E	393.79 ± 15.31*	31.23 ± 2.29*	326.17 ± 9.23*	32.39 ± 0.94*	33.85 ± 0.79*	39.22 ± 1.39*	33.53 ± 1.00*	31.23 ± 2.29*
SHP	92.92 ± 5.99¥	12.69 ± 1.02¥	67.75 ± 5.07¥	5.39 ± 1.01¥	4.60 ± 1.12¥	8.85 ± 1.04¥	12.49 ± 1.22¥	12.69 ± 1.02¥
SHT	95.86 ± 6.66¥	12.64 ± 1.99¥	68.91 ± 6.17¥	6.02 ± 1.17¥	4.99 ± 0.97¥	8.12 ± 1.12¥	12.79 ± 1.65¥	12.64 ± 1.99¥

Table (2): Effects of the Experimental Conditions on the Glutathione Circuits

Group	GSH mg/100 ml & mg/g wet tissue			GPx U/g wet tissue		GSH-R U/g wet tissue		GST U/g wet tissue	
	sGSH	bGSH	lGSH	bGPx	lGPx	bGSH-R	lGSH-R	bGST	lGST
C	2.87 ± 0.13	4.47 ± 0.11	3.15 ± 0.06	48.88 ± 1.14	41.47 ± 2.08	140.23 ± 5.95	52.01 ± 2.99	49.89 ± 6.06	88.93 ± 4.92
E	0.39 ± 0.09*	0.61 ± 0.12*	0.49 ± 0.06*	14.06 ± 2.51*	11.87 ± 1.90*	43.02 ± 1.97*	15.98 ± 3.36	231.19 ± 9.97*	300.02 ± 7.34*
SHP	2.88 ± 0.23¥	4.46 ± 0.19¥	3.19 ± 0.25¥	48.84 ± 1.18¥	41.05 ± 1.46¥	140.63 ± 4.80¥	52.29 ± 3.96¥	49.66 ± 8.46¥	89.01 ± 6.07¥
SHT	2.77 ± 0.31¥	4.11 ± 0.07¥	2.99 ± 0.29¥	48.03 ± 1.16¥	41.01 ± 0.97¥	139.63 ± 4.02¥	51.79 ± 3.08¥	50.71 ± 4.97¥	91.98 ± 4.98¥

(*) Represents significant difference between the control and envenomed groups at $P \leq 0.05$.
 (¥) Represents significant difference between the envenomed and treated groups at $P \leq 0.05$.

Table (3): Effects of the Experimental Conditions on the Enzymatic and Non-Enzymatic Antioxidants

Group	ICAT U/g wet tissue	H ₂ O ₂ mg / 100 ml & mg / g wet tissue			SOD mg / g wet tissue			TAC mM / L & mM / g tissue		
		sH ₂ O ₂	bH ₂ O ₂	lH ₂ O ₂	sSOD	bSOD	lSOD	sTAC	bTAC	lTAC
C	40.01 ± 0.31	124.68 ± 2.97	171.44 ± 3.90	162.16 ± 6.40	28.24 ± 1.096	6.89 ± 0.87	34.25 ± 0.94	3.95 ± 0.52	19.14 ± 1.05	19.83 ± 0.99
E	16.89 ± 0.16*	250.10 ± 3.864*	403.11 ± 12.44*	385.22 ± 9.87*	10.94 ± 0.57*	3.03 ± 0.28*	19.05 ± 1.14*	1.01 ± 0.75*	6.96 ± 0.81*	5.04 ± 0.99*
SHP	39.95 ± 1.55¥	124.89 ± 6.16¥	171.64 ± 6.13¥	161.95 ± 8.08¥	28.43 ± 0.71¥	6.28 ± 1.09¥	34.00 ± 0.79¥	3.94 ± 0.41¥	19.04 ± 1.88¥	19.86 ± 0.63¥
SHT	38.97 ± 1.92¥	126.58 ± 8.88¥	174.80 ± 5.33¥	163.61 ± 5.89¥	27.01 ± 0.77¥	6.00 ± 0.90¥	33.15 ± 0.95¥	3.19 ± 0.33¥	18.05 ± 0.71¥	18.98 ± 0.96¥

(*) Represents significant difference between the control and envenomed groups at $P \leq 0.05$.
 (¥) Represents significant difference between the envenomed and treated groups at $P \leq 0.05$.




Fig. (1): Serum Glucose Content




Fig. (2): Tissue Malondialdehyde (MDA) Content




Fig. (3): Tissue Protein Carbonyl Content (PCOC)




Fig. (4): Serum and Tissue Nitric Oxide (NO) Content




Fig. (5): Serum and Tissue Reduced Glutathione (GSH) Content




Fig. (6): Tissue Glutathione Peroxidase (GPx) Activity




Fig. (7): Tissue Glutathione Reductase (GSH-R) Activity




Fig. (8): Tissue Glutathione-S-Transferase (GST) Activity




Fig. (9): Liver Catalase (CAT) Activity




Fig. (10): Serum and Tissue Superoxide Dismutase (SOD) Activity




Fig. (11): Serum and Tissue Hydrogen Peroxide (H₂O₂) Content




Fig. (12): Serum and Tissue Total Antioxidant Capacity (TAC)

Discussion:

Although the mechanism of the hyperglycemia induced by scorpions' venom; indicated as one of the pathophysiological effects of scorpion envenomation in this work (table "1" & fig."1"), and are in line with other investigators (Abd Rabo, 2006; Choudhry et al., 2011; Azzam, 2012; Azzam et al., 2013; de Oliveira et al., 2016 & Chaubey, 2017), is not clearly understood, there are some suggestions. This can occur through the reduced insulin secretion, excessive release of catecholamines, decreased thyroid hormone levels

and increased cortisol and glucagon levels (Murthy and Zare, 1998; Goncalves et al., 2003; Oliveira et al., 2016 & Chaubey, 2017). This appears to be secondary to cause an autonomic storm that activates both sympathetic and parasympathetic autonomic nervous system (Mirakabadi et al., 2006), where envenomation is a powerful stimulant of the autonomic nervous system, especially the peripheral and central adrenergic systems, causing intense autonomic discharge and leading to a massive release of neurotransmitters (Murthy and Haghazari, 1999; Dittrich et al., 2002; Vasconcelos

et al., 2004; Azzam, 2012 & Azzam et al., 2013) and activation of b-receptors with catecholamine and serotonin secretion (Adam, 1959), which give rise to several clinical manifestations including hyperglycemia (Murthy and Haghazari, 1999; Dittrich et al., 2002; Vasconcelos et al., 2004; Mirakabadi et al., 2006; Azzam, 2012; Azzam et al., 2013 & Murthy et al., 2015). The released catecholamines results in a variety of metabolic effects and mobilizes substrates from fuel depots; as mobilization and break down of liver glycogen causing an increase in blood sugar levels and activation, via cyclic AMP, of hepatic glycogen phosphorylase, the enzyme responsible for converting glycogen to glucose-1-phosphate, the rate limiting step in glycogenolysis (Ganong, 1987; Keele et al., 2000; Jackson, 2006 & Murthy et al., 2015). Also, the epinephrine-like effects, mediated by the serotonin, present in the scorpion venom, was said to be responsible for the predicted hyperglycemia and liver and muscle glycogenolysis after envenomation (Mohamed et al., 1972) where it acts as a stimulant of gluconeogenesis (Murthy, 2000) and inhibits glycolysis (Murthy et al., 2003). Otherwise envenomation was estimated to enhance glucose production by a combined epinephrine, glucagon, and cortisol infusion that occurs as a result of an additive effect on hepatic glucose production, without any additional change in glucose clearance (Goldstein et al., 1995). Additionally, the suppressed insulin secretion, block of insulin secretion (El-Asmar et al., 1974; Mirakabadi et al., 2006 & Murthy et al., 2015), and insulin resistance (Zare, 1993) by scorpion envenomation may be causative. Furthermore, the actions of the scorpion neurotoxins on voltage-gated sodium (Nav) or potassium (Kv) channels have demonstrated a gamut of physiological responses,

including the increasing of plasma glucose levels (Vasconcelos et al., 2004; Ribeiro et al., 2010; Pucca et al., 2012; 2015 & Oliveira, et al., 2016). Goncalves and his team work clarified a direct effect of a scorpion toxin on isolated islets of Langerhans (not provided by catecholamine's action), enhancing the β -cells membrane depolarization and significantly potentiating glucose-induced insulin secretion (Goncalves et al., 2003 & Oliveira et al., 2016). This is in consistent with imitating OS to be the major causative factor of venom-induced toxicity and has been associated with renal failure, hepatic impairment and acute pancreatitis in viper and other envenomed experimental animals and humans.

Of great importance, the significant increases of the oxidative stress (OS) biomarkers in the envenomed animals, revealed by our results (table "1" & figures "2" -"4") and previously by others (De Roodt et al., 2003; Meki et al., 2003; Hammoudi-Trik et al., 2004; Fatani et al., 2006; 2010; Azzam, 2012; Azzam et al., 2013; Al Asmari et al., 2016 & Al-Sheikh et al., 2017), indicating that scorpion venom could induce oxidative damage in the envenomed animals. Besides causing many human pathologies (Valko et al., 2007) (Yamasaki et al., 2008; Valenta et al., 2010; Sagheb et al., 2011; Al Asmari et al., 2014; 2016 & Al-Sheikh et al., 2017).

Oxidative stress is a cellular phenomenon or condition simply defined as a case of physiological imbalance between the levels of antioxidants and oxidants (free radicals; FRs or reactive species; RS) (Boveris and Navarro, 2008; Kalam et al., 2012 & Moustafa, 2015) in favor of oxidants (Ighodaro and Akinloye, 2017). OS occurs as a deleterious process implicated in the development of many diverse diseases including hypertension, cardiac dysrhythmia and myocardial damage (Satoh and

Nishida, 2004; Fatani et al., 2006 & Liang and Juan, 2011), all of which are present in scorpion envenomation (Ismail, 1995; Amitai, 1998; de Roodt et al., 2003; Hammoudi-Triki et al., 2004; Dousset et al., 2005; Fatani et al., 2006; Liang and Juan, 2011 & Aksoy et al., 2012). An oxidant/free radical is defined as any chemical species containing one or more unpaired electrons in the outer atomic or molecular orbital/s (Kohen and Nyska, 2002; Pryor et al., 2006; Aksoy et al., 2012 & Aprioku, 2013). Because of these unpaired electrons, free radicals are highly reactive and readily take part in chemical reactions with virtually all cell components (lipids, proteins, complex carbohydrates and nucleic acids) in the body. These reactions occur through a chain of oxidative reactions to cause tissue injury (Kohen and Nyska, 2002 & Aprioku, 2013). FRs can be formed from two groups of sources; endogens, mostly mitochondrial electron transport chains, and exogens; as cigarettes, pesticides, solvents, petrochemical products, drugs, alcohol, solar rays, stress, x-rays and heavy physical activity (Valko et al., 2007 & Aksoy et al., 2012). Although they have a very short life cycle, FRs may be harmful to organisms due to their activity levels (Braunersreuther and Jaquet, 2012). The FRs of interest, in most biological systems are referred as reactive oxygen species (ROS); where the most significant free radicals are oxygen-centered (Aprioku, 2013). ROS in the cells include singlet oxygen ($^1\text{O}_2$), superoxide anion ($\text{O}_2^{\bullet-}$), hydroxyl radical ($^{\bullet}\text{OH}$), hypochlorous acid (HOCl) and hydrogen peroxide (H_2O_2) (Slater, 1984; Halliwell, 2006; Pryor et al., 2006 & Aprioku, 2013). Although H_2O_2 contains no unpaired electrons, it is considered a reactive oxygen species because of its ability to generate highly reactive hydroxyl free

radicals through interactions with reactive transition metals like iron and copper (Slater, 1984; Aruoma et al., 1991 & Valko et al., 2005). Reactive nitrogen species (RNS) is another group of free radicals referring collectively to nitrogen-centered radicals that alters cellular functions. RNS includes nitric acid (NO), peroxyxynitrate radical (ONOO^{\bullet}), nitrogen dioxide radical ($^{\bullet}\text{NO}_2$) and other oxides of nitrogen and products arising when NO reacts with $\text{O}_2^{\bullet-}$, peroxy radical (RO^{\bullet}) and alkoxy radical (ROO^{\bullet}) (Kohen and Nyska, 2002; Victor et al., 2004; Valko et al., 2005 & Aprioku, 2013). ROS and RNS are generated as a consequence of normal cellular metabolism (Lobo et al., 2010), but increased under conditions of both physiological and psychological stress (Jackson, 2000) as in scorpionism (Azzam, 2012 & Azzam et al., 2013). They are well recognized to play a dual role; both deleterious and beneficial species (Valko et al., 2007). The human body has antioxidative mechanisms of two groups; enzymatic (SOD, CAT, GPx, GST, GSH-R, etc.) and nonenzymatic (GSH, ascorbic acid; Vitamin C, α -tocopherol; Vitamin E, carotenoids, flavonoids, melatonin, albumin, uric acid, etc) to overwhelm oxidants (Al-Gayyar et al., 2007; Valko et al., 2007; Gill et al., 2011; Zengin et al., 2014; Thatoi et al., 2014 & Moustafa, 2015). Total antioxidant capacity (TAC) of an organism represents the sum of all antioxidant activities in the organism, including both enzymatic and non-enzymatic activities of antioxidants, that can be regarded as the ability of an organism to suppress ROS (Lesser, 2006 & Dann, 2017).

OS can occur due to an overproduction of oxidants, over exposure to exogenous oxidants, a decrease in antioxidant defenses or a combination of these factors (Zengin et al., 2014). On this context, the scorpion-induced cytotoxic oxidative damage

was shown to cause lipid and protein oxidation to the experimental animals as compared with the controls, indicated by significant increases in MAD, NO and PCOC levels in brain and liver tissues (table "1"). It is well known that scorpion venoms, by their action mainly on sodium channels, enhance the release of various neurotransmitters, such as adrenaline and noradrenaline, (Ismail, 1995; Meki et al., 2003 & Azzam, 2012) that can induce the generation of FRs leading to OS and apoptosis (Khorchid et al., 2002).

The mechanisms by which FRs and lipid peroxidation (LPO) are generated in scorpion envenomation are complex and may depend on a multitude of interacting factors (Fatani et al., 2006 & Azzam, 2012); 1) Scorpion venoms contain neurotoxins that act on a number of ionic channels resulting in alteration of ionic transport and cytosolic calcium overload (Harvey et al., 1992; Fatani et al., 2006 & Azzam, 2012), which 2) elevates adenosine monophosphate (AMP) concentration and increases its catabolism, leading subsequently to generation of FRs. 3) These FRs attack membrane phospholipids causing their peroxidation, that 4) eventually causes impairment of biological membrane functioning, decreased fluidity, and inactivation of membrane-bound receptors, all of which may culminate in multiple organ dysfunction (MOD) (Bhaumik et al., 1995; Love, 1999; Al-Omar et al., 2004; Fatani et al., 2006 & Azzam, 2012); a feature encountered in scorpion envenomation (Meki et al., 2003 & Azzam, 2012).

LPO is a natural metabolic process under normal aerobic conditions and it is one of the most investigated consequences of ROS action on membrane structure and function (Blokhina et al., 2003). Generally, LPO can be described as a process under which oxidants such as FRs or non-radical

species attack lipids containing C = C bond/s; especially polyunsaturated fatty acids (PUFAs) that involve hydrogen abstraction from a carbon, with oxygen insertion, resulting in lipid peroxy radicals and hydroperoxides (Yin et al., 2011) causing its deterioration (Devasagayam et al., 2003). Also, the presence of phospholipases (PLA), as a constituent in the toxins produced by venomous animals (De Maria et al., 2007), including scorpions (Zamudio et al., 1997; Conde et al., 1999; Valdez-Cruz et al., 2004; Hariprasad et al., 2007; Hmed et al., 2013 & Incamnoi et al., 2013) may be causative. PLA is a group of enzymes that hydrolyze one of the ester bonds on phospholipids to release a variety of products (Shimizu et al., 2006), as arachidonic acid whose metabolism results in the formation of potentially toxic ROS and lipid peroxides (Adibhatla and Hatcher, 2008 & Abou-Elezz et al., 2017). Currently, LPO is considered as the main molecular mechanisms involved in the oxidative damage to cell structures and in the toxicity process that lead to cell death (Repetto et al., 2012). Peroxidation process of membrane lipids include loss of polyunsaturated fatty acids (PUFAs), decreased lipid fluidity, altered membrane permeability, effects on membrane associated enzymes, altered ion transport, release of material from subcellular compartments and the generation of cytotoxic metabolites or lipid hydro-peroxides (Santamaria et al., 2002 & Abou-Elezz et al., 2017). LPO produces a wide variety of oxidation products (Ayala et al., 2014). MDA which is one of the aldehydes formed as secondary end-product product during LPO as is a final metabolite of LPO formed from a variety of unsaturated fatty acids in biological membranes stimulated by ROS and RNS overproduction (Valko et al., 2007), and generated by decomposition of unsaturated fatty acids

(arachidonic acid and larger PUFAs) (Esterbauer et al., 1991; Valko et al., 2007; Yin et al., 2011 & Ayala et al., 2014), through enzymatic or non-enzymatic processes, appears to be the most mutagenic product of LPO (Esterbauer et al., 1990), widely used as a convenient biomarker for LPO (Ayala et al., 2014).

Additionally, as predicted from our results, the single SC dose of 1/5 LD₅₀ (0.12 mg / kg BW) of the crude venom of Saudi scorpion *Androctonus crassicauda*, induced evidenced protein oxidation, as indicated by accumulation of protein carbonyl (PCO) in the examined tissues (table "1" & fig. "3"). Previous studies agreed with these data (Krishnan and Kodrik, 2006; Abdel-Rahman et al., 2009; Azzam, 2012; Azzam et al., 2013 & Béchohra et al., 2016). Proteins are possibly the most immediate vehicle for inflicting oxidative damage on cells, because they are often used as catalysts (Pandey and Rizvi, 2010). Exposure of proteins to reactive oxygen species (ROS) alter the physical and chemical structure of the proteins causing oxidation of side-chain groups, protein scission, backbone fragmentation, cross-linking and unfolding. These alterations are consequent to formation of new reactive groups, generation of protein carbonyls (PCOs), oxidation of -SH groups, formation of dityrosine containing cross-linked protein products known as protein products (AOPPs) and many others (Levine et al., 1990; Witko-Sarsat et al., 1996; Cakatay et al., 2008 & Pandey and Rizvi, 2010). Protein carbonyl content (PCOC) is actually the most general indicator and the most commonly used marker of protein oxidation (Berlett and Stadtman, 1997; Shacter, 2000; Chevion et al., 2000; Beal, 2002 & Deori et al., 2016), particularly associated with OS derived from free radicals and other activated oxygen species (Levine et al., 1990 &

Halliwell and Whiteman, 2004). Protein carbonyls are formed early during OS conditions and are not a result of one specific oxidant, thus they can be called a marker of overall protein oxidation (Weber et al., 2015). PCO is a product of irreversible non-enzymatic oxidation or carbonylation of proteins (Dalle-Donne et al., 2006) that represents an irreversible form of protein modification, demonstrated to be relatively stable (degradation/clearance in hours/days) in contrast to LPO products that are removed within minutes (Grune et al., 1995 & 1996). The usage of PCO group as a biomarker of OS has some advantages in comparison with the measurement of other oxidation products because of the relative early formation and the relative stability of carbonylated protein (Berlett and Stadtman, 1997 & Dalle-Donne et al., 2003).

Carbonyl (C=O; CO) is a functional group consisting of a carbon atom that has a double bond to an oxygen atom, thus aldehydes, ketones, carboxylic acids, esters and amides are all carbonyl groups (Danielsen, 2015). The carbonyl groups can be introduced on protein side chains, when they are oxidized, by a variety of oxidative pathways (Dalle-Donne et al., 2003; Fedorova et al., 2014 & Abou-Elezz et al., 2017). Oxidation of certain amino acids within proteins can result in formation of PCO which are disturbers of protein function (Dalle-Donne et al., 2003 & Danielsen, 2015), either by modifying their structural function or by causing loss of catalytic activity. Oxidation can arise from direct oxidation of most protein residues or with secondary oxidation products (Beal, 2002). Oxidative cleavage of proteins; by either the α -amidation pathway or by oxidation of glutamyl side chains, leading to formation of a peptide in which the N-terminal amino acid is blocked by an α -

ketoacyl derivative, is a way (Berlett and Stadtman, 1997 & Dalle-Donne et al., 2003). In addition, CO groups may be introduced into proteins by secondary reaction of the nucleophilic side chains of Cys, His, and Lys residues, with the aldehydes produced during LPO, or with reactive carbonyl derivatives generated as a consequence of the reaction of reducing sugars, or their oxidation products with lysine residues of proteins with the eventual formation of the advanced glycation / lipoxidation end products (AGEs/ALEs) (Berlett and Stadtman, 1997; Stadtman and Berlett, 1997; 1998; Uchida, 2000; Dalle-Donne et al., 2003 & Halliwell and Gutteridge, 2015). The venom-induced oxidative stress which elevates the steady level of ROS causes protein carbonyls to accumulate on the side chains of proteins, decreases the TAC, also indicated in this work; table "3" and fig. "12", (Chaudhuri et al., 2006; Krishnan and Kodrik, 2006; Abdel-Rahman et al., 2009; 2010, Azzam, 2012 & Azzam et al., 2013) and degrades the ability to oxidized proteins by either decreased proteases concentration or increased levels of proteases inhibitors (Stadtman, 2004).

On contrary, there is a relation between the observed increase in nitric oxide (NO) concentration after envenomation and the resulted elevations in PCOC; where the increase in PCOC might be in part due to the increase in the NO concentration. NO; a free radical generated as a byproduct of a five-electron oxidation of the amino acid L-arginine mediated by one of three isoforms of nitric oxide synthase (Moncada et al., 1991; Annane et al., 2000 & Abdoon, 2004), readily reacts with superoxide ($O_2^{\bullet-}$) to form peroxynitrite ($ONOO^-$), a potent oxidant and nitrating agent capable of attacking and modifying proteins and depleting antioxidant

defenses (Mc Cord, 2000 & Abdel-Rahman et al., 2010).

On this respect, our work results (table "1" & fig. "4") were in harmony with the previous works (Dousset et al., 2005; Petricevich, 2010; Abdel-Rahman et al., 2010; Azzam, 2012; Sahan-Firat et al., 2012; Azzam et al., 2013; & Ahmed et al., 2015) denoting that scorpion envenomation causes elevated NO content. Within a cell, the effects of NO are concentration dependent and its actions are dependent on its levels, the redox status of the cell and the amount of metals, proteins, and thiols, amongst other factors (Rosselli et al., 1998 & Agarwal et al., 2012). NO is known to have a dual role; as a ubiquitous intracellular messenger in many mammalian systems and a contributor to lethal processes (Ignarro, 2000), and the cardiovascular autonomic nervous system has proven to be no exception (Chowdhary and Townend, 1999). NO as one of the smallest and most known diffusible signal molecules, can freely crosses membrane barriers to exert its effects through direct binding and/or reaction with target proteins (Stamler et al., 1992 & Stanley, 2000). It is known to be involved in multiple reactions (Stamler et al., 1992) of biological importance; such as being an endogenous regulatory mediator in the control of vascular tone (Moncada et al., 1991) and vasomotor function (Petricevich, 2004; 2010; Ahmed et al., 2015 & Abou-Elezz et al., 2017), its implication in the genesis of various pathological conditions such as inflammation, arteriosclerosis, hypertensive vascular lesions in which the infiltration of inflammatory cells occurs (Olsen, 1972; B'usse and M'ulsch, 1990; Libby and Hansson, 1991) and insulin resistance, as which resulted from envenomation (Petricevich, 2002; Petricevich and Pe'Na, 2002 & Azzam, 2012). The high levels of

NO in the venom-treated animals may abolish the effect of catecholamines, as a reflex control of sympathetic nerve activity (Chowdhary and Townend, 1999; Abdel-Rahman, 2009; Azzam, 2012 & Azzam et al., 2013). Several inducers of insulin resistance; including pro-inflammatory cytokines and oxidative stress that can modify the levels of nitrites, and nitrate production have been associated with several conditions as envenomation. Envenomation by scorpions, bees, snakes, spiders and wasps involves release and activation of pro-inflammatory cytokines that may be at least partially responsible for NO synthesis (Petricevich, 2004; 2010 & Azzam, 2012). NO could be released by tissues after scorpionism via induction of the inducible nitric oxide synthase (iNOS) (Stamler et al., 1992) and production of ROS (Meki and Mohey El-Deen, 1988). Also, the significantly high serum level of NO supports the evidence for the role of these potent vasodilators in the terminal hypotension that is usually observed in humans and animals after envenomation (Abdoon and Fatani, 2009). In our findings, NO secretion may be increased for its prominent protective role against the neurotoxic effects of the scorpion toxins, either by direct activation of Ca^{2+} -activated- K^+ -channel (K^+ -Ca) in cell-free membrane patches (Bolotina et al., 1994 & Azzam, 2012) or by inhibiting the mitochondrial Ca^{2+} uptake and increasing the probability for opening the mitochondrial ATP-sensitive K^+ -channel by modulating affinity to ATP (Rakhit et al., 2001 & Azzam, 2012). On the other hand, NO can give rise to toxic secondary radical species (Stamler et al., 1992).

Naturally, cells have developed an extensive defense system (antioxidants) to neutralize the harmful oxidative effects of excessive ROS generation (Volko et al., 2007). Antioxidant enzyme

activity and oxidant generation rates have been postulated to be useful in estimating the level of oxidative damage (Volko et al., 2007 & Aboul-Soud et al., 2011). Antioxidants are important inhibitors of LPO, not only for protection but also as a defence mechanism of living cells against oxidative damage (Dauqan et al., 2011 & Padmanabhan et al., 2012). Apart from being involved in damaging cellular components, ROS seem to play a major role in venom-induced toxicity (Dousset et al., 2005). This is evident from the present results; as the fine balance, normally maintained between antioxidant activity and oxidant generation, has shifted in favor of oxidative reactions, causing OS in the E group (tables "1" - "3" & figures "5" - "12"). Cellular antioxidant defense systems including SOD, CAT and glutathione redox cycle; which reduce ROS to water (Volko et al., 2007; Lee et al., 2015 & Patlevič et al., 2016; Ighodaro and Akinloye, 2017) are the most important organic mechanisms against tissue damage induced by FRs (Kochar and Umathe, 2009 & Azzam, 2012), that may prevent disturbances in ROS homeostasis, or reduce the effect of oxidative stress in cells.

GSH is the first essential defense mechanism for the maintenance of homeostasis and redox balance and the prevention of LPO and depletion of cellular thiols (Halliwell and Gutteridge, 1989 & Couto et al., 2013). It is considered the most abundant intracellular ubiquitous thiol-based antioxidant involved in numerous processes essential for normal biological functions, as DNA and protein synthesis (Peña et al., 2003 & Azzam, 2012). In the GSH redox cycle, GPx catalyzes the decomposition of H_2O_2 and lipid peroxides to water. The present results (table "2" & fig."5") indicated very significant decreases in GSH, thus producing a pronounced OS. Furthermore, there was a very

significant reduction in GSH-R activity (Table "2" & fig. "7"). The drop in GSH levels of the venom-treated animals may be caused by the pronounced decrease in GSH-R activity or as a result of GSH reacting directly with excessively generated H_2O_2 . Valko and his co-workers reported that GSH levels can be regulated by its denovo synthesis catalyzed by γ -glutamate cysteine ligase and glutathione synthase, hence, the noted drop in GSH levels of examined tissues of the E group may be a result of the crude venom acting to lower the gene expression levels and consequently the activities of these enzymes (Valko et al., 2007 & Ghneim, 2017).

Remarkable decline in liver CAT activity was noticed in parallel with the decrements in GSH in the tested organs of envenomed animals (table "3" and fig. "9"). These changes were properly due to the dexterous effects of reactive oxygen species as $O_2^{\cdot-}$, H_2O_2 and NO. The depletion of liver GSH, as a nucleophilic scavenger of reactive metabolites, may be the main cause of the fallen serum GSH levels (Ballatori and Rebbeor, 1998). On the other hand this may be likely to reduce the ability of the liver and brain tissues to eliminate OFR (Yu, 1999; Basu et al., 2001 & Azzam, 2012). Also, GPx, and GSH-R were clearly decreased while GST was elevated in brain and liver tissues of envenomed animals (table "2" & figures "6", "7" & "8"). GSTs; a family of enzymes catalyzing the addition of GSH to endogenous and xenobiotic substrates with electrophilic functional groups, play important role in detoxification and metabolism of many xenobiotic and endobiotic compounds (Ji et al., 1992). The increment in GST activity (table "2" & fig. "8") is in consistent with the generation of FRs; as one of the defense mechanisms to detoxify or neutralize the toxic metabolites generated in the liver (El-Demerdash et al., 2005). GPx plays a

primary role in minimizing oxidative damage. GPx and GST work together with glutathione in the decomposition of H_2O_2 or other organic hydroperoxides to non-toxic products at the expense of GSH. On contrary, significant decreases in SOD activity and TAC (table "3" & figures "10" & "12") were reported. The role of SOD is to protect cells against oxidative damage caused by superoxide (SO) anion radicals. SODs cause rapid dismutation of SOD to a less reactive H_2O_2 molecule which is further degraded by CAT to water and oxygen (Wetscher et al., 1995). Thus the decreased SOD activity, in our study, may be related to adaptive response to the generated FRs to $O_2^{\cdot-}$ in the treated animals (Giller and Winge, 1984; Kale et al., 1999; Azzam, 2012 & Azzam et al., 2013). An association between GSH depletion and reduced SOD activity was suggested (Loven et al., 1986). Also, the decrements in SOD and CAT activities may be due to inactivation of mitochondrial and cytosolic enzymes, leading to a decrease in the metabolic activity (Kochar and Umathe, 2009). Alternatively, SO anion radicals are dismutated and catalyzed by SOD to H_2O_2 (Schild et al., 2003). Both pathways of SO anion radical reactions; the formation of H_2O_2 and the formation of $ONOO^-$, lead to a decrease in the activity of both SOD and CAT in the current study.

With respect to recovery trials of scorpion venom toxicity in animal models with safer, more effective, cheaper and more sustainable products with therapeutic impact, this work is an attempt and a developing effort pointing towards an upcoming transition from the wholesale coarsity of the serum-based antivenom to discover therapeutic natural solutions based on modern biotechnology. Thus scorpion haemolymph (SH) was used in this work, as discussed earlier, to ensure its protective

and/or treating ability of the scorpion-evoked OS. Scorpions are known to be quite resistant to their own venom. Some kill the scorpion and apply the scorpion's fluid into the site of sting thinking that it will neutralize the venom. SH was insured to have beneficial effects when applied to envenomed individuals in a controlled environment (Tripod.com, 2011; Azzam, 2012; Azzam et al., 2013 & Abdou et al., 2015). Our results; presented in tables "1", "2" & "3", investigated that the venom-induced hyperglycemia was returned to euglycemia in the pre and post-envenomed groups treated with SH. Thus, the noticed cellular dysfunctions subsequent to hyperglycemic oxidative stress after envenomation conditions in this study were normalized to the control values. The haemolymph have reported to have antihyperglycemic effects due to the presence of insulin like peptides; structurally related to that of insulin of mammals (Padmanabhan et al., 2012), so can reverse the metabolic disturbances by changes in the hormonal environment, especially insulin (Kankonkar et al., 1992; Murthy et al., 2002; 2003; Natu et al., 2006 & Azzam. 2012). Otherwise, the neutralization effect may be attributed to the non-enzymatic antioxidant activity predicted on the oxygenated haemocyanin (Quéinnec et al., 1999; Bolton et al., 2009 & Dann, 2017); a large multisubunit extracellular pigmented metalloprotein component in the SH that performs the important function of oxygen transport (Ali et al., 1995 & Azzam, 2012). It is worth to mention that the haemolymph is known by its ability to inhibit ROS by its bioactive peptides (Moselhy, 2012 & Padmanabhan et al., 2012). It was found to contain a number of effective antitoxins that may reverse the venom-toxins-metabolic disturbances (Harpaz, 1973 & Balasubramanian et al., 2017). Also, the

haemolymph was reported to have potent peptides showing diverse array of activity (Arul Prakash et al., 2011 & Padmanabhan et al., 2012) and plays a prominent role in host defense response including superoxide anion activity, (Kramer et al. 1977; Sahoo et al., 2005 & Moselhy, 2012). On other hand, a number of studies have reported that scorpion haemolymph have cross-species toxin inhibitory effects (Smith et al., 2000 & Abdou et al., 2015) but its mechanism of action remains unclear (Azzam, 2012 & Azzam et al., 2013). This may explain its activity to reverse the venom-toxins-metabolic disturbances. The attainment of normal blood glucose level following the SH indicated that SH may act as the scorpion antivenom (SAV) to neutralize its toxicity. Also, the amelioration efficacy of SH, under our experimental conditions, ensured that the time of treatment, the IV injection and the desired dose effectively and fastly neutralize the circulating venom and durably clears of free toxins from the circulation as well as their rapid redistribution from the peripheric compartment to the vascular one (Azzam, 2012 & Azzam et al., 2013).

Recommendations:

The management of diseases without any side effects remains a challenge in medicine. More trials to elucidate the therapeutic value of other scorpions' haemolymph against the cytotoxicity of venom and to separate its novel active molecule (s) are very important. More efforts to discover natural antivenins to prevent and treat scorpion sting symptoms are recommended. Also, a well-planned health education program might be useful to overcome the problems of sting bites especially scorpionism, predominately in developing countries.

References:

- 1- Abd Rabo, H. A. E. M. (2006): **Study of the influence of gamma radiation on certain pharmacological and biochemical actions of *Leiurus quinquestriatus* scorpion venom.** MSc. Thesis, National Center for Radiation Research and Technology Atomic Energy Authority, ARE.
- 2- Abdel-Rahman, M. A.; Harrison, P. L. and Strong, P. N. (2015): **Snapshots of scorpion venomics.** J. Arid Environ, 112 (B): 170–176.
- 3- Abdel-Rahman, M. A.; Omran, M. A. A.; Abdel-Nabi, I. M., Ueda, H. and McVean, A. (2009): **Intraspecific diversity in the Egyptian scorpion *Scorpio maurus palmatus* venom collected from different biotopes.** Toxicon, 53 (3): 349-359.
- 4- Abdel-Rahman, M. A.; Omran, M. A.; Abdel-Nabi, I. M.; Nassier, O. A. and Schemerhorn, B. J. (2010): **Neurotoxic and cytotoxic effects of venom from different populations of the Egyptian *Scorpio maurus palmatus*.** Toxicon, 55 (2-3): 298-306.
- 5- Abdel-Rahman, M. A.; Quintero-Hernandez, V. and Possani, L. D. (2013): **Venom proteomic and venomous glands transcriptomic analysis of the Egyptian scorpion *Scorpio maurus palmatus* (Arachnida: Scorpionidae).**Toxicon 74: 193–207.
- 6- Abdoon, N. A. S. (2004): **Toxigenesis and treatment of scorpion envenomation in man and experimental animals in Sudan.** PhD Thesis, College of Pharmacy, Khartoum University, Khartoum, Sudan.
- 7- Abdoon, N. A. S. and Fatani, A. J. (2009): **Correlation between blood pressure, cytokines and nitric oxide in conscious rabbits injected with *Leiurus quinquestriatus* scorpion venom.** Toxicon, 54 (4): 471-80.
- 8- Abdou, R. H.; Abd El-Ghany, F. S.; Abdel-Nabi, I. M. and Abdel-Rahman, M. A. (2015): **Toxicological effects of the horned viper venom *Cerastes cerastes* and their neutralization with the Egyptian scorpion *Androctonus australis* haemolymph.** Ind. J. Appl. Res., 5 (3): 37- 41.
- 9- Abou-Elezz, M. F.; Moustafa, A. Y. and El-Naggar, M. S. (2017): **Assessment of biochemical and histopatho-logical effects of crude venom of cone snail *Conus flavidus* on albino mice.** Intern. J. Ecotoxicol. Ecobiol., 2 (1): 33-44
- 10- Aboul-Soud, M. A. M.; Al-Othman, A. M.; El-Desoky, G. E.; Al-Othman, Z. A.; Yusuf, K.; Ahmad, J. and Al-Khedhairi, A. A. (2011): **Hepatoprotective effect of vitamin E/selenium against malathion-induced injuries on the antioxidant status and apoptosis-related gene expression in rats.** J. Toxicol. Sci., 36 (3): 285-96.
- 11- Adam, K. R. and Weiss, C. (1959): **Distribution of 5-hydroxytryptamine in scorpion venoms.** Nature, 183: 1389-99.
- 12- Adi-Bessalem, s.; Hammoudi-Triki, D. and Laraba-Djebari, F. (2014): **Scorpion venom interactions with the immune system.** Scorpion Venoms: Part of the Toxinology book series (TOXI), 4: 87-107.
- 13- Adibhatla, R. M. and Hatcher, J. F. (2008): **Altered lipid metabolism in brain injury and disorders.** In: Lipids in Health and

- Disease, Chapter 9, Quinn, P. J. & Wang, X. edns., Subcell. Biochem., 49: 241–268.
- 14- Aebi, H. (1984): **Catalase in vitro**. In Packer, L. Edn., Methods in Enzymology, Academic Press, San Diego, 105 (c): 121-126.
- 15- Agarwal, A.; Aponte-Mellado, A.; Premkumar, B. J.; Shaman, A Gupta, S. A. (2012): **The effects of oxidative stress on female reproduction: A review**. Reprod. Biol. Endocrinol., 10 (1): 49-79.
- 16- Ahmed, A. E.; Abdel-Baseer, K. A.; Saad, K.; Hassan, A. F. and El-Houfey, A. A. (2015): **Endocrinological and biochemical changes of scorpionism in children in Upper Egypt**. Therap. Adv. Endocrinol. Metabol., 6 (5): 210 – 216.
- 17- Aispuro-Hernandez, E.; Garcia-Orozeo, K. D.; Muthlia-Almazan, A.; Del-Toro-Sanchez, L.; Robles-Sanchez, R. M.; Hernandez, J.; Gonzalez-Aguilar, G.; Yepiz-Plascencia, G.; Sotelo-Mundo, R. R. (2008): **Shrimp thioredoxin is a potent antioxidant protein**. Comp Biochem Physiol C Toxicol Pharmacol., 148 (1): 94-99.
- 18- Aksoy, S.; Cam, N.; Gurkan, U.; Oz, D.; Özden, K.; Altay, S.; Durmus, G. and Agirbasli, M. (2013): **Oxidative stress and severity of coronary artery disease in young smokers with acute myocardial infarction**. Cardiol. J., 19 (4): 381-6.
- 19- Al Asmari, A. K.; Al-Saief, A. A. and Abdo, N. M. (2007): **Morphological identification of scorpion species from Jazan and Al-Madina Al-Munawwara Regions, Saudi Arabia**. J. Venom. Anim. Toxins Incl. Trop. Dis., 13 (4): 821-843
- 20- Al Asmari, A. K.; Al-Saief, A. A.; Abdo, N. M. and Al-Moutaery, K. R. (2009b): **New additions to the scorpion fauna of Riyadh region, Saudi Arabia**. J. Venom. Anim. Toxins Incl. Trop. Dis., 15 (4): 612-632.
- 21- Al Asmari, A. K.; Al-Saief, A. A.; Abdo, N. M. and Al-Moutaery, K. R. (2009a): **The scorpion fauna of Al-Baha and Hail Regions, Saudi Arabia**. J. Biolo. Sci., 9 (2): 96-108.
- 22- Al Asmari, A. K.; Al-Saief, A. A.; Abdo, N. M.; Al-Moutaery, K. R. and Al-Harbi, N. O. (2013): **A review of the scorpion fauna of Saudi Arabia**. Egypt. J. Nat. Hist., 6: 1 - 21.
- 23- Al Asmari, A. K.; Al-Saief, A. A.; Hassen, M. A. and Abdulkaksood, N. A. (2008): **Role of prazosin on cardiovascular manifestations and pulmonary edema following severe scorpion stings in Saudi Arabia**. Saudi Med. J., 29 (2): 299-302.
- 24- Al Asmari, A. K.; Al-Zahrani, A. G. and Al-Jowhary, S. (2012): **Clinical aspects and frequency of scorpion stings in the Riyadh Region of Saudi Arabia**. Saudi Med. J., 33 (8): 852-858.
- 25- Al Asmari, A. K.; Khan, H. A. and Manthiri, R. A. (2012): **Rapid profiling of crude scorpion venom using liquid chromatography and its relevance to species identification**. Acta Chromatogr., 24 (3): 501-509.
- 26- Al Asmari, A. K.; Khan, H. A. and Manthiri, R. A. (2015): **Effect of *Androctonus bicolor* scorpion venom on the activities of serum enzymes in rats**. Int. J. Clin. Exp. Med., 8 (7): 11734-11737.
- 27- Al Asmari, A. K.; Khan, H. A. and Manthiri, R. A. (2016): **Effect of *Androctonus bicolor* scorpion venom on**

- serum electrolytes in rats: A twenty four hour time course study. Hum. Exp. Toxicol., 35 (3): 293-296.
- 28- Al Asmari, A. K.; Khan, H. A.; Manthiri, R. A.; Al Yahya, K. M. and Al Otaibi, K. E. (2014): **Effects of *Echis pyramidum* snake venom on hepatic and renal antioxidant enzymes and lipid peroxidation in rats.** J. Biochem. Mol. Toxicol., 28: 407- 412.
- 29- Al Asmari, A. K.; Kunnathodi, F.; Al Saadon, K. and Idris, M. M. (2016): **Elemental analysis of scorpion venoms.** J Venom Res., 7: 16-20.
- 30- Al Asmari, A. K.; Manthiri, R. A.; Abdo, N. M.; Duaiji, F. A. and Khan, H. A. (2017): **Saudi medicinal plants for the treatment of scorpion sting envenomation.** Saudi J. Biol. Sci., 24 (6): 1204-1211.
- 31- Al Asmari, A. K.; Manthiri, R. A.; Osman N. M. A.; Al-Otaibi, A. F. and Al-Asmari, S. A. (2016): **Beneficial role of quercetin on *Echis coloratus* snake venom induced hepato-renal toxicity in rats.** J. Biol. Sci., 16 (4): 102-111.
- 32- Al-Gayyar, M. M.; Eissa, L. A.; Rabie, A. M. and El-Gayar, A. M. (2007): **Measurements of oxidative stress status and antioxidant activity in chronic leukaemia patients.** J. Pharm. Pharmacol, 59 (5): 409-417.
- 33- Ali, S. A.; Zaidi, Z. H. and Abbasi, A. (1995): **Oxygen transport proteins: Structure and organization of hemocyanin from scorpion (*Buthus indicus*).** 112 (1): 225-32.
- 34- Al-kahlout, B. H.; Muhammad, M.; Abid, M. M.; Kasim, M. M.; Shumaila, M. and Haneef, M. D. (2015): **The need for regional management guidelines in emergency departments: Epidemiological review of scorpion stings in Qatar.** Saudi Med. J. 36 (7): 851-855.
- 35- Allen, R. G.; Toy, R. K.; Newton, R. K.; Farmer, K. J. and Sohal, R. S. (1985_a): **Effect of experimentally altered glutathione levels on life span, metabolic rate, superoxide dismutase, catalase and inorganic peroxides in the adult housefly, *Musca domestica*.** Comparative Biochemistry and Physiology Part C Comparative Pharmacology and Toxicology, 82 (2): 399-402.
- 36- Al-Omar, M. A.; Beedham, C. and Alsarra, I. A. (2004): **Pathological roles of reactive oxygen species and their defence mechanisms.** J. Saudi Pharm., 12: 1-18.
- 37- Al-Ramahi, H. M. and Al-Hasnawi, M. H. (2002): **diagnostic and epidemiologic study of fat tail scorpion (*Androctonus crassicauda*) in Mid Euphrates Region/ Iraq.** College of Veterinary Medicine, Babylon University, Iraq, 10 (10): 263 - 269.
- 38- Al-Sadoon, M. K. and Jarrar, B. A. (2003): **Epidemiological study of scorpion stings Saudi Arabia between 1993 and 1997.** J. Venom. Anim. Toxins, 9 (1): 1- 8.
- 39- Al-Sheikh, Y. A.; Ghneim, H. K.; Aljaser, F. S.; Aboul-Soud, M. A. M. (2017): **Ascorbate ameliorates *Echis coloratus* venom-induced oxidative stress in human fibroblasts.** Exp. Therap. med., 14 (1): 703- 713.
- 40- Amitai, Y. (1998): **Clinical manifestations and management of scorpion envenomation.** Public Health Rev., 26 (3):

- 257-263.
- 41- Annane, D.; Sanquer, S.; Sebillé, V.; Faye, A.; Djuranovic, D.; Raphaël, J. C.; Gajdos, P. and Bellissant, E. (2000): **Compartmentalized inducible nitric oxide-synthase activity in septic shock.** Lancet, 355 (9210): 1143-1148.
- 42- Aprioku, J. S. (2013): **Pharmacology of free radicals and the impact of reactive oxygen species on the testis.** J. Reprod. Infertil., 14 (4): 158–172.
- 43- Arul Prakash, A.; Balasubramanian, S.; Gunasekaran, G.; Prakash, M. and Senthil Raja, P. (2011): **Control of Clinical Pathogens by the Haemolymph of *Paratelphusa hydrodromous*; a Freshwater Crab.** ISRN Pharma., 2011: 642768, 4 pages
- 44- Aruoma, O. I.; Halliwell, B.; Gajewski, E. and Dizdaroglu, M. (1991): **Copper-ion-dependent damage to the bases in DNA in the presence of hydrogen peroxide.** Biochem. J., 273 (3): 601- 604.
- 45- Ayala, A.; Muñoz, M. F. and Argüelles, S. (2014): **Lipid Peroxidation: Production, metabolism and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal.** Oxi. med. Cell. Long., 2014 (6): 360438.
- 46- Azevedo A. D.; Cunha Melo, J. R. and Freire-Maia, L. (1979): **Arterial hypertension and pulmonary edema induced by catecholamines in unanesthetized rats.** Pharmacol. Res. Commun., 11: 157-67.
- 47- Azzam, N. N. (2012): **Biochemical and physiological studies on the biological activities of the venom and haemolymph of the Egyptian scorpion *Androctonus australis*.** PhD Thesis, Zoology Dept., Suez Canal University, Ismailia, ARE.
- 48- Azzam, N. N.; Abdel-Nabi, I. M. and Abdel-Rahman; M. A. (2013): **Evidence for free radicals generation by the Egyptian *Androctonus australis* scorpion venom and the potency of the haemolymph to attenuate these metabolic disorders.** The 10th Annual Congress of Medical Sciences (ACMS), Under the theme: "Medical Research towards Healthy Society", NRC, Dokki, Cairo, Egypt, 12th - 16th Jan.
- 49- B'usse, R. and M'ulsch, A. (1990): **Induction of nitric oxide synthase by cytokines in vascular smooth muscle cells.** FEBS Lett ., 275 (1-2): 87-90.
- 50- Balasubramanian, S.; Priya, K.; Revathi, I.; Revathi, A.; Venkatesh, P. and Gunasekaran, G. (2017): **Screening of antibacterial activity and biochemical assay from haemolymph of cockroach *Blatta orientalis* (Linnaeus, 1758).** JEZS, 5 (3): 753-758
- 51- Ballatori, N. and Rebbeor, J. E. (1998): **Roles of MRP₂ and OTP₁ in hepatocellular export of reduced glutathione.** Seminars in liver disease, 18 (4): 377-387.
- 52- Basu, S.; Michaëlsson, K.; Olofsson, H.; Johansson, S. and Melhus, H. (2001): **Association between oxidative stress and bone mineral density.** Biochem. Biophys. Res. Commun., 288 (1): 275-279.
- 53- Bawaskar, H. S. and Bawaskar, P. H. (2007): **Utility of scorpion antivenin in the management of severe *Mesobuthus***

- tamulus (Indian red scorpion) envenoming at rural setting.** J. Assoc. Physicians India (JAPI), 55: 14-21.
- 54- Bawaskar, H. S. and Bawaskar, P. H. (2012): **Scorpion sting: Update.** J. Assoc. Physicians India, 60 (1): 46-55 (9):797-803.
- 55- Béchohra, L.; Laraba-Djebari, F. and Hammoudi-Triki, D. (2016): **Cytotoxic activity of *Androctonus australis hector* venom and its toxic fractions on human lung cancer cell line.** J. Venom. Animals Toxins incl. Trop. Dis., 22(1): 29-44.
- 56- Berger, M.; Beys-da-Silva, W. O.; Santi, L.; de Oliveira, I. M.; Jorge, P. M.; Henriques, J. A. P.; Driemeier, D.; Vieira, M. A. R. and Guimarães, J. A. (2013): **Acute *Lonomia oblique* caterpillar envenomation-induced physiopathological alterations in rats: Evidence of new toxic venom activities and the efficacy of serum therapy to counteract systemic tissue damage.** Toxicon 74: 179 - 192
- 57- Berlett, B. S. and Stadtman, E. R. (1997): **Protein oxidation in aging; disease, and oxidative stress.** J. Biol. Chem., 272 (33): 20313-20316.
- 58- Beutler, E.; Doron, O. and Kelly, B. (1963): **Improved method for the determination of blood glutathione.** J. Lab. Clin. Med., 61 (5): 882-888.
- 59- Bewick, V.; Cheek, L. and Ball, J. (2004): **Statistics review 9: One-way analysis of variance.** Crit. Care, 8 (2): 130–136.
- 60- Bhattacharjee, S. (2014): **Membrane lipid peroxidation and its conflict of interest: The two faces of oxidative stress.** CURR. SCI., 10 (11): 1181- 1193.
- 61- Bhaumik, G.; Srivastava, K. K.; Selvamurthy, W. And, Purkayastha, S. S. (1995): **The role of free radicals in cold injuries.** Int. J. Biometeorol., 38 (4): 171-175.
- 62- Blokhina, O.; Virolainen, E. and Fagerstedt, K. V. (2003): **Antioxidants, oxidative damage and oxygen deprivation stress: A review.** Annals of Botany, 91 (2): 179 -194.
- 63- Bolotina V. M.; Najibi, S.; Palacino, J. J.; Pagano, P. J.; Cohen, R. A. 1994): Nitric oxide directly activates calcium-dependent potassium channels in vascular smooth muscle. Nature, 368 (6474): 850-853.
- 64- Bolton, J. C.; Collins, S.; Smith, R.; Perkins, B.; Bushway, R.; Bayer, R. and Vetelino, J. (2009): **Spectroscopic analysis of haemolymph from the American lobster (*Homarus americanus*).** J. Shellfish Res., 28 (4): 905-912.
- 65- Bosmans, F. and Tytgat, J. (2007): **Voltage-gated sodium channel modulation by scorpion α -toxins.** Toxicon, 49 (2): 142–158.
- 66- Boşnak, M.; Ece, A.; Yolbaş, İ.; Boşnak, V.; Kaplan, M. and Gürkan, F. (2009): **Scorpion sting envenomation in children in southeast Turkey.** Wilderness Environ. Med., 20: 118-124.
- 67- Bouaziz, M.; Bahloul, M.; Kallel, H.; Samet, M.; Ksibi, H.; Dammak, H.; Ahmed, M. N.; Chtara, K.; Chelly, H.; Hamida, C. B. and Rekik, N. (2008): **Epidemiological, clinical characteristics and outcome of severe scorpion envenomation in South Tunisia: Multivariate analysis of 951 cases.** Toxicon, 2 (8): 918-926.
- 68- Boveris, A. and Navarro, A. (2008): **Brain**

- mitochondrial dysfunction in aging.** Life, 60 (5): 308-314.
- 69- Bowles, D. E. and Swaby, J. A. (2006): **Field guide to venomous and medically important invertebrates affecting military operations: Identification, Biology, Symptoms, Treatment.** Version 2, US Air Force Institute for Operational Health (AFIOH), 2513 Kennedy Circle Drive Brooks City-Base, TX 78235.
- 70- Bowles, D. E. and Swaby, J. A. (2006): **Field guide to venomous and medically important invertebrates affecting military operations: Identification, Biology, Symptoms, Treatment.** Version 2, US Air Force Institute for Operational Health (AFIOH) 2513 Kennedy Circle Drive Brooks City-Base, TX 78235.
- 71- Bradley, R. (1984): **The influence of the quantity of food on fecundity in the desert grassland scorpion (*Paruroctonus utahensis*) (Scorpionida, Vaejovidae): An experimental test.** Oecologia, 62: 53-56.
- 72- Braunersreuther, V. and Jaquet, V. (2012): **Reactive oxygen species in myocardial reperfusion injury; from physiopathology to therapeutic approaches.** Curr. Pharm. Biotechnol., 13 (1): 97–114.
- 73- Bücherl, W. and Buckley, E. E. (1971): **Venomous Animals and Their Venoms. Venomous Vertebrates, Vol. II, 1st edn.,** New York and London, Academic Press, 712 Pages
- 74- Cakatay, U.; Kayali, R. and Uzun, H. (2008): **Relation of plasma protein oxidation parameters and paraoxonase activity in the ageing population.** Clin. Exp. Med., 8 (1): 51-57.
- 75- Caliskan, F.; Quintero-Hernandez, V.; Restano-Cassulini, R.; Coronas-Valderrama, F. I.; Corzo, G. and Possani, L. D. (2013): **Molecular cloning and biochemical characterization of the first Na⁺-channel and α -type toxin peptide (Acra4) from *Androctonus crassicauda* scorpion venom.** Biochimie., 95 (6):1216-1222.
- 76- Candido, D. M. and Lucas S. (2004): **Maintenance of scorpions of the genus *Tityus Koch* (scorpiones, buthidae) for venom obtention at Instituto Butantan, São Paulo, Brazil.** J. Venom Anim. Toxins. Incl. Trop. Dis., 10 (1): 86 – 97.
- 77- Cao, Z. J.; Luo, F.; Wu, Y. L.; Mao, X. and Li, W. X. (2006): **Genetic mechanisms of scorpion venom peptide diversification.** Toxicon, 47: 348-355.
- 78- Cao, Z.; Di, Z.; Wu, Z. and Li, W. (2014): **Overview of scorpion species from China and their toxins.** Toxins (Basel), 6 (3): 796–815.
- 79- Carmo, A.; Chatzaki, M.; Horta, C.; Magalhães, B.F.; Oliveira-Mendes, B. B. R.; Chavez-Olortegui, C. and Kalapothakis, E. (2015): **Evolution of alternative methodologies of scorpion antivenoms production.** Toxicon, 97 (2015): 64-74
- 80- Chakravarthy, A. K. and Sridhara, S. (2016): **Arthropod Diversity and Conservation in the Tropics and Sub-tropics.** Springer Singapore, 1stedn., 435 pages.
- 81- Chaubey, M. K. (2017): **Pharmacological changes after black scorpion *Heterometrus fastigiosus* Couzijn**

- envenomation in mice.** Journal of Medical Sciences, 17 (2): 75-80
- 82- Chaudhuri, K. R.; Healy, D. G. and Schapira, A. H. (2006): **Non-motor symptoms of Parkinson's disease: diagnosis and management.** Lancet Neurol., 5 (3): 235-45.
- 83- Chevion, M.; Berenshtein, E. and Stadtman, E. (2000): **Human studies related to protein oxidation: Protein carbonyl content as a marker of damage.** Free Rad. Res., 33 (Suppl): S99-108
- 84- Chippaux, J. P. and Goyffon, M. (2010): **Epidemiology of scorpionism: A global appraisal.** Acta Trop., 107 (2): 71-79.
- 85- Chippaux, J.-P. (2012): **Emerging options for the management of scorpion stings.** Drug Des. Devel. Ther., 6: 165-173.
- 86- Choudhry, N. K.; Fischer, M. A.; Avorn, J.; Liberman, J. N.; Schneeweiss, S.; Pakes, J.; Brennan, T. A. and Shrank, W. H. (2011): **The implications of therapeutic complexity on adherence to cardiovascular medications.** Arch. Intern. Med., 171 (9): 814-822.
- 87- Choudhry, P. K. R. Pandey, R. and Deshpande, S. B. (2011): **2-Deoxy-D-glucose reverses the Indian red scorpion venom-induced cardiopulmonary abnormalities in anesthetized rats.** Ind. J. Exp. Biol., 49 (1): 16-23
- 88- Chowdhary, S. and Townend, J. N. (1991): **Role of nitric oxide in the regulation of cardiovascular autonomic control.** Clin. Sci. (Lond), 97 (1): 5-17.
- 89- Conde, R.; Zamudio, F. Z.; Becerril, B. and Possani, L.D. (1999): **Phospholipin; a novel heterodimeric phospholipase A2 from *Pandinus imperator* scorpion venom.** FEBS Lett., 460 (1): 447-450.
- 90- Couto, N.; Malys, N.; Gaskell, S. J. and Barber, J. (2013): **Partition and turnover of glutathione reductase from *Saccharomyces cerevisiae*: A Proteomic Approach.** J. Proteome. Res., 12 (6): 2885–2894
- 91- Curry, S. C.; Vance, M. V.; Ryan, P. J.; Kunkel, D. B. and Northey, W. T. (1984): **Envenomation by the scorpion *Centruroides sculpturatus*.** J. Toxic. Clin. Toxic., 21 (4-5): 417-449.
- 92- Dalle-Donne, I.; Rossi, R.; Colombo, R.; Giustarini, D. and Milzani, A. (2006): **Biomarkers of oxidative damage in human disease.** Clin. Chem., 52 (4): 601-623.
- 93- Dalle-Donne, I.; Rossi, R.; Giustarini, D.; Milzani, A. and Colombo, R. (2003): **Protein carbonyl groups as biomarkers of oxidative stress.** Clin. Chim. Acta, 329 (1): 23-38.
- 94- Dancey, C. P. and Reidy, J. (2002): **Statistics without Maths for Psychology: Using SPSS for Windows™.** Harlow, Essex: Prentice-Hall, Harlow, UK, 2ndedn., 604 pages.
- 95- Danielsen, T. (2015): **Protein markers of oxidative stress damage :Applications to experimental research and effects of fatty acids.** MSc Thesis, Biosciences Dept., Faculty of Mathematics and Natural Sciences, Oslo University, Norway.
- 96- Dann, L. E. (2017): **antioxidant capacity in the haemolymph of the marine isopod *Pentidotea resecata*.** MSc Thesis, Walla Walla University, Washington, USA.

- 97- Danny, L. (2015): **Why A Single Vial Of Antivenom Can Cost \$14,000.** Smithsonian, Smartnews, <https://www.smithsonianmag.com/smart-news>.
- 98- Daoudi, K.; Chgoury, F.; Rezzak, M.; Bourouah, Q.; Boussadda, L.; Soukri, A.; Sabatier, J.-M. and Oukkache, N. (2017): **Consequences of *Androctonus mauretanicus* and *Buthus occitanus* scorpion venoms on electrolyte levels in rabbits.** Heliyon, 3 (2017): e 00221.
- 99- Dauqan, E. M. A.; Abdullah, A. and Sani, H. A. (2011): **Natural antioxidants, lipid profile, lipid peroxidation, antioxidant enzymes of different vegetable.** Adv. J. Food Sci. Tech., 3 (4): 308- 316.
- 100- Davila, S.; Pinero, D.; Bustos, P.; Cevallos, M. A. and Davila, G. (2005): **The mitochondrial genome sequence of the scorpion *Centruroides limpidus* (Karsch 1879) (Chelicerata; Arachnida).** Gene., 360 (2): 92-102.
- 101-de la Vega, R. c. R. Schwartz, E. F. and Possani, L. D. (2010): **Mining on scorpion venom biodiversity.** Toxicon, 56 (7): 1155-61.
- 102-De Maria, L.; Vind, J.; Oxenboll, K. M.; Svendsen, A. and Patkar, S. (2007): **Phospholipases and their industrial applications.** Appl. Microbiol. Biotechnol., 74 (2): 290-300.
- 103-de Oliveira, G. H.; Cerni, F. A.; Cardoso, L. A.; Arantes, E. C. and Berto, M. (2016): ***Tityus serrulatus* envenoming in non-obese diabetic mice: A risk factor for severity.** J. Venom. Anim. Toxins Incl. Trop. Dis., 22 (1): 26.
- 104- de Roodt, A. R.; Coronas, F. I. V.; Lago, N.; Gonzalez, M. E.; Laskowicz, R. D.; Beltramino, J. C.; Saavedra, S.; López, R. A.; Reati, G. J.; Vucharchuk, M. G.; Bazán, E.; Varni, L.; Salomón, O. D. and Possani, L. D. (2010): **General biochemical and immunological characterization of the venom from the scorpion *Tityus trivittatus* of Argentina.** Toxicon, 55 (2-3): 307–319.
- 105- de Roodt, A. R.; Garcia, S. I.; Salomon, O. D.; Segre, L.; Dolab, J. A.; Funes, R. F. and De Titto, E. H. (2003): **Epidemiological and clinical aspects of scorpionism by *Tityus trivittatus* in Argentina.** Toxicon, 41 (8): 971-977.
- 106- Dean, R. T.; Fu, S.; Stocker, R. and Davies, M. J. (1997): **Biochemistry and pathology of radical-mediated protein oxidation.** J. Biochem., 324: 1–18.
- 107- Deori, M.; Devi, D.; Kumari, S.; Hazarika, A.; Kalita, H.; Sarma, R. and Devi, R. (2016): **Antioxidant effect of sericin in brain and peripheral tissues of oxidative stress induced hypercholesterolemic rats.** Front. Pharmacol., 15 (7): 319, eCollection 2016.
- 108- Desouky, M. M. and Al-Shammari, A. M. (2011): **Scorpions of Hail region, northern Saudi Arabia, and molecular phylogenetics of two common species; *Androctonus crassicauda* and *Scorpio maurus kruglovi*.** Bulletin of the British Arachnological Society, 15 (6): 193-200.
- 109- Devasagayam, T. P.; Boloor, K. K. and Ramasarma T. (2003): **Methods for estimating lipid peroxidation: An analysis of merits and demerits.** IJBB, .40 (5): 300-308.

- 110- Ding, J.; Chua, P.-J.; Bay, B.-H. and Gopalakrishnakone, P. (2014): **Scorpion venoms as a potential source of novel cancer therapeutic compounds**. Experimental Biology and Medicine, 239: 387–393.
- 111- Dittrich, K., Ahmed, R. and Ahmed, Q. A. (2002): **Cardiac arrest following scorpion envenomation**. Annals. Saudi Med., 22 (1 -2): 87-90.
- 112- Dousset, E.; Carrega, L.; Steinberg, J. G.; Clot-Faybesse, O.; Jouirou, B.; Sauze, N.; Devaux, C.; Autier, Y.; Jammes, Y.; Martin-Eauclaire, M. F. and Guieu, R. (2005): **Evidence that free radical generation occurs during scorpion envenomation**. Comp. Biochem. Physiol. Toxicol. Pharmacol., 140 (c) : 221-226.
- 113- Dumas, B. T., Watson, W. A. and Biggs, H. G. (1997): **Albumin standards and the measurement of serum albumin with bromocresol green**. Clin. Chim. Acta., 258 (1): 21-30.
- 114- Dunlop, J. A. and Selden P. A. (2013): **Scorpion fragments from the Silurian of Powys, Wales**. Arachnology, 16 (1): 27-32.
- 115- Dutertre, S. and Lewis, R. J. (2010): **Use of venom peptides to probe ion channel structure and function**. J. Biol. Chem., 285 (18): 13315-13320.
- 116- Ebrahimi, M.; Azizi, K.; Moemenbellah-Fard, M. D.; Fakoorziba, M. R. and Soltani, A. (2015): **Morphometry indices of the black fat-tailed scorpion *Androctonus crassicauda* (Scorpiones Buthidae), from Fars Province, southern Iran**. Entomology, 12 (1): 39-47
- 117- Ebrahimi, V.; Hamdami, E.; Moemenbellah-Fard, M. D. and Jahromi, S. E. (2017): **Predictive determinants of scorpion stings in a tropical zone of south Iran: Use of mixed seasonal autoregressive moving average model**. J. Venom. Anim. Toxins Incl. Trop. Dis., 23 (1): 39- 51.
- 118- EL-Asmar, M. F.; Farag R. M.; Shoukry, S. and EL-Smimi, I. R. (1979): **Effect of scorpion (*Leiurus quinquestriatus H and E*) venom on lipid metabolism**. Toxicol., 17 (3): 279-83.
- 119- El-Asmar, M.; Soliman, S. F.; Ismail, M. and Osman, O. H. (1974): **Glycemic effect of venom from the scorpion *Buthus minax* (L. Koch)**. Toxicol., 12 (3): 249-51.
- 120- El-Demerdash, F. M.; Yousef M. I. and Abou El-Naga N. I. (2005): **Biochemical study on the hypoglycemic effects of onion and garlic in alloxan-induced diabetic rats**. Food Chem. Toxicol., 43 (1): 57-63.
- 121- Esterbauer, H.; Eckl, P. and Ortner, A. (1990): **Possible mutagens derived from lipids and lipid precursors**. Mutation Research, 238 (3): 223–233.
- 122- Esterbauer, H.; Schaur, R. J. and Zollner, H. (1991): **Chemistry and Biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes**. Free Radical Biology and Medicine, 11 (1): 81–128.
- 123- Fatani, A. J.; Ahmed, A. A. E.; Abdel-Halim; R. M.; Abdoon, N. A. and Darweesh, A. Q. (2010): **Comparative study between the protective effects of Saudi and Egyptian antivenoms, alone or in combination with ion channel modulators, against**

- deleterious actions of *Leiurus quinquestriatus* scorpion venom.** *Toxicon*, 55 (4): 773-786.
- 124- Fatani, A. J.; Al-Zuhair H. A.; Yaquob H. I.; Abdel-Fattah A. A .; El-Sayed M. I. and El-Sayed F. A . (2006): **Protective effects of the antioxidant *Ginkgo biloba* extract and the protease inhibitor aprotinin against *Leiurus quinquestriatus* venom-induced tissue damage in rats.** *J. Venom. Anim. Toxins incl. Trop. Dis.*, 12 (2): 255-275
- 125- Fedorova, I.; Mikov, A.; Maleeva, E.; Andreev, A.; Bocharov, E.; Tikhonov, D. and Kozlov, S. A. (2014): **Novel insect-toxin from *Tibellus oblongus* spider venom.** *JFEBS*, 281: 414-415.
- 126- Field, A. P. (2000): **Discovering statistics using SPSS for windows: Advanced techniques for the beginner (Introducing Statistical Methods series).** London, Sage Publications Ltd; Pap/Cdr edition, 512 pages.
- 127- Fossati, P.; Prencipe, L. and Berti, G. (1980): **Use of 3,5-dichloro-2-hydroxybenzene-sulfonic acid/4-aminophenazne chromogenic system in direct assay of uric acid in serum and urine.** *Clin. Chem.*, 26 (2): 227-231.
- 128- Fox, J. W. and Serrano, S. M. T. (2008): **Exploring snake venom proteomes: Multifaceted analyses for complex toxin mixtures.** *Proteomics*, 8: 909-920.
- 129- Francielle, A.; Cordeiro, F. A.; Amorim, F. G.; Anjolette, F. A. P. and Arantes, E. C. (2015): **Arachnids of medical importance in Brazil: Main active compounds present in scorpion and spider venoms and tick saliva.** *J. Venom. Anim. Toxins. Incl. Trop. Dis.*, 21 (24):1-14.
- 130- Fraser, C. (1991): **Biological variation in clinical chemistry: An update: collated data (1988-1991).** *Arch. Pathol. Lab. Med.*, 116: 916-923.
- 131- Freire-Maia, L.; Campos, J. A. and Amaral, C. F. S. (1994): **Approaches to the treatment of scorpion envenoming.** *Toxicon*, 32: 1009-1014.
- 132- Frembgen, J. W. (2004): **The Scorpion in Muslim Folklore.** *Asian Folklore Studies*, 63 (1): 95 -123
- 133- Fry, B. G.; Roelants, K. and Norman, J. A. (2009): **Tentacles of venom: Toxic protein convergence in the Kingdom *Animalia*.** *J. Mol. Evol.*, 68 (4): 311–321
- 134- Fry, B. G.; Roelants, K.; Champagne, D. E.; Scheib, H.; Tyndall, J. D. A.; King, G. F.; Norman, J. A.; Lewis, R. J.; Norton, R. S.; Renjifo, C. and de la Vega, R. C. R. (2009): **The Toxicogenomic Multiverse: Convergent recruitment of proteins into animal venoms.** *Annu. Rev. Genom. Hum. Genet.*, 10: 483-511.
- 135- Ganong, W. F. (1987): **Review of Medical Physiology.** 13th edn., Prentice Hall Publisher, New York: Appleton and Lange, p. 283.
- 136- Gazarian, K. G.; Gazarian, T.; Hernández, R. and Possani, L. D. (2005): **Immunology of scorpion toxins and perspectives for generation of anti-venom vaccines.** *Vaccine*, 23 (2005): 3357-3368.
- 137- Geethabali and Rajashekhar, K . P. (1988): **Ionic composition of the haemolymph of the whip scorpion; *Thelyphonus indicus***

- stoliczka*, and its saline formula. Acta Arachnol., 36 (2): 87-92.
- 138- Ghneim, H. K. (2017): **The effect of *Echis coloratus* venom on biochemical and molecular markers of the antioxidant capacity in human fibroblasts.** Libyan J. Med., 12 (1): 1304515.
- 139- Gill, S. S.; Khan, N. A.; Anjum, N. A. and Tuteja, N. (2011): **Amelioration of cadmium stress in crop plants by nutrients management: morphological, physiological and biochemical aspects.** Plant Stress, 5:1–23
- 140- Giller, B. L. and Winge, D. R. (1984): **Subcellular disturbance in rat liver.** Methods Enzymol., 105: 105-115.
- 141- Goldberg, D.M. and Spooner, R. J. (1983): **Assay of glutathione reductase.** In: Methods of enzymatic analysis, Bergmeyer, H.U. Edn., Verlag Chemie, Weinheim, Germany, 3rd edn., 3: 258-265.
- 142-Goldstein, R. E.; Abumrad, N. N.; Wasserman, D. H. and Cherrington, A. D. (1995): **Effects of an acute increase in epinephrine and cortisol on carbohydrate metabolism during insulin deficiency.** Diabetes, 44 (6): 672-81.
- 143- Gönenç, A.; Erten, D.; Aslan, S.; Akıncı, M.; Şimşek, B. and Torun, M. (2006): **Lipid peroxidation and antioxidant status in blood and tissue of malignant breast tumor and benign breast disease.** Cell Biol. Int., 30 (4): 376-380.
- 144- Goyffon, M. and El-Ayeb, M. (2002): **Epidemiologie du scorpionism.** Infotox, 15 (2): 2–6.
- 145- Green, L. C.; Wagner, D. A.; Glogowski, J., Skipper, P. L.; Wishnok, J. S. and Tannenbaum, S. R. (1982): **Analysis of nitrate, nitrite and [15N] nitrate in biological fluids.** Analyt. Biochem., 126(1): 131-138.
- 146- Grune, T.; Reinheckel, T. and Davies, K. J. (1996): **Degradation of oxidized proteins in K562 human hematopoietic cells by proteasome.** J. Biol. Chem., 271 (26): 15504-15509.
- 147- Grune, T.; Reinheckel, T.; Joshi, M. and Davies, K. J. (1995): **Proteolysis in cultured liver epithelial cells during oxidative stress: Role of the multicatalytic proteinase complex, proteasome.** J. Biol. Chem., 270 (5): 2344–2351.
- 148- Gwee, M. C.; Nirthanan, S.; Khoo, H. E.; Gopalakrishnakone, P.; Kini, R. M. and Cheah, L. S. (2002): **Autonomic effects of some scorpion venoms and toxins.** Clin. Exp. Pharmacol. Physiol., 29 (9): 795-801.
- 149- Habig, W. H.; Pabst, M. J. and Jakoby, W. B. (1974): **Glutathione-S-transferase: The first enzymatic step in mercapturic acid formation.** J. Biol. Chem., 249 (22): 7130-7139.
- 150- Halliwell, B. (2006): **Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life.** Plant Physiol., 141 (2): 312-322.
- 151- Halliwell, B. and Gutteridge, J. (2015.): **Free radicals in biology and medicine.** NY: Oxford Univ. Press., 5th edn.
- 152- Halliwell, B. and Gutteridge, M. (1989): **Free Radicals in Biology and Medicine.** 2nd edn., Oxford Univeristy Press, New York.
- 153- Halliwell, B. and Whiteman, M. (2004): **Measuring reactive species and oxidative**

- damage in vivo and in cell culture: How should you do it and what do the results mean?** Br. J. Pharmacol.,142 (2): 231-55.
- 154- Hammoudi-Triki D.; Ferquel, E.; Robbe-Vincent, A.; Bon, C.; Choumet, V. and Laraba-Djebari, F. (2004): **Epidemiological data, clinical admission gradation and biological quantification by ELISA of scorpion envenomation in Algeria: Effect of immunotherapy.** Trans. R. Soc. Trop. Med. Hyg., 98(4): 240-50.
- 155- Hariprasad, G.; Singh, B.; Das, U.; Ethayathulla, A. S.; Kaur, P.; Singh, T.P. and Srinivasan, A. (2007): **Cloning, sequence analysis and homology modeling of a novel phospholipase A2 from *Heterometrus fulvipes* (Indian black scorpion).** DNA Seq., 18(3): 242–246.
- 156- Harpaz, I (1973): **Early entomology in the Middle East.** Smith, R. F.; Mittler, T. E. and Smith, C. N. edn., History of Entomology, Annual Reviewers Inc., California.
- 157- Harvey, A. L.; Anderson, A. J.; Braga, M. F. M.; Marshall, D. L.; Rowan, E. G.; Vatanpour, H.; Castañeda, O. and Karlsson, E. (1992): **Toxins affecting neuronal ion channels.** In: Gopalakrishnakone P. & Tan, C. K. edn., Recent advances in toxicology research, Singapore: National University of Singapore, 59-70.
- 158- Hmed, B.; Serria, H. T. and Mounir, Z. K. (2013): **Scorpion peptides: Potential use for new drug development.** J. Toxicol., 2013 (10): 958797, 15 pages.
- 159- Hosseini, Z.; Khosravi, M.; Ghorbanpoor, M. and Mayahi, M. (2017): **Oral absorption of *Mesobuthus eupeus* scorpion venom in mice.** IJT, 11 (2): 15-19.
- 160- Ighodaro, O. M. and Akinloye, O. A. (2017): **First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid.** Alex. J. Med. (AJM), xxx (2017) xxx–xxx.
- 161- Ignarro, L. J. (2000): **Nitric Oxide: Biology and Pathobiology.** Sec. III: Principles of Pathobiology. Academic Press., Losangles, California, USA, P.454
- 162- Incamnoi, P.; Patramanon, R.; Thammasirirak, S.; Chaveerach, A.; Uawonggul, N.; Sukprasert, S.; Rungsa, P.; Daduang, J.; Daduang, S. (2013): **Heteromtoxin (HmTx), a novel heterodimeric phospholipase A2 from *Heterometrus laoticus* scorpion venom.** Toxicon, 61 (2013): 62–71.
- 163- Inceoglu, B.; Lango, J.; Jing, J.; Chen, L.; Doymaz, F.; Pessah, N. I. and Hammock, B. D. (2003): **One scorpion, two venoms: Prevenom of *Parabuthus transvaalicus* acts as an alternative type of venom with distinct mechanism of action.** Proc. Natl. Acad. Sci., U S A, 100 (3): 922–927.
- 164- Isbister, G. K.; Graudins, A.; White, J. and Warrell, D. (2003): **Antivenom treatment in arachnidism.** J. Toxicol. Clin. Toxicol., 41: 291-300.
- 165- Ismail M. (1995): **The scorpion envenoming syndrome.** Toxicon, 33(7): 825-58.
- 166- Ismail M. and Abd-Elsalam, M. A. (1988): **Are the toxicological effects of scorpion**

- envenomation related to tissue venom concentration? *Toxicon*, 26: 233-56.
- 167- Ismail M.; Fatani, A. J. Y. and Dabeas T. T. (1992): **Experimental treatment protocols for scorpion envenomation; a review of common therapies and on effect of kallikrein – kinin inhibitors.** *Toxicon*, 30: 1257-79.
- 168- Ismail, M.; Abd-Elsalam, M. A. and al-Ahaidib, M. S. (1995): ***Androctonus crassicauda (Olivier)*; a dangerous and unduly neglected scorpion-I: Pharmacological and clinical studies.** *Toxicon* 32 (12): 1599-618.
- 169- Jackson, E. K. (2006): **Renin and Angiotensin.** In: Laurence L. Brunton, John S, Lazo, Keith L. Parker, editors. Goodman & Gilman's: The Pharmacological Basis of Therapeutics, 11th edition, New York, New Delhi: McGraw-Hill Medical Publishing Division, pp. 795.
- 170-Jackson, M. J. (2000): **Exercise and oxygen radical production by muscle.** In: Sen, C. K., Packer, L. & Hanninen, O. (eds). Handbook of Oxidants and Antioxidants in Exercise, Amsterdam, Elsevier Science, 57-68.
- 171- Jalali, A. and Rahim, F. (2014): **Epidemiological Review of Scorpion Envenomation in Iran.** *Iranian Journal of Pharmaceutical Research*, 13 (3):743-756.
- 172- Jalali, A.; Pipelzadeh, M. H.; Taraz, M.; Khodadadi, A. Makvandi, M. and Rowan, E. G. (2011): **Serum TNF- α levels reflect the clinical severity of envenomation following a *Hemiscorpius lepturus* sting.** *Eur. Cytokine Netw.*, 22 (1): 5-10.
- 173- Jarrar,B. M. and Al-Rowaily, M. A. (2008): **Epidemiological aspects of scorpion stings in Al-Jouf Province, Saudi Arabia.** *Ann. Saudi Med.*, 28 (3): 183-187
- 174- Jeran, A. J. (2001): **Scorpion Biology and Research.** In: Brownell, P., and Polis, G. A. editors, Paleontology, New York: Oxford Univ., Press 2001, 370-92.
- 175- Ji, X.; Zhang, P.; Armstrong, R. N. and Gilliland, G. L. (1992): **The three dimensional structure of glutathione-S-transferase from the Mu gene class: Structural analysis of the binary complex of isozyme 3-3 and glutathione at 2.2 Å resolution.** *Biochem.*, 3131 (42): 10169-84.
- 176- Judah, V. and Nuttall, K. (2016): **Exotic Animal Care and Management.** Cengage Learning, USA, 2nd edition, 384 pages.
- 177- Kalam, S.; Singh, R.; Mani, A.; Patel, J.; Khan, F. N. and Pandey, A. (2012): **Antioxidants: Elixir of life.** . *J. Int. Multidiscipl. Res.*, 2 (1): 18-34.
- 178- Kale, M.; Rathore, N.; John, S. and Bhtangar, D. (1999): **Lipid peroxidative damage on pyrethroid exposure and alternations in antioxidants status in rat erythrocytes: A possible involvement of reactive oxygen species.** *Toxicol. Lett.*, **105 (3): 197-205.**
- 179- Kankonkar, R. C.; Murthy, K. R. K.;Zare, A. M.; Balasubramaniam, P. and Yeolekar, M. E. (1992): **Reversal of cardiovascular and hemodynamic disturbances by scorpion antivenin administration in myocarditis due to envenomation by Indian red scorpion (*Buthidae* family) venom.** *Rec. Adv. Toxinol. Res.*, 2: 61-70.

- 180- Karnad, D. R. (2009): **Management of scorpion envenomation: Need for a standard treatment protocol using drugs and antivenom.** J.Assoc. Phys. India, 57: 299–300.
- 181- Kassiri, H.; Kasiri, E.; Veys-Behbahani, R. and Kasiri, A. (2014): **Epidemiological survey on scorpionism in Gotvand County, Southwestern Iran: An analysis of 1067 patients.** J. Acute Disease, 2014: 314-319.
- 182- Keele, C. A.; Neil, E. and Joels, N. (2000): **Samson Wright’s Applied Physiology.** 13th edn., Oxford, Oxford University Press., pp.405–512.
- 183- Khatony, A.; Abdi, A.; Fatahpour, T. and Towhidi, F. (2015): **The epidemiology of scorpion stings in tropical areas of Kermanshah province, Iran, during 2008 and 2009.** Venom. Anim. Toxins. Incl. Trop. Dis., 21(1): 45-52.
- 184- Khorchid, A.; Fragoso, G.; Shore, G.; and Almazan, G. (2002): **Catecholamine-induced oligodendrocyte cell death in culture is developmentally regulated and involves free radical generation and differential activation of caspase-3.** GLIA, 40 (3): 283-99.
- 185- Kim, w. R.; Flamm, S. L.; Di Bisceglie, A. M. and Bodenheimer, H. C. (2008): **Serum activity of alanine aminotransferase (ALT) as an indicator of health and disease.** Hematology, 47(4): 1363-1370.
- 186- Kochar, N. I. and Umathe S. N. (2009): **Beneficial effects of L-arginine against diabetes-induced oxidative stress in gastrointestinal tissues in rats.** Pharmacology Reports, 61(4): 665-672.
- 187- Kohen, R. and Nyska, A. (2002): **Oxidation of biological systems: Oxidative stress phenomena, antioxidants, redox reactions, and methods for their quantification.** Toxicol. Pathol., 30(6): 620–50.
- 188- Koracevic, D.; Koracevic, G.; Djordjevic, V.; Andrejevic, S. and Cosic, V. (2001): **Method for the measurement of antioxidant activity in human fluids.** J. Clin. Pathol., 54 (5): 356-361.
- 189- Krifi, M. N.; Amri, F.; Kharrat, H. and El Ayeb, M., (1999): **Evaluation of antivenom therapy in children severely envenomed by *Androctonus australis garzonii* (Aag) and *Buthus occitanus tunetanus* (Bot) scorpions.** Toxicon, Nov., 37 (11), 1627-34.
- 190- Krifi, M. N.; Marrakchi, N.; El Ayeb, M. and Dellagi, K. (1998): **Effect of some variables on the in vivo determination of scorpion and viper venom toxicities.** Biologicals, 26: 277-88.
- 191- Krishnan, A.; Sonawane, R. V.; Karnad, D. R. (2007): **Captopril in the treatment of cardiovascular manifestations of Indian red scorpion (*Mesobuthus Tamulus Concanesis Pocock*) envenomation.** J. Assoc. Physicians Ind., 55: 22-26.
- 192- Krishnan, N. and Kodrík, D. (2006): **Antioxidant enzymes in *Spodoptera littoralis* (Boisduval): Are they enhanced to protect gut tissues during oxidative stress?** J. Insect. Physiol., 52 (1): 11-20.
- 193- Kshirsagar, V. Y.; Ahmed, M. A. and Colaco, S. M. (2012): **Motor aphasia: A rare complication of scorpion sting.** Pediatr. Neurosci., 7 (3): 231–3.

- 194- Laustsen, A. H.; Solà, M.; Jappe, E. C.; Oscoz, S.; Lauridsen, L. P. and Engmark, M. (2016): **Biotechnological trends in spider and scorpion antivenom development.** *Toxins (Basel)*, 8 (8): 226-258.
- 195- Lee, C.C.; Hsieh, H. J.; Hsieh, C. H. and Hwang, D.F. (2015): **Plancitoxin I from the venom of crown-of-thorns starfish (*Acanthaster planci*) induces oxidative and endoplasmic reticulum stress associated cytotoxicity in A375.S2 cells.** *Exp. Mol. Pathol.*, 99 (1): 7–15.
- 196- Lesser, M. P. (2006): **Oxidative stress in marine environments: Biochemistry and physiological ecology.** *Ann. Rev. Physiol.*, 68(1):253-78.
- 197- Lesser, M. P.; Levine, R. L.; Garland, D.; Oliver, C. N.; Amici, A.; Climent, I.; Lenz, A. G.; Ahn, B. W.; Shaltiel, S. and Stadtman, E. R. (1990): **Determination of carbonyl content in oxidatively modified proteins.** *Methods Enzymol.*, 186: 464-78.
- 198- Levine, R. L.; Garland, D.; Oliver, C. N.; Amici, A.; Climent I.; Lenz, A. G.; Ahn, B. W.; Shaltiel, S. and Stadtman, E. R. (1990): **Determination of carbonyl content in oxidatively modified proteins.** *Methods Enzymol.*, 186: 464-78.
- 199- Levine, R. L.; Williams, J. A.; Stadtman, E. R. and Shacter, E. (1994): **Carbonyl assays for determination of oxidatively modified proteins.** *Methods Enzymol.*, 233: 346-357.
- 200- Liang, S. and Juan, J. (2011): **Effect of jujube extract on oxidative injury in heart muscles of exhausted training rats.** *Afr. J. Micro. Res.*, 5 (14): 1896-1899.
- 201- Libby, P. and Hansson, G. K. (1991): **Biology of disease: Involvement of the immune system in human atherogenesis: Current knowledge and unanswered questions.** *Lab. Invest.*, 64 (1): 5-15.
- 202- Lighton, J. R. B.; Brownell, P. H.; Joos, B. and Turner, R. J. (2001): **Low metabolic rate in scorpions: implications for population biomass and cannibalism.** *J. of Exp. Biol.*, 204: 607–613.
- 203- Lobo, V.; Patil, A.; Phatak, A. and Chandra, N. (2010): **Free radicals, antioxidants and functional foods: Impact on human health.** *Pharmacogn. Rev.*, 4 (8): 118–126.
- 204- Lourenço, W. R. (2016): **Scorpion incidents, misidentification cases and possible implications for the final interpretation of results.** *J. Venom. Anim. Tox. Incl. Trop. Dis.*, 22: 21.
- 205- Love, S. (1991): **Oxidative stress in brain ischemia.** *Brain Pathol.*, 9 (1): 119-131.
- 206- Loven, D.; Schedl, H.; Wilson, H.; Daabees, T. T.; Stegink, L. D.; Diekus, M. and Oberley, L. (1986): **Effect of insulin and oral glutathione on glutathione levels and superoxide dismutase activities in organs of rats with streptozotocin-induced diabetes.** *Diabetes*, 5 (5): 503-7.
- 207- Lowe, R. M. and Farrell, P. M. (2011): **A portable device for the electrical extraction of scorpion venom.** *Toxicon*, 57 (2): 244-247.
- 208- Mahadevan, S. (2000): **Scorpion sting.** *Indian Pediatr.*, 37 (5): 504-514.
- 209- Mansour, N. (2001): **Waiting time and characteristics of taking in charge scorpion priced persons in Sidi-Bouzid region Institute Pasteur de Tunis, Tunis,**

- Tunisia. Archives de l'Institut de Pasteur, Tunis, 78: 25-31.
- 210- Maral, J.; Puget, K. and Michelson, A. (1977): **Comparative study of superoxide dismutase, catalase and glutathione peroxidase levels in erythrocytes of different animals.** Biochem. Biophys. Res. Commun., 77: 1525-1535.
- 211- Mc Cord, J. M. (2000): **The evolution of free radicals and oxidative stress.** Am. J. Med., 108 (8): 652-659.
- 212- McIntosh, M.; Ghomashchi, F.; Gelb, H.; Dooley, J.; Stoehr, J.; Giordani, B.; Naisbitt, R. and Olivera, M. (1995): **Conodipine-M; a novel phospholipase A2 isolated from the venom of marine cone snail *Conus magus*.** J. Biol. Chem., 270 (8): 3518-3526.
- 213- Meadows P. E. and Russell F. E. (1970): **Milking of arthropods.** Toxicon, 8: 311–312.
- 214- Mebs, D. (2002): **Scorpions and snakes, such as cobras, mambas and vipers made the African continent famous for venomous animals.** Bulletin de la Societe de Pathologie Exotique, 95 (3): 131.
- 215- Meier, J. and Theakston, R. D. (1986): **Approximate LD50 determinations of snake venoms using eight to ten experimental animals.** Toxicon, 24 (4): 395-401.
- 216- Meki A. R.; Hasan, H. A.; El-Deen, Z. M. And Bakkar, S. (2003): **Dysregulation of apoptosis in scorpion envenomed children: Its reflection on their outcome.** Toxicon, 42: 229-37.
- 217- Meki, A. R. and Mohey El-Dean, Z. M. (1998): **Serum interleukin-1-beta, 1L-6, interleukin-6, nitric oxide and alpha-1-antitrypsin in scorpion envenomed children.** Toxicon, 36 (12): 1851-1859.
- 218- Mirakabadi, A. Z.; Jalali A.; Jahromi A. E.; Vatanpur, H. and Akbary, A. (2006): **Biochemical changes and manifestations of envenomation produced by *Odonthobuthus doriae* venom in rabbits.** J. Venom. Anim. Toxins. Incl. Trop. Dis., 12 (1): 67-77.
- 219- Mistry P. and Herbert, K. (2003): **Modulation of hOGG1 NDA repair enzyme in human cultured cells in response to prooxidant and antioxidant challenge.** Free Rad. Biol. Med., 35: 397-405.
- 220- Moghadam, A. T.; Akbari, A.; Kavosh, F.; Mohammadian, A. and Soleimani, P. (2013): **Evaluation of the differences in protein profile, osmolarity and electrolyte composition of haemolymph of five species of scorpion in Khuzestan province (South-Western Iran).** J. Exp. Zool. India, 16 (2): 615-617.
- 221- Mohamad, I.; Elsayh, K.; Mohammad, H.; Saad, K.; Zahran, A. and Abdallah, A. (2014): **Clinical characteristics and outcome of children stung by scorpion.** Eur. J. Pediatr., 173: 815–818.
- 222- Mohamed, A. H.; Hani-Ayobem, M.; Mervat A.; Beskharoun, M. A. and El-Damarawy, N. A. (1972): **Glycaemic responses to scorpion venom.** Toxicon, 10 (2): 139-149.
- 223- Moncada, S.; Palmer, R. M. J. and Higgs, E. A. (1991): **Nitric oxide: Physiology, pathophysiology and pharmacology.** Pharmacol. Res., 43 (2): 109–142.

- 224- Morrow, S. (2017): **A safer way to milk a scorpion.**
<http://blogs.discovermagazine.com/d-brief>.
- 225- Moselhy, W. A. (2012): **Cytotoxicity and antimicrobial activity of the haemolymph peptides of the silk worm; *Bombyx mori***
L. J. Appl. Sci. Res., 8 (6): 3038-3043
- 226- Moustafa, S. R. (2015): **Association of superoxide dismutase, glutathione peroxidase, catalase, and xanthine oxidase with incidence of bladder cancer.**
J. Cancer Res., 3 (2): 17-27.
- 227- Murthy K. R. K. and Krishna, R. (2002): **On scorpion envenoming syndrome: Problems of medical ethics and accountability in medical research in India.** *J. Venom. Anim. Toxins*: 8 (1): 3–17.
- 228- Murthy, K. R. K. (2000): **The scorpion envenoming syndrome: A different perspective: The physiological basis of the role of insulin in scorpion envenoming.** *J. Venom. Anim. Toxins*, 6 (1): 4-51.
- 229- Murthy, K. R. K. and Haghazari, L. (1999): **The blood levels of glucagon, cortisol and insulin following the scorpion (*Mesobuthus tamulus concanesis*, Pocock) envenoming in dogs.** *J. Venom. Anim. Toxins*, 5 (1): 47-55.
- 230- Murthy, K. R. K. and Zare, A. Z. (2002): **Scorpion antivenom reverses metabolic, electrocardiographic, and hormonal disturbances caused by the Indian red scorpion *Mesobuthus tamulus concanesis*, Pocock Envenomation.** *J. Venom. Anim. Toxins Incl. Trop. Dis.*, 8: 30-48.
- 231- Murthy, K. R. K.; As, D.; M. Zare Abbas, Z. and Haghazari, L. (2003): **Investigations on the role of insulin and scorpion antivenom in scorpion envenoming syndrome.** *J. Venom. Anim. Toxins incl. Trop. Dis.*, 9 (2): 202-38
- 232- Murthy, K. R.; Shenoy, R.; Vaidyanathan, P.; Kelkar, K.; Sharma, N.; Birevar, N.; Rao, S. and Mehta, M. N. (1991): **Insulin reverses haemodynamic changes and pulmonary edema in children stung by the Indian red scorpion *Mesobuthus tamulus concanesis* Pocock.** *Ann. Trop. Med. Parasitol.*, 85: 651–657.
- 233- Natu, V. S.; Kamerkar, S. B.; Geeta, K.; Vidya, K.; Natu, V.; Sane, S.; Kushte, R.; Thatte, S.; Uchil, D. A.; Rege, N. N. and Bapa, R. D. (2010): **Efficacy of anti-scorpion venom serum over prazosin in the management of severe scorpion envenomation.** *J. Postgrad. Med.*, 56 (4): 275-80.
- 234- Natu, V. S.; Kumar, Z. N. and Lavanya, M. (2015): **Suppressed insulin secretion, elevated mediators of inflammation, hyperinsulinemia-insulin resistance: Insulin administration reverses cardiovascular, metabolic changes, pulmonary edema and all other clinical manifestations in scorpion envenoming syndrome.** *Ind. J. Mednodent. Allied. Sci.*, 3 (2): 90-107
- 235- Nebauer, A. (2013): **Keeping *Androctonus sp.* in captivity.**
<http://thereptiletimes.wordpress.com>.
- 236- Newton, M. (2007): **Help!! How do I sex my scorpion?** <http://arachnoboards.com>.

- 237- Nisani, Z.; Dunbar, S. G. and Hayes, W. K. (2007): **Cost of venom regeneration in *Parabuthus transvaalicus* (Arachnida: Buthidae)**. Comp. Biochem. Physiol., 147 (2): 509-513.
- 238- Ohkawa, H. W.; Ohishi, N. and Yagi, K. (1979): **Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction**. Anal. Biochem., 95: 351-358.
- 239- Olivera, B. M. and Teichert, R. W. (2007): **Diversity of the neurotoxic conus peptides: A model for concerted pharmacological discovery**. Mol. Interv., 7 (5): 251-60.
- 240- Olsen, F. (1972): **Inflammatory cellular reaction in hypertensive vascular disease in man**. Acta Pathol. Microbiol. Scand. (A), 80 (2): 253-6.
- 241- Omran, M. A. A. (2003): **Cytotoxic and apoptotic effects of scorpion *Leiurus quinquestriatus* venom on 293T and C₂C₁₂ eukaryotic cell lines**. J. Venom. Anim. Toxins Incl. Trop. Dis., 9 (2): 255-275.
- 242- Omran, M. A. A. and Mc Vean, A. (2000): **Intraspecific variation in scorpion *Leiurus quinquestriatus* venom collected from Egypt (Sinai and Aswan deserts)**. J. Toxicol. Toxin Rev., 19: 247-64.
- 243- ORKIN (2017): **Are scorpions poisonous?** <https://www.orkin.com/stinging-pests/scorpions/poisonous-scorpions>.
- 244- ORKIN (2017): **Scorpions: Facts, Identification and Control**. <https://www.orkin.com/stinging-pests/scorpions>.
- 245- Oukkache, N.; Chgoury, F.; Lalaoui, M.; Cano, A. A. and Ghalim N. (2013): **Comparison between two methods of scorpion venom milking in Morocco**. J. Venom. Anim. Toxins incl. Trop. Dis., 19 (5): 1-5.
- 246- Özkan, O. and Filazi, A. (2004): **The determination of acute lethal dose-50 (LD₅₀) levels of venom in mice, obtained by different methods from scorpions; *Androctonus crassicauda* (Olivier 1807)**. Acta Parasitol. Turcica., 28 (1): 50–53.
- 247- Özkan, O.; Adiguzel, S.; Kar, S.; Yakistiran, S.; Cesaretli, Y. and Karaer, K. Z. (2007): **Determination of potency and paraspecific effects of *Androctonus crassicauda* (Olivier, 1807) antivenom against *Mesobuthus gibbosus* (Brullé, 1832) venom (Scorpiones: Buthidae)**. J. Venom. Anim. Toxins incl. Trop. Dis., 13 (2): 500-508.
- 248- Özkan, O.; Adigüzel, S; Ates, C.; Bozyigit, I. and Filazi, A. (2006): **Optimization of anti-scorpion venom production**. J. Venom. Anim. Toxins Incl. Trop. Dis., 12 (3): 390-399
- 249- Padmanabhan, P. K.; Krishnan, R. G.; Kalimuthu, K.; Perumal, D.; Elumalai, S.; Arulappan, J. P. and Arumugam Perumal, A. (2012): **In vitro antioxidant activity of haemolymph from *Camponotus compressus* and its antihyperglycemic activity on alloxan Induced diabetic albino rats**. Asian Pac. J. Trop. Biomed., (2012): 1-6
- 250- Paglia, D. E. and Valentine, W. N. (1967): **Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase**. J. Lab. Clin. Med., 70: 158-169.

- 251- Pandey, K. B. and Rizvi, S. I. (2010): **Resveratrol may protect plasma proteins from oxidation under conditions of oxidative stress in vitro.** J. Braz. Chem. Soc., 21 (5): 909-913.
- 252- Patlevič, P.; Vašková, J.; Jr., P. S.; Vaško, L. and Švorc, P. (2016): **Reactive oxygen species and antioxidant defense in human gastrointestinal diseases.** Integr. Med. Res., 5 (4): 250–258.
- 253- Peña, L. M.; Dolores, F. and Peña, J. P. (2003): **Fish tolerance to organophosphate-induced oxidative stress is dependent on the glutathione metabolism and enhanced by N-acetyl cysteine.** Aquat. Toxicol., 65: 337-360.
- 254- Petricevich, L. (2004): **Cytokine and nitric oxide production following severe envenomation.** Current Drug Targets- Inflammation and Allergy, 3 (3): 325-332.
- 255- Petricevich, V. L. (2002): **Effect of *Tityus serrulatus* venom on cytokine production and the activity of murine macrophages.** Mediat. Inflamm., 11 (1): 23-31.
- 256- Petricevich, V. L. (2010): **Scorpion venom and the inflammatory response.** Mediat. infl., 2010: 903295.
- 257- Petricevich, V. L. and Peña, C. F. (2002): **The dynamics of cytokine and nitric oxide secretion in mice injected with *Tityus serrulatus* scorpion venom.** Mediators of Inflammation, 11 (3):173–180
- 258- Polis, G. A. (1990): **The biology of scorpions.** Stanford Univ., Pr., 10th edn., 587 pages.
- 259- Possani, E.; Merino, M.; Corona, F.; Bolivar, F. and Becerril, B. (2000): **Peptides and genes coding for scorpion toxins that affect ion-channels.** Biochim., 82 (9-10): 861-868.
- 260- Prendini, L. and Wheeler, W. C. (2005): **Scorpion higher phylogeny and classification, taxonomic anarchy, and standards for peer review in online publishing.** Cladistics, 21: 446 - 494
- 261- Pryor, W. A.; Houk, K. N.; Foote, C. S.; Fukuto, J. M.; Ignarro, L. J.; Squadrito, G. L. and Davies, K. J. (2006): **Free radical biology and medicine: It's a gas, man!** Am. J. Physiol. Regul. Integr. Comp. Physiol., 291 (3): R491- 511.
- 262- Pucca, M. B.; Carlos, J.; Roncolato, E. C.; Bertolini, T. B.; Fossa, C. M.; Varanda, W. A.; Arantes, E. C. and Barbosa, J. E. (2011): **Monoclonal antibody (Scfv) against the venom of the scorpion *Tityus serrulatus*; produced by phage display technic, is capable to recognize and inhibit the action of the ts1 toxin.** Epeq/Fafibe, 1: 18–23.
- 263- Quéinnec, E.; Gardès-Albert, M.; Goyffon, M.; Ferradini, C. and Vuillaume, M. (1999): **Antioxidant activity of haemocyanin: A pulse radiolysis study.** Biochim. Biophys. Acta(BBA), 1041 (2): 153-159.
- 264- Quintero-Hernández, V.; Jiménez-Vargasa, J. M.; Gurrolaa, G. B.; Valdiviab, H. H. F. and Possani, L. D. (2013): **Scorpion venom components that affect ion-channels function.** Toxicol., 15 (76): 328–342.
- 265- Rakhit, R. D.; Mojat, M. H.; Marber, M. S. and Duchon, M. R. (2001): **Mitochondria as targets for nitric oxide-induced protection during simulated ischemia and reoxygenation in isolated neonatal**

- cardiomyocytes.** Circulation, 103 (21): 2617-2623.
- 266- Raynova, Y.; Marchev, A.; Doumanova, L.; Pavlov, A.; Idakieva, K. (2015): **Antioxidant Activity of *Helix aspersa maxima* (Gastropod) Hemocyanin.** Acta microbiologica Bulgarica, 31 (2): 127-131.
- 267- Repetto, M.; Semprine, J. and Boveris, A. (2012): **Lipid Peroxidation: Chemical Mechanism, Biological Implications and Analytical Determination.** Tech Open Access Pub., Argentina, Chapter 1.
- 268- Reznick, A. Z. and Packer, L. (1994): **Oxidative damage to proteins: A spectrophotometric method for carbonyl assay.** Methods Enzymol., 233: 357-363.
- 269- Rodrigo, C. and Gnanathasan, A. (2017): **Management of scorpion envenoming: A systematic review and meta-analysis of controlled clinical trials.** Rodrigo and Gnanathasan Systematic Reviews, 6:74-85.
- 270- Rodríguez, E. R. R.; Umbarila, L. R.; Possani, L. D. and Becerril B. (2015): **Recombinant neutralizing antibodies: A new generation of antivenoms.** Part of the Toxinology book series: Scorpion Venoms. TOXI, volume 4, Springer; Berlin, Germany, pp. 139-159.
- 271- Rodríguez-Rodríguez, E. R.; Olamendi-Portugal, T.; Serrano-Posada, H.; Arredondo-López, J. N.; Gómez-Ramírez, I.; Fernández-Taboada, G.; Possani, L. D.; Anguiano-Vega, G. A.; Riaño-Umbarila, L. and Becerril, B. (2016): **Broadening the neutralizing capacity of a family of antibody fragments against different toxins from Mexican scorpions.** Toxicon, 119: 52-63.
- 272- Rosselli, M.; Keller, P. J. and Dubey, R. K. (1998): **Role of nitric oxide in the biology, physiology and pathophysiology of reproduction.** Hum. Reprod. Update., 4 (1): 3-24.
- 273- Sagheb, M. M.; Sharifian, M.; Moini, M. and Salehi, O. (2011): **Acute renal failure and acute necrotizing pancreatitis after *Echis carinatus sochureki* bite: Report of a rare complication from southern Iran.** Prague. Med. Rep., 12 (1): 67-71
- 274- Sahan-Firat,, S.; Canacankatan, N.; Korkmaz, B.; Yildirim, H. and Tamer, L. (2012): **Increased production of nitric oxide mediates selective organspecific effects of endotoxin on oxidative stress.** Antiinflamm. Antiallergy Agents Med. Chem., 11: 161-172.
- 275- Santamaria, A.; Rodriguez, S.; Zugasti, A.; Martinez, A.; Galvan-Arzate, S. and S. Puertas, (2002): **A venom extract from the sea anemone *Bartholomea annulata* produces hemolysis and lipid peroxidation in mouse erythrocytes.** Toxicology, 173 (3): 221-228.
- 276- Santibáñez-López, C. E.; Francke, O. F.; Ureta, C. and Possani, L. D. (2016): **Scorpions from Mexico: From species diversity to venom complexity.** Toxins, 8 (1): 2-19.
- 277- Satoh, H. and Nishida, S. (2004): **Electropharmacological actions of *Ginkgo biloba* extract on vascular smooth and heart muscles.** Clin. Chim. Acta, 342 (1-2): 13-22.
- 278- Satoh, K. (1978): **Serum lipid peroxides in cerebrovascular disorders detected by a**

- new colorimetric method. Clin. Chim. Acta, 90: 37-43.
- 279- Schild, L.; Reinheckel, T.; Reiser, M.; Horn, T. F.; Wolf, G. and Augustin W. (2003): **Nitric oxide, produced in rat liver mitochondria, causes oxidative stress and impairment of respiration after transient hypoxia.** FASEB J., 17 (15): 2194-2201.
- 280- Selmane, S.; Benferhat, L.; L'Hadj, M. and Zhu, H. (2017): **Scorpionism in Sidi Okba, Algeria: A cross-sectional study of 2016 stung patients between 2014 and 2015.** Trop. Biomed., 34 (2): 425-432.
- 281- Shacter, E. (2000): **Quantification and significance of protein oxidation in biological samples.** Drug. Metab. Rev., 32 (3-4): 307-26
- 282- Shimizu, T.; Ohto, T. and Kita, Y. (2006): **Cytosolic phospholipase A2: Biochemical properties and physiological roles.** IUBMB Life, 58 (5-6): 328-333.
- 283- Shrestha, B. R.; Pandey, D. P.; Acharya, K. P.; Thapa-Magar, C.; Mohamed, F. and Isbister, G. K. (2017): **Effective, polyvalent, affordable antivenom needed to treat snakebite in Nepal.** Bulletin of the World Health Organization, 95: 718-719
- 284- Slater, T. F. (1984): **Free-radical mechanisms in tissue injury.** Biochem. J., 222 (1): 1-15.
- 285- Snedecor, G. W. (1956): **Statistical Methods.** Ames, Iowa, USA, Iowa State University Press, 4th edn.
- 286- Soleglad, M. E. and Fet, V. (2003): **High-level systematics and phylogeny of the extant scorpions (*Scorpiones: Orthosterni*).** Euscorpius, 11: 1-175.
- 287- Sollod, B. L.; Wilson, D.; Zhaxybayeva, O.; Gogarten, J. P.; Drinkwater, R. and King, G. F. (2005): **Were arachnids the first to use combinatorial peptide libraries?.** Peptides, 26 (1): 131-39.
- 288- Stadtman, E. R. (2004): **Role of oxidant species in aging.** Cur. Med. Chem., 11 (9): 1105-1112.
- 289- Stadtman, E. R. and Berlett, B. S. (1997): **Reactive oxygen-mediated protein oxidation in aging and disease.** Chem. Res. Toxicol., 10 (5):485-94.
- 290- Stadtman, E. R. and Berlett, B. S. (1998): **Reactive oxygen-mediated protein oxidation in aging and disease.** Drug Metab. Rev., 30 (2): 225-43.
- 291- Stamler, J. S.; Singels, D. J. and Loscalzo, J. (1992): **Biochemistry of nitric oxide and its redox activated forms.** Science, 258 (5090): 1898 -1902.
- 292- Stamler, J.S.; Simon, D.; Osborne, J. A.; Mullins, M. E.; Jaraki, O.; Michel, T.; Singel, D. J. and Loscalzo, J. (1992): **S-nitrosylation of proteins with nitric oxide: Synthesis and characterization of biologically active compounds.** Proc. Natl. Acad. Sci., 89 (1): 444-448.
- 293- Stanley, K. (2000): **Nitric Oxide and Free Radicals in Peripheral Neurotransmission.** Springer Science and Business Media, 1st edn., New York.
- 294- Sunagar, K.; Undheim, E. A. B.; Chan, A. H. C; Koludarov, I.; Muñoz-Gómez, S. A.; Antunes, A.; and Fry, B. G. (2013): **Evolution stings: The origin and diversification of scorpion toxin peptide scaffolds.** Toxins, 5 (12): 2456-2487.
- 295- Tan, P. T.; Veeramani, A.; Srinivasan, K. N.; Ranganathan, S.; Brusica, V. (2006):

- Scorpion 2: A database for structure-function analysis of scorpion toxins.** Toxicon, 47 (3): 356 – 363.
- 296- Telmisany, A. K. and Othman, O. H. (1993): **Snake and scorpion venom and envenomation: A review.** JKAU., Med. Sci., 3: 11-18.
- 297- Teruel, R.; Kovařík, F. and Turiel, C. (2013): **A new species of *Androctonus Ehrenberg, 1828* from north western Egypt (*Scorpiones: Buthidae*).** Euscorpius: Occasional Publications in Scorpiology, 177.
- 298- Thatoi H. N.; Patra, J. K. and Das, S. K. (2014): **Free radical scavenging and antioxidant potential of mangrove plants: A review.** Acta Physiol. Plant, 36: 561–579.
- 299- Todos Santos Eco Adventures' Blog; TSEA (2011): **Baby love: Scorpion motherhood and family values.** <https://tosea.wordpress.com>.
- 300- Trinder, P. (1969): **Determination of glucose in blood using glucose oxidase with an alternative oxygen receptor.** Ann. Clin. Biochem., 6: 24-7.
- 301- Tripod.com (2018): **Venom and toxin: What to do when stung by a scorpion?** <http://members.tripod.com/~cianwee/welcome.html>.
- 302- Uawonggul, N.; Chaveerach, A.; Thammasirirak, S.; Arkaravichien, T.; Chuachan, C. and Daduang, S. (2006): **Screening of plants acting against *Heterometrus laoticus* scorpion venom activity on fibroblast cell lysis.** J. Ethnopharmacol., 103 (2): 201–207.
- 303- Uawonggul, N.; Sukprasert, S.; Rungsa, P.; Daduang, J. and Daduang, S. (2012): **Heteromtoxin (HmTx); a novel heterodimeric phospholipase A2 from *Heterometrus laoticus* scorpion venom.** Toxicon, 61 (1) 62–71.
- 304- Uchida K. (2000): **Role of reactive aldehyde in cardiovascular diseases.** Free Rad. Biol. Med., 28 (12): 1685 - 1696.
- 305- Uluğ, M.; Yaman, Y.; Yapıcı, F. and Can-Uluğ, N. (2012): **Scorpion envenomation in children: An analysis of 99 cases.** Turkish J. Pediatrics, 54: 119-127.
- 306- Valdez-Cruz, N. A.; Batista, C. V. and Possani, L.D. (2004): **Phaiodactylipin; a glycosylated heterodimeric phospholipase A from the venom of the scorpion *Anuroctonus phaiodactylus*.** Eur. J. Biochem., 271 (8): 1453-1464.
- 307- Valenta, J.; Stach, Z. and Svítek, M. (2010): **Acute pancreatitis after viperid snake *Cerastes cerastes* envenoming: A case report.** Prague. Med. Rep., 111: 69–75.
- 308- Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M.; Mazur, M. and Telser, J. (2007): **Free radicals and antioxidants in normal physiological functions and human disease.** Int. J. Biochem. Cell Biol., 39 (1): 44-84.
- 309- Valko, M.; Morris, H. and Cronin, M. T. (2005): **Metals, toxicity and oxidative stress.** Curr. Med. Chem., 12 (10): 1161–208.
- 310- Vasconcelos, F.; Sampaio, S. V.; Garófalo, M. A. R.; Guimarães, L. F. L Giglio, J. R. and Arantes, E. C. (2004): **Insulin-like effects of *Bauhinia forficata* aqueous extract upon *Tityus serrulatus* scorpion envenoming.** J. Ethnopharmacol., 95 (2-3): 385-392.

- 311- Venancio, E. J.; Portaro, F. C.; Kuniyoshi, A. K.; Carvalho, D. C.; Piddé-Queiroz, G. and Tambourgi, D. V. (2013): **Enzymatic properties of venoms from Brazilian scorpions of *Tityus* genus and the neutralisation potential of therapeutical antivenoms.** *Toxicon*, 69 (2013):180-90.
- 312- Victor, V. M.; Rocha, M. and De la Fuente, M. (2004): **Immune cells: Free radicals and antioxidants in sepsis.** *Int. Immunopharmacol.*, 4 (3): 327–47.
- 313- Waddington, J.; Rudkin, D. M. and Dunlop, J. A. (2015): **A new mid-Silurian aquatic scorpion: One step closer to land?** *Biol. Lett.*, 11 (1): 20140815.
- 314- Weber, D.; Davies, M. J. and Grune, T. (2015): **Determination of protein carbonyls in plasma, cell extracts, tissue homogenates, isolated proteins: Focus on sample preparation and derivatization conditions.** *Redox Biol.*, 5: 367–380.
- 315- Wetscher, G. J.; Bagchi, M.; Bagchi, D.; Perdakis, G.; Hinder, P. R.; Glaser, K. and Hinder, R. A. (1995): **Free radicals production in nicotine-treated pancreatic tissue.** *Free Rad. Biol. Med.*, 18 (5): 877-882.
- 316- Whittemore, F. W.; Keegan, Jr. H. L.; Fitzgerald, C. M.; Bryant, H. A. and Flanigan, J. F. (1963): **Studies of scorpion antivenins: Venom collection and scorpion colony maintenance.** *Bull. World Health Organ.*, 28 (4): 505–511.
- 317- Wikipedia (2018): **Black fat-tailed scorpion.** Wikipedia; the free encyclopedia, <https://en.wikipedia.org/wiki>.
- 318- Williams, D. J. (2015): **Snake bite: A global failure to act costs thousands of lives each year.** *BMJ*, 351: h5378.
- 319- Winterbourne, C. C.; Hawkins, R. E.; Brian, M. and Carrell, R. W. (1974): **The estimation of red cells super-oxide dismutase activity.** *J. Lab. Clin. Med.*, 85: 337-342.
- 320- Witko-Sarsat, V.; Friedlander, M.; Capeillère-Blandin, C.; Nguyen-Khoa, T.; Nguyen, A. T.; Zingraff, J.; Jungers, P. and Descamps-Latscha, B. (1996): **Advanced oxidation protein products as a novel marker of oxidative stress in uremia.** *Kidney Int.*, 49 (5): 1304-1313.
- 321- Wu, X. S.; Jian, X. C.; Yin, B. and He, Z.J. (2010): **Development of the research on the application of chlorotoxin in imaging diagnostics and targeted therapies for tumors.** *Chin. J. Cancer*, 29 (6): 626 – 630.
- 322- Yamasaki, S. C.; Villaruel, J. S.; Barone, J. M.; Zambotti-Villela, L. and Silveira, P. F. (2008): **Amino peptidase activities, oxidative stress and renal function in *Crotalus durissus terrificus* envenomation in mice.** *Toxicon*, 52: 445 - 454.
- 323- Yaqoob, R.; Tahir, H. M.; Arshad, M.; Naseem, S. and Ahsan, M. M. (2016): **Optimization of the conditions for maximum recovery of venom from scorpions by electrical stimulation.** *Pakistan J. Zool.*, 48 (1): 265-269.
- 324- Yilmaz, N. F.; Arslan, E. D.; Demir, A.; Kavalci, C.; Durdu, T.; Yilmaz, M. S.; Yel, C. and Akbulut, S. (2013): **Epidemiologic and clinical characteristics and outcomes of scorpion sting in the southeastern**

- region of Turkey.** Ulus. Travma. Acil. Cerrahi. Derg., 19 (5): 417–422.
- 325- Yin, H.; Xu, L. and Porter, N. A. (2011): **Free radical lipid peroxidation: Mechanisms and analysis.** Chemical Reviews, 111 (10): 5944 - 5972.
- 326- Yu, B. P. (1999): **Approaches to anti-aging intervention: The promises and the uncertainties mechanisms of aging and development.** J. Biochem., 111 (2): 73-907.
- 327- Yugandhar, B.; Murthy, K. R. and Sattar, S. A. (1999): **Insulin administration in severe scorpion envenoming.** J. Venom. Anim. Toxins, 5: 200 - 219.
- 328- Zamudio, F. Z.; Conde, R.; Arevalo, C.; Becerril, B.; Martin, B. M.; Valdivia, H. H. and Possani, L. D. (1997): **The mechanism of inhibition of ryanodine receptor channels by imperatoxin I; a heterodimeric protein from the scorpion *Pandinus imperator*.** J. Biol. Chem., 272 (18):11886-94.
- 329- Zare, M. A. (1992): **Scorpion venom poisoning: Reversal of biochemical hormonal and pathophysiological disturbances by antivenom and insulin therapy.** PhD Thesis, University of Bombay, Bombay, India, pp229 .
- 330- Zargan, J.; Tirgari, S.; Tahernejad, K.; Lotfi, H. and Farahmandzad, A. (2003): **Study of scorpion fauna in Abomosa, great and small tonbs and hengam islands of the Persian Gulf.** Iran South Med J., 6 (1): 20 - 24.
- 331- Zeng, X.; Wang, S.; Nie, Y.; Zhanga, L. and Luo, X. (2012): **Characterization of BmKbpps; a multifunctional peptide from the Chinese scorpion *Mesobuthus martensii* Karsch: gaining insight into a new mechanism for the functional diversification of scorpion venom peptides.** Peptides, 33: 44 - 51.
- 332- Zengin, S.; Al, B.; Yarbil, P.; Guzel, R.; Orkmez, M.; Yildirim, C. and Taysi, S. (2014): **An assessment of oxidant/antioxidant status in patients with snake envenomation.** Emerg. Med. J., 31 (1): 48-52.
- 333- Zhao, R.; Dai, H.; Qiu, S.; Li, T.; He, Y.; Ma, Y.; Chen, Z.; Wu, Y.; Li, W. and Cao, Z. (2011): **SdPI; the first functionally characterized kunitz-type trypsin inhibitor from scorpion venom.** PLoS ONE, 6: e27548.
- 334- Zhu, S.; Peigneur, S.; Gao, B.; Luo, L.; Jin, D.; Zhao, Y. and Tytgat, J. (2011): **Molecular diversity and functional evolution of scorpion potassium channel toxins.** Mol. Cell Proteomics, 10 (2): M110.002832.