




BioBacta



Journal of Bioscience and Applied Research

www.jbaar.org



Knowledge, Attitude and Practice toward Prevention of Hepatitis B Virus Infection among Somalian Immigrant in the State of Selangor, Malaysia and their HBV Infection Status.

Abdirauf Mohamed Abdi^a, Mohd Nazil Salleh^{a*}

^aDepartment of Health Sciences, Faculty of Engineering and Life Sciences, Universiti Selangor, Shah Alam City Campus, Jalan Zikron A 7/A, Seksyen 7, 40000 Shah Alam Selangor, Malaysia

*Corresponding author: Associate Professor Dr. Mohd Nazil Salleh, Department of Health Sciences, Faculty of Engineering and life Sciences, Universiti Selangor, Shah Alam City Campus, Jalan Zikron A 7/A, Seksyen 7, 40000 Shah Alam Selangor, Malaysia. Email: drnazil@unisel.edu.my. Tel: +603 -5522 3471

Abstract

Background: Hepatitis B virus (HBV) infection is a serious health problem that can lead to liver cirrhosis and cancer. Assessing the public knowledge, attitude and practice towards HBV infection can be useful in planning public health policies in HBV management. **Objective:** The current study sought to determine the association between knowledge, attitude, and practice towards HBV infection with the HBV infection status of Somalian immigrants in Malaysia. **Method:** A cross-sectional study was conducted from September 2017 to April 2018 among Somalian immigrants in the state of Selangor, Malaysia. Data was collected using validated, self-administered structured questionnaire. Their blood samples was collected for detection of HBV DNA using nested PCR. Descriptive statistics and chi square analysis were done to determine this association. **Result:** A total of 145 participants were recruited. Majority of participants were in the age group of 20-31 (84.1%), male (62.8%), single (52.4%), had educational level of undergraduate degree (70.3%), and unemployed (88.3%). Majority of the participants show good knowledge (82.8%) and attitude (78.6%) but generally poor practice (32.4%) towards HBV infection. The mean knowledge, attitude and practice score among them were $16.9 \pm (4.89)$, $9.20 \pm (2.94)$ and $9.39 \pm (2.93)$ respectively. The sociodemographic characteristic variables and KAP of HBV were not significantly associated. Selected samples collected from the participant tested negative for HBV DNA via nested PCR assay. **Conclusion:** Somalian immigrants in Malaysia have a good level of knowledge and attitude but poor level of practice towards HBV infection. All participants were negative for HBV infection.

Key Words: Hepatitis B virus (HBV), Prevention, Somalian Immigrant, Infection Status, State of Selangor.

1. Introduction

Hepatitis B virus (HBV) Infection is one of the major public health issues worldwide. It is the most common and serious form of infection that can progress to chronic, lifelong infection of the liver (Kim et al. 2019). According to the World Health Organization (WHO), nearly 2 million people has been reported with evidence of ongoing or previous HBV infection globally. More than 2 million people, 350 million of them suffered from chronic, lifelong infection of the virus (Abdela et al. 2016).

HBV infection targeted the liver, leading to development of a plethora of liver diseases ranging from acute hepatitis to chronic hepatitis, liver cirrhosis, and hepatocellular carcinoma. It is estimated that 15–40% of chronic HBV carriers were susceptible to develop liver cirrhosis and hepatocellular carcinoma. Consequently, one million people die each year as a result of HBV infection and its complications (Noubiap et al. 2015).

Attributed to the development of vaccine against HBV, regions of developed nations such as North America, Northern and Western Europe and Australia have seen significant reduction in the disease burden (Baig et al. 2015). These regions are considered as having low endemicity of HBV with prevalence ranging from 5 to 7% of the population and 0.5 to 2% of them are chronic carrier (Baig VN et al. 2015).

However, moderate endemic regions such as part of Eastern and Southern Europe, the Middle East, Japan, and part of South America with 10 to 60% prevalence and 2 to 7% chronic carriers, as well as high endemic regions such as South East Asia, China, sub-Saharan Africa and the Amazon Basin with 70 to 95% prevalence and more than 8% chronic carriers, still exists due the lack of

enforcement and awareness towards the control of the disease (Coppola et al. 2015).

HBV can be transmitted through contact with infected bodily fluids that include blood, semen, and vaginal secretion (Sagnelli et al. 2014). Moreover, HBV can also be transmitted from infected mothers to their newborn babies and is believed to be a major contributor to the prevalence in high endemic regions (Adjei et al. 2016). In terms of practice, HBV is transmitted through unprotected sexual intercourse with infected individual, sharing of needles and syringes either in health-care settings or among persons who inject drugs, or during medical, surgical, or dental procedures, or getting a tattoo, or shaving in the barbershop through the use of equipment that is contaminated with infected blood (Sagnelli et al. 2014).

By adhering to universal precautions which include using protective barriers such as gloves, vaccination, appropriate sterilization of medical equipment, and proper hospital waste management system, the spread of HBV infection can be prevented. In addition, HBV cannot be transmitted by holding hands, sharing food, kissing, hugging, coughing, sneezing, or breastfeeding (Sagnelli et al. 2014).

Assessment of knowledge, attitude, and practice (KAP) towards a certain health issue is a useful tool in planning public health policies. Strategies in disease management can be tailored to maximize result by taking into account the current KAP of the target population. It is a valuable and most frequently used study tool in health-seeking behavior research. The study collects information on what is known, believed and done by the member of the population in relation to a particular topic (Hamissi et al. 2014).

Knowledge is usually assessed in order to see how

far the community members knew regarding the biomedical concepts revolve around the topic of interest such as causes and symptoms of the disease. When the knowledge deviates from biomedical concepts, it is termed as “beliefs”. People’s beliefs, feelings, and values toward a particular object or phenomenon culminated into their attitude towards it (Hassan-kadle et al. 2018). Finally, the knowledge and attitude towards a disease influence their practices such as preventive measures towards a certain disease or lifestyle modification (Kim et al. 2019).

HBV can also spread via immigration, particularly within migrants from countries with a high HBV endemicity to those with a low HBV endemicity (Li et al. 2015). Like many other sub-Saharan African countries, Somalia also have high endemicity of HBV with evidence of current and past infection found in 18.9% to 29% of the population (Mansour-Ghanaei et al. 2013). On the other hand, Malaysia, despite being one of the countries located in highly endemic region of South East Asia, become one of the countries in Asia who has reduced burden of HBV infection with reported HBV prevalence of 5% (Mesfin and Kibret 2013). The aim of the current study is to determine the association between knowledge, attitudes and practices (KAP) towards HBV infection among Somalian immigrants in the state of Selangor, Malaysia, and their HBV infection status.

2. Methods

Ethical Approval

This research has been approved by the Ethic Committee of the Faculty of Engineering and Life Science UNISEL (J150056E).

Study Design

Data on knowledge, attitude and practice toward HBV among Somalian immigrants in Malaysia and their sociodemographic characteristics were

assessed through validated, self-administered structured questionnaires (Nelson et al. 2017). HBV infection status of the participants were confirmed via detection of the HBV DNA in their blood (Rajamoorthy et al. 2019). Data were collected from September 2017 to April 2018.

Study Participants

The study population were recruited from Somalian immigrants to Malaysia who reside in the state of Selangor, particularly the three different regions of Shah Alam, Serdang and Gombak. Inclusion criteria include adults with Somalia citizenship in the state of Selangor that is older than 20 years old regardless of gender, employment, and social status. Somalian immigrants that has recurrent evidence of HBV infection, who are younger than 20 years old, that have citizenship other than Somalia, or not residing in Malaysia were excluded from the study.

Data Collection

Data was collected by using a self-administered structured questionnaire that collects information about the sociodemographic characteristics of the participants and their KAP towards prevention HBV. Data were collected by means of personal interview with the study population. The investigator gave a brief introduction to participants before filling the questionnaire. The face validity and the reliability of the questionnaire were evaluated by conducting a pilot study with approximately 5% of the sample size (10 people).

Blood Sample Collection

To determine the HBV infection status among participants, their blood was tested for the presence of HBV DNA. Venipuncture was performed by the certified phlebotomist and 4 ml of blood was collected in vacutainer tube together with age and gender data. All blood sample was centrifuged and the plasma was separated. All samples were stored at -80°C until analysis.

Nested Polymerase Chain Reaction

Presence of HBV DNA in participants blood was detected using nested polymerase chain reaction (PCR). Nested PCR is a technique that utilized two sets of primers to amplify low occurring DNA sequence. DNA from the samples was extracted

using a QIAamp DNA Blood Mini Kit (QIAGEN GmbH, Hilden, Germany). Two different sets of primers (S1 and nested S2) were used to target the HBV DNA as shown in Table 1. Presence of amplified product were assessed with agarose gel electrophoresis.

Table 1. Primers used for amplification of HBV infection.

Primer Name	Sequence (5'–3')
S1 forward	TCCTGCTGGTGGCTCCAG
S1 reverse	CGTTGACATACTTTCCAATCAA
Nested S2 forward	ACCCTGYRCCGAACATGGA
Nested S2 reverse	CAACTCCCAATTACATARCCCA

Data Analysis

Data analysis was checked for comprehension and reliability. Data was entered and analyzed using IBM SPSS software version 21.0. The findings were presented as numbers and percentages for categorical data. For quantitative data, the findings were presented as mean and standard deviation. The relationship between the variables were determined using descriptive statistics. Chi-square test was used for determine the effect of different factors on the event of HBV infection. The chi-square result with a p-value less than 0.05 was considered statistically significant and while more than 0.05 was considered to be insignificant.

Results

Sociodemographic Characteristics

A total of 145 questionnaires were distributed and 145 were received with a response rate of 100%. The gender distribution was 91 (62.8%) males and 54 (37.2%) females. Mean age of the study participants was 21.2 (0.44) years old. Most participants were in the 21 to 30 years old age group, 122 (84.1%). In terms of marital status, there were 76 (52.4%) singles, 52 (35.9%) married individuals, and 17 (11.7%) divorcees. Regarding educational level, 11 (7.6%) individuals had up to secondary level education, 102 (70.3%) individuals had undergraduate degree, and 32 (22.1%) individuals had postgraduate degree. Majority of the participants were students, hence, 128 (88.3%) of them identified themselves as unemployed. The sociodemographic characteristics of the Somalian immigrants in Malaysia is shown in Table 2.

Table 2. Sociodemographic characteristics among Somalian immigrants in the state of Selangor, Malaysia.

Variable	Group	Frequency (n)	Percent (%)
Age (21.2 ± 0.44)	< 20	4	2.8
	21-30	122	84.1
	31-40	17	11.7
	> 41	2	1.4
Gender	Male	91	62.8
	Female	54	37.2
Marital status	Single	76	52.4
	Married	52	35.9
	Divorced	17	11.7
Education level	Up to Secondary	11	7.6
	Undergraduate degree	102	70.3
	Postgraduate degree	32	22.1
Working status	Employed	17	11.7
	Unemployed	128	88.3

Assessment of Knowledge towards Hepatitis B

The participants knowledge on HBV infection, etiology, transmission and vaccination was evaluated with 24 questions related to public knowledge on HBV infection. Scoring of more than 12 was considered to be an indicator of good knowledge on HBV infection. Among the

participants, 120 (82.8%) of them showed good knowledge regarding HBV in general as well as its etiology, transmission and vaccination. The mean knowledge score among the participants was 16.9 (4.88). Table 3 illustrates the participants knowledge on different aspects of HBV infection.

Table 3. Level of knowledge on HBV infection among Somalian immigrants in the state of Selangor, Malaysia.

No	Items	Yes	No
		N (%)	N (%)
1.	Have you ever heard of disease termed as Hepatitis B?	86 (59.3)	59 (40.7)
2.	Do you know what is the Hepatitis B infection?	81 (55.9)	64 (44.1)
3.	Do you know the causes of Hepatitis B?	100 (69.0)	45 (31.0)
4.	Can Hepatitis B virus infections result in chronic hepatitis and liver cancer?	125 (86.2)	20 (13.8)
5.	Can Hepatitis B affect liver function?	127 (87.6)	18 (12.4)
6.	Can Hepatitis B affect any age group?	96 (66.2)	49 (33.8)
7.	Can Hepatitis B be transmitted by un-sterilized syringes, and surgical instruments?	116 (80.0)	29 (20.0)
8.	Can Hepatitis B be transmitted by using blades of the barber/ear and nose piercing?	115 (79.3)	30 (20.7)
9.	Can Hepatitis B be transmitted from mother to child?	110 (75.9)	35 (24.1)
10.	Can Hepatitis B be transmitted by contaminated water/food prepared by person suffering with these infections?	47 (32.4)	98 (67.6)
11.	Is Hepatitis B virus transmission hereditary?	61 (42.1)	84 (57.9)
12.	The early symptoms of Hepatitis B are same like cold and flu (runny nose and cough)	57 (39.3)	88 (60.7)
13.	Do you know the treatment for Hepatitis B virus infection?	67 (46.2)	78 (53.8)
14.	Can Hepatitis B be self-cured by body?	77 (53.1)	68 (46.9)
15.	Is vaccination available for Hepatitis B?	122 (84.1)	23 (15.9)
16.	Do you recommend vaccination against Hepatitis B among your family members?	126 (86.9)	19 (13.1)
17.	Are you vaccinated against Hepatitis B?	121 (83.4)	24 (16.6)
18.	Are you aware of the appropriate intervals of the Hepatitis B vaccination?	128 (88.3)	17 (11.7)
19.	Do you know the precautionary measures to be taken against Hepatitis B in your routine practice?	115 (79.3)	30 (20.7)
20.	Are you aware of the risk of exposure to Hepatitis B at the dentist?	114 (78.6)	31 (21.4)
21.	Do you know the symptoms of Hepatitis B infection?	123 (84.8)	22 (15.2)
22.	Does HBV spread by casual contact such as hand shaking?	76 (52.4)	69 (47.6)
23.	Do you know the mode of disease transmission?	129 (89.0)	16 (11.0)
	- Sexual intercourse	37 (25.5)	-
	- Blood transfusion	76 (52.4)	-
	- Infection needle	32 (22.1)	-
24.	What are the methods of prevention of hepatitis B virus?		
	- Avoid unprotected sex	61 (42.1)	-
	- Vaccination	39 (26.9)	-
	- Material sterilization	20 (13.8)	-
	- Avoid diagnosed HBV patients	15 (10.3)	-
	- Proper disposal of sharps	2 (1.4)	-
	- Avoid needle stick injury	5 (3.4)	-
	- Avoid food that is not well cooked	3 (2.1)	-

Assessment of Attitude towards Hepatitis B

The participants attitude towards HBV transmission, prevention, vaccination, and consequences was evaluated with 15 questions related to public perception towards HBV infection. Scoring of more than 7 was considered to be an indicator of good attitude towards HBV infection.

Among the participants, 114 (78.6%) of them showed good attitude towards HBV transmission, prevention, vaccination, and consequences. The mean attitude score among the participants was 9.2 (2.94). Table 4 illustrates the participants attitude towards different aspects of HBV infection.

Table 4. Level of attitude towards HBV infection among Somalian immigrants in the state of Selangor, Malaysia.

No	Items	Yes	No
		N (%)	N (%)
1.	Do you think you can get Hepatitis B?	102 (70.3)	43 (29.7)
2.	Would you share a meal with an infected person?	103 (71.0)	42 (29.0)
3.	Should an infected person be terminated from school/work?	101 (69.7)	44 (30.3)
4.	Is vaccination against Hepatitis B necessary?	107 (73.8)	38 (26.2)
5.	Have you ever thought you could get Hepatitis B?	82 (56.6)	63 (43.4)
6.	Should Hepatitis B patient be allowed to work routinely?	75 (51.7)	70 (48.3)
7.	Should Hepatitis B patient be isolated?	116 (80.0)	29 (20.0)
8.	Is education effective in improving the knowledge on Hepatitis B?	108 (74.5)	37 (25.5)
9.	How do you deal with Hepatitis B patient? - I avoided dealing with them - I deal with them cautiously - I deal with them normally	- 83 (57.2) 32 (22.1) 30 (20.7)	- - - -
10.	What will be your reaction when you found that you have Hepatitis B? - Fear - Shame - Surprise - Sadness	- 39 (26.9) 15 (10.3) 13 (9.0) 78 (53.8)	- - - - -
11.	Who will you talk to about your illness? - Physician - Spouse - Parents - Child - Others relatives - Friends - No one	- 54 (37.2) 5 (3.4) 32 (22.1) 20 (13.8) 4 (2.8) 4 (2.8) 26 (17.9)	- - - - - - -
12.	What will you do if you think that you have symptoms of Hepatitis B? - Self-treatment - Going to a health facility - Going to a homeopath - Going to a traditional healer	- 67 (46.2) 51 (35.2) 3 (2.1) 24 (16.6)	- - - -
13.	When you have symptoms of Hepatitis B, at what stage will you come to the health facility? - After self-treatment fails - Three to four weeks after appearance of symptoms - As soon as I realized the symptoms of HBV - Will not see a physician at all	- 24 (16.6) 34 (23.4) 45 (31.0) 42 (29.0)	- - - -
14.	How expensive do you think the diagnosis and treatment of Hepatitis B is? - Free - Reasonable - Somewhat expensive - Expensive - Do not know	- 34 (23.4) 10 (6.9) 17 (11.7) 19 (13.1) 65 (44.8)	- - - - -
15.	What will worry you the most after you are diagnosed with Hepatitis B? - Fear of death - Fear that the disease spread to a family member - The treatment cost - Isolation from society	- 66 (45.5) 35 (24.1) 13 (9.0) 31 (21.4)	- - - -

Assessment of Practice towards Hepatitis B

The participants practice towards HBV screening, prevention and vaccination was evaluated with 16 questions related to the public action in terms of controlling HBV infection. Scoring of more than 8 was considered to be an indicator of good practice

towards HBV infection. Among the participants, only 47 (32.4%) of them showed good practice towards HBV screening, prevention, and vaccination, tabulated in table 5. The mean practice score among the participants was 9.4 (2.93).

Table 5. Level of practice towards HBV infection among Somalian immigrants in the state of Selangor, Malaysia.

No	Items	Yes	No
		N (%)	N (%)
1.	Have you done screening for Hepatitis B?	89 (61.4)	56 (38.6)
2.	Have you got yourself vaccinated against Hepatitis B	74 (51.0)	71 (49.0)
3.	Do you ask for screening of blood before transfusion?	101 (69.7)	44 (30.3)
4.	Do you ask for new syringe before use?	82 (56.6)	63 (43.4)
5.	Do you ask your barber to change blade/or for safe equipment's for ear and nose piercing?	104 (71.7)	41 (28.3)
6.	Do you share food/utensils/water etc. with others?	70 (48.3)	75 (51.7)
7.	Do you avoid meeting people?	59 (40.7)	86 (59.3)
8.	Will you continue and complete the treatment for the infection?	102 (70.3)	43 (29.7)
9.	Have you ever participated in health education program related to Hepatitis B?	96 (66.2)	49 (33.8)
10.	Have you had any injury or wound while working?	89 (61.4)	56 (38.6)
11.	Have you been subjected to Hepatitis B examination?	97 (66.9)	48 (33.1)
12.	Have you ever accidentally splashed blood or body fluids in your mouth, eyes, and nose?	38 (26.2)	107 (73.8)
13.	Have you been subjected to premedical examination on appointment?	94 (64.8)	51 (35.2)
14.	Is there a periodical medical examination? If yes, how frequent?	99 (68.3)	46 (31.7)
	- Every one to two months	49 (33.8)	-
	- Every three to six months	34 (23.4)	-
	- Once a year	36 (24.8)	-
	- Do not response	26 (17.9)	-
15.	Have you received full dose of vaccination against Hepatitis B?	55 (37.9)	90 (62.1)
16.	How many doses of Hepatitis B vaccine did you receive?	-	-
	- One dose	24 (16.6)	-
	- Two doses	33 (22.8)	-
	- Three doses	55 (37.9)	-
	- Do not know	33 (22.8)	-

Association of Sociodemographic Characteristics and KAP Scores

The relationship between the sociodemographic characteristics among the participants and the outcome of their knowledge, attitude and practice

towards HBV infection scores were determined using descriptive statistics and Chi-square test. Sociodemographic characteristics were found to have no effect on the KAP scores as depicted in Table 6.

Table 6. Sociodemographic characteristics among Somalian immigrants in the state of Selangor ,Malaysia.


Description	N (%)	Good Knowledge	P-value	Good Attitude	P-value	Good Practice	P-value
Age							
< 20	4 (2.8)	3 (75.0)	0.899	3 (75.0)	0.639	0 (0.0)	0.388
21-30	122 (84.1)	101 (82.8)		94 (77.0)		41 (33.6)	
31-40	17 (11.7)	14 (82.4)		15 (88.2)		6 (35.3)	
> 41	2 (1.4)	2 (100.0)		2 (100.0)		0 (0.0)	
Gender							
Male	91 (62.8)	76 (83.5)	0.754	75 (82.4)	0.148	30 (33.0)	0.853
Female	54 (37.2)	44 (81.5)		39 (72.2)		17 (31.5)	
Marital status							
Single	76 (52.4)	58 (76.3)	0.085	56 (73.7)	0.159	30 (39.5)	0.145
Married	52 (35.9)	46 (88.5)		42 (80.8)		12 (23.1)	
Divorced	17 (11.7)	16 (94.1)		16 (94.1)		5 (29.4)	
Education level							
Up to Secondary	11 (7.6)	9 (81.8)	0.723	7 (63.6)	0.204	4 (36.4)	0.174
Undergraduate Degree	102 (70.3)	83 (81.4)		84 (82.4)		37 (36.3)	
Postgraduate Degree	32 (22.1)	28 (87.5)		23 (71.9)		6 (18.8)	
Working status							
Employed	17 (11.7)	16 (94.1)	0.187	13 (76.5)	0.818	6 (35.3)	0.787
Unemployed	128 (88.3)	104 (81.3)		101 (78.9)		41 (32.0)	

HBV Infection Status

All the participants included in this study were known to be free of HBV infection. Molecular analysis for the presence of occult HBV infection was done via PCR detection of HBV DNA. Successful amplification of the HBV DNA will produce a 350 base pair product that can be seen in the agarose gel electrophoresis. All samples were

tested negative for HBV DNA. None of the samples demonstrated successful amplification of the 350 base pair HBV DNA PCR product as shown in Figure 1. Hence, the association between knowledge, attitudes and practices towards HBV infection among Somalian immigrants in the state of Selangor, Malaysia, and their HBV infection status could not be evaluated.

Figure 1. PCR Product of HBV DNA Nested PCR.



Discussion

The existence of vaccine against HBV infection, since 1982, has led to the really low prevalence of HBV infection in many developed countries (Siraj, Fareed, and Mahajan 2016). However, prevalence of HBV infection is still high in sub-Saharan Africa and South East Asia due to the lack of awareness and action taken towards controlling the disease (Terrault et al. 2018).

Malaysia, a country located in South East Asia, is one of the few countries in Asia who has low prevalence of HBV infection (Elbur et al. 2017). Considering the potential of HBV transmission via immigration, particularly within migrants from countries with a high HBV endemicity to those with a low HBV endemicity, the Malaysian government requires a mandatory health screening for all traveler within the first five days of arrival in Malaysia. This mandatory screening also applies to the Somalian immigrants, whose home country have reported prevalence of 18.9% to 29% of the population (Mansour-Ghanaei et al. 2013).

The current study sought to evaluate knowledge, attitude and practice towards HBV infection among Somalian immigrants in Malaysia and their relationships to the HBV infection status of the Somalian immigrants. Results of the study showed overall good level of knowledge and attitude of the participants toward HBV infection. However, a poor level of practice towards HBV infection was detected among the participants. Nevertheless, all participants were devoid of molecular evidence of HBV infection. Moreover, participant's sociodemographic characteristics also shown no association their KAP.

Numerous studies had been done in the past to assess the level of KAP towards HBV infection in variety of population, mainly healthcare professionals (Haq et al. 2013). In terms of knowledge and attitude, many of the population studied demonstrates fair level of understanding and perception towards the disease.

However, vaccination compliance has been continuously identified as a gap in practice in many of these studies. In the current study, Somalian immigrants demonstrated similar pattern, where they have good knowledge and attitude but poor practice.

A high level of knowledge on the biomedical concepts revolving around HBV was expected in healthcare professional's population. This typically translates to good attitude towards the disease. Study focusing on dentist, physician, healthcare workers, medical students, and dental students, (Haq et al. 2013) all revealed a fair level of knowledge and attitude towards HBV. In non-clinical university students, a study conducted among students of Fayoum University, Egypt, revealed that three-quarters of the participants had a satisfactory level of knowledge and attitude towards HBV (Zampino et al. 2015), (Vahdat et al. 2014), (Wang 2009), WHO, 2018, WHO, 2019.

In the current study, 82.8% out of the 145 participants, demonstrated good level of knowledge towards HBV. Good knowledge was apparent among the participants because majority (88.3%) of them were students who had prior knowledge about the infection. The mean knowledge score for the complete cross-sectional study was 16.9 (4.88). A good level of knowledge is indicated by the score of 12. As expected, level of attitude follows level of knowledge with 78.6% of participants showed good attitude towards HBV. The mean attitude score for the complete cross-sectional study was 9.2 (2.94). A good level of attitude is indicated by the score of 7.

Sufficient level of knowledge and good attitude towards a certain disease can lead to a good practice in the disease prevention and control (Kim H et al. 2019). However, many KAP studies in HBV has shown a relatively poor level of practice despite a good level of knowledge and attitude towards it. When inspected deeper, vaccination compliance was identified as the recurrent gap in the practice of

dentist (Elbur et al. 2017), physician, healthcare workers, medical students, and dental students. Unavailability of the vaccine in developing country such as Ethiopia has contributed to the low vaccination practice among study populations.

In the current study, only 32.4% of the participants demonstrated good level of practice towards HBV. More than half of the population (62.1%) did not received the recommended three doses of the HBV vaccine. However, the mean knowledge score for the complete cross-sectional study indicated a good practice with a score of 9.4 (2.93), slightly higher than the cut off score of 8.

In terms of HBV infection status, all study participants were found to be devoid of traces of HBV DNA in their blood. Nested PCR is a method that utilized two sets of primers to amplify low occurring DNA sequence. This method has been proven to be able to detect occult HBV infection in 4% of the 1000 HBV negative Iranian (Raihan R et al. 2019). The negative result in Somalian immigrants in Malaysia suggest a positive correlation between good KAP and the HBV infection status among Somalian immigrants in Malaysia.

Although the finding may not be representative of the entire Somalian immigrants in Malaysia, the current study demonstrates a gap regarding practice on prevention of HBV among Somalian immigrants in the state of Selangor. This study can serve as a reference to urge the Somalian Ministry of Health to prioritize the awareness and education on the HBV prevention measures, particularly its vaccination program. This highlights the need to make the appropriate HBV vaccine freely available in the health centers along with educating the public, especially pregnant mothers, on the importance of vaccination against HBV.

Conclusion

Knowledge and attitude towards HBV infection among Somalian immigrants in Malaysia were acceptable but their practice towards HBV infection were poor. Nevertheless, all of the participants were devoid of molecular evidence of HBV infection, suggesting positive correlation between good KAP and the HBV infection status among Somalian immigrants in Malaysia.

Acknowledgements

This research was supported by an internal funds for postgraduate project of University Selangor.

Conflict of interest

The authors have no conflicts of interest to disclose.

References

Abd El-Nasser G, Abed El baset E. Assessment of Students Knowledge and Attitude Toward Hepatitis B and C in Sohag University/Egypt. *KUFA J Nurs Sci*. 2013;3(1).

Abdela A, Woldu B, Haile K, Mathewos B, Deressa T. Assessment of knowledge, attitudes and practices toward prevention of hepatitis B virus infection among students of medicine and health sciences in Northwest Ethiopia. *BMC Res Notes* [Internet]. 2016 Dec 19 [cited 2019 Jan 30];9(1):410. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27543117>

Abiola A-HO, Agunbiade AB, Badmos KB, Lesi AO, Lawal AO, Alli QO. Prevalence of HBsAg, knowledge, and vaccination practice against viral hepatitis B infection among doctors and nurses in a secondary health care facility in Lagos state, South-western Nigeria. *Pan Afr Med J* [Internet]. 2016 [cited 2019 Jan 30];23. Available from: <http://www.panafrican-med-journal.com/content/article/23/160/full/>

- Adjei, Charles Ampong et al. 2016.** “Mother-to-Child Transmission of Hepatitis B: Extent of Knowledge of Physicians and Midwives in Eastern Region of Ghana.” *BMC Public Health*: 1–7. <http://dx.doi.org/10.1186/s12889-016-3215-6>.
- Baig VN, Gupta PK, Sharma AK, Madhusudan S.** Assessment of knowledge, attitude and practice about hepatitis B among Clinicians & Medical students: a cross sectional study. *Natl J Community Med*. 2015;6(2):415–22.
- Coppola N, Alessio L, Pisaturo M, Macera M, Sagnelli C, Zampino R, et al.** Hepatitis B virus infection in immigrant populations. *World J Hepatol* [Internet]. 2015 Dec 28 [cited 2019 Jan 30];7(30):2955–61. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26730274>
- Chen CJ, Wang LY, Yu MW.** Epidemiology of hepatitis B virus infection in the Asia-Pacific region. *J Gastroenterol Hepatol* [Internet]. 2000 May [cited 2019 Jan 30];15 Suppl:E3-6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10921373>
- Hamissi J, Tabari ZA, Najafi K, Hamissi H, Hamissi Z.** Knowledge, Attitudes and Practice of Hepatitis B vaccination among Iranian dentists. *Int J Collab Res Intern Med Public Heal*. 2009;6(7).
- Hassan-Kadle MA, Osman MS, Ogurtsov PP.** Epidemiology of viral hepatitis in Somalia: Systematic review and meta-analysis study. *World J Gastroenterol* [Internet]. 2018 Sep 14 [cited 2019 Jan 30];24(34):3927–57. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30228786>
- Kim H, Yang JD, El-Serag HB, Kanwal F.** Awareness of Chronic Viral Hepatitis in the United States: An Update from National Health and Nutrition Examination Survey. *J Viral Hepat*. 2019 Jan 10;
- Li X, Kang H, Wang S, Deng Z, Yang T, Jia Y, et al.** Knowledge, Attitude, and Behavior of Hepatitis B Virus Infection among Chinese Dental Interns. *Hepat Mon* [Internet]. 2015 May 23 [cited 2019 Jan 30];15(5):e25079. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26045703>
- MacLachlan JH, Cowie BC.** Hepatitis B virus epidemiology. *Cold Spring Harb Perspect Med* [Internet]. 2015 May 1 [cited 2019 Jan 30];5(5):a021410. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25934461>
- Mesfin YM, Kibret KT.** Assessment of Knowledge and Practice towards Hepatitis B among Medical and Health Science Students in Haramaya University, Ethiopia. Schildgen O, editor. *PLoS One* [Internet]. 2013 Nov 21 [cited 2019 Jan 30];8(11):e79642. Available from: <https://dx.plos.org/10.1371/journal.pone.0079642>
- Nelson NP, Easterbrook PJ, McMahon BJ.** Epidemiology of Hepatitis B Virus Infection and Impact of Vaccination on Disease. *Clin Liver Dis*. 2016;20(4):607–28.
- Raihan R, Mohamed R, Radzi Abu Hassan M, Md Said R.** Chronic Viral Hepatitis in Malaysia: “Where are we now?” *Euroasian J hepato-gastroenterology* [Internet]. 2017 [cited 2019 Jan 30];7(1):65–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29201775>
- Sagnelli, Evangelista et al. 2014.** “Epidemiology of Acute and Chronic Hepatitis B and Delta over the Last 5 Decades in Italy.” 20(24): 7635–43.
- Siraj F, Fareed P, Mahajan N.** Assessment of knowledge attitude and practice towards hepatitis B among health care workers in a tertiary care hospital. *Int J Reprod Contraception, Obstet Gynecol* [Internet]. 2016 Dec 16 [cited 2019 Jan 30];5(1):58–61. Available from: <http://www.ijrcog.org/index.php/ijrcog/article/view/501>
- Terrault NA, Lok ASF, McMahon BJ, Chang K-M, Hwang JP, Jonas MM, et al.** Update on

prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology* [Internet]. 2018 Apr 1 [cited 2019 Jan 30];67(4):1560–99. Available from: <http://doi.wiley.com/10.1002/hep.29800>

Elbur, Abubaker Ibrahim, Nawaf Almalki, Abdualziz Alghamdi, and Housain A Alqarni Alqarni. 2017. “Knowledge , Attitude and Practice on Hepatitis B : A Survey among the Internet Users in Taif , Kingdom of Saudi Arabia.” 3(3): 1–7.

Ul Haq N, Hassali MA, Shafie AA, Saleem F, Farooqui M, Aljadhey H. A cross sectional assessment of knowledge, attitude and practice towards Hepatitis B among healthy population of Quetta, Pakistan. *BMC Public Health*. 2012 Aug 23;12(1):692.

Vahdat S, Hamzehgardeshi L, Hessam S, Hamzehgardeshi Z. Patient involvement in health care decision making: a review. *Iran Red Crescent Med J*. 2014 Jan;16(1):e12454.

Vaezjalali M, Rashidpour S, Rezaee H, Hajibeigi B, Zeidi M, Gachkar L, et al. Hepatitis B viral DNA among HBs antigen negative healthy blood donors. *Hepat Mon*. 2013 Mar;13(3):e6590.

Wang T. Model of life expectancy of chronic hepatitis B carriers in an endemic region. *J Epidemiol*. 2009;19(6):311–8.

World Health Organization. Hepatitis B [Internet]. 2018. Available from: <https://www.who.int/en/news-room/fact-sheets/detail/hepatitis-b>

World Health Organization. Hepatitis B vaccination has dramatically reduced infection rates among children in Europe, but more is needed to achieve elimination [Internet]. World Health Organization; 2017 [cited 2019 Jan 30]. Available from: <http://www.euro.who.int/en/health-topics/communicable-diseases/hepatitis/news/news/2017/04/hepatitis-b->

[vaccination-has-dramatically-reduced-infection-rates-among-children-in-europe,-but-more-is-needed-to-achieve-elimination](http://www.euro.who.int/en/health-topics/communicable-diseases/hepatitis/news/news/2017/04/hepatitis-b-vaccination-has-dramatically-reduced-infection-rates-among-children-in-europe,-but-more-is-needed-to-achieve-elimination)

Zampino R, Boemio A, Sagnelli C, Alessio L, Adinolfi LE, Sagnelli E, et al. Hepatitis B virus burden in developing countries. *World J Gastroenterol* [Internet]. 2015 Nov 14 [cited 2019 Jan 30];21(42):11941–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26576083> .

