



Journal of Bioscience and Applied Research

JBAAR

WWW.JBAAR.ORG



## The Antioxidant defense Responses of Sea cucumber *Holothuria polii* Against Rickettsia-like organism (RLOs) Infection and Heavy Metal Pollution in Alexandria coast

Mohamed M. Elghazaly and Aml Z. Ghoneim

Department of Zoology, Faculty of Science, Damanhour University, El Behiera,  
Egypt

CORRESPONDING AUTHOR: E-mail: [m\\_ghzaly@yahoo.com](mailto:m_ghzaly@yahoo.com)

### Abstract

*Holothuria polii*, the common black sea cucumber (Echinodermata, Holothuroidea), is abundant in Mediterranean Sea along Alexandria coast. This species was given the great valued commercially and for human consumption. The present study was objective to assess the biological effect of marine pollution on the sea cucumber, *Holothuria polii* collected from two locations in Alexandria coast; Abo-Qir station as an industrial area compared with Miami station as a reference area, using the oxidative stress biomarkers approach, heavy metals bio-accumulation, and Rickettsia-like organism (RLOs) infection. Sea cucumber samples were collected from both stations in Alexandria, during two seasons; winter (January) and summer (July) 2016. Atomic absorption spectrophotometer was used to detect Zn, Mn, pb, Cu and Cd in the body wall of the animal as well as in water samples.

The results showed a significant decrease in the antioxidant parameters; catalase (CAT), superoxide dismutase (SOD), Glutathione reductase (GSH), and Glutathione transferase(GST) activities, while the Malondialdehyde (MDA) level was significantly increased in the body wall tissue of sea cucumber collected from Abo-Qir station as compared with Miami station. The concentrations of heavy metals (Zn, Mn, Pb, Cu and Cd) in body wall of sea cucumber and sea water samples at the two stations differed significantly, showed increasing in the industrial area (Abo-Qir station) compared with Miami station. As parasites are ubiquitous in the marine pollutant environment, so the present study proved that the sea cucumber samples collected from Abo-Qir station were infected by Rickettsia-like organism (RLOs), found in the digestive tract and the respiratory tree of this marine animal. The present results of pollutant biomarkers indicated that, the degree of pollution increased in summer as compared with winter season in the two locations.

This study also suggested that the sea cucumber was an ideal bioindicator animal for marine pollution. The suppression of antioxidant biomarkers in the tissue of sea cucumber compared with high values of MDA confirmed the presence of anthropogenic contaminants in Abo-Qir station, which essentially due to the industrial rejections. Also, Rickettsia-like organism (RLOs) infection and the bio-accumulation of the heavy metals in the tissues of sea cucumber obstructed the antioxidant defense responses in the animal.

**Keywords:** Oxidative stress, Heavy metals, Rickettsia-like organism (RLOs), Marine Pollution, Alexandria.

## **Introduction**

Sea cucumber *Holothuria polii*, is a soft-bodied and worm-like marine invertebrate lives on the sea floor worldwide, but most abundant in tropical shallow-water coral reefs. It belongs to the Phylum Echinodermata; class Holothuroidea, order Aspidochirotida, and family Holothuriidae (Delle Chiaje, 1823). The commercial Sea cucumbers have been well recognized as a tonic and traditional remedy in Japan, China and Malaysia for their benefits against hypertension, asthma, rheumatism, constipation, burns and impotence (Hu *et al.*, 2010). These medicinal benefits and health functions of sea cucumbers can be attributed to the presence of appreciable amounts of bioactive compounds, such as the triterpene glycosides (saponins), chondroitin sulfates, glycosaminoglycan, sulfated polysaccharides, sterols (glycosides and sulfates), phenolics, peptides, cerberosides and lectins (Bordbar *et al.*, 2011).

In marine environment, sea cucumbers may accumulate many contaminants such as heavy metals under chronic exposure to rather high levels of chemical stress (Luo *et al.* 2010). Heavy metals are considered as critical contaminants since they are toxic, persistent, and non-biodegradable and would inevitably be accumulated and biomagnified in food chain, exerting deleterious effects on the animals and human health when their concentrations are highly elevated (Spanopoulos-Zarco *et al.*

2014). In marine animals, it has been found that heavy metal accumulation is positively correlated with metal concentration in diet and ambient water (Julshamn, Grahl- Nielsen, 1996). Water pollution is a serious problem in the global context, heavy metal pollution in marine environment caused by diverse activities such as phosphate fertilizers, industrial effluent, foundry wastes, paints, mining and rock weathering (Muduli & panda, 2010). The bioaccumulation of contaminants by the tissues and organs of marine organisms has been extensively studied throughout the world and led to the adoption of the bioindicator concept for the environmental quality assessment (Langston & Spence, 1995). The origin of trace elements in the sampled water of the Egyptian Mediterranean sea is mainly the waste water discharge (Radwan *et al.*, 2014).

The importance of oxidative stress response as potential biomarkers of environmental pollution has been addressed by different experimental approaches (Ferreira *et al.*, 2005; Orbea *et al.*, 2002). The biomarkers utilized include components of oxidative adaptive responses, such as antioxidant enzyme activities (catalase (CAT), superoxide dismutases (SODs) or glutathione peroxidase (GPX), or the estimation of oxidative damages in lipids, proteins and DNA (Filho, 1996). When the defense mechanisms are unbalanced regarding the increased presence of ROS

generated compounds, e.g. by the presence of pollutants, oxidative damage will occur, indicating a mechanism of toxicity in aquatic organisms. Ferreira *et al.*, (2007) mentioned that, antioxidant systems can be considered as non-specific biomarkers of exposure to pollutants, and also as an indicator of toxicity.

Rickettsia-like organisms (RLOs) are small, pleomorphic, rod-shaped coccid prokaryotes, most of which are obligate intracellular Gram-negative parasites (Edgerton *et al.*, 2002) some RLOs have been associated with diseases in mollusks and other marine animals (Lauckner, 1983). Despite their importance as causative agents of severe mortality outbreaks in farmed aquatic species, little is known about the life cycle of Rickettsia-like organisms and their host range (Ferrantini *et al.* 2009). RLOs may have severe physiological implications in cultured organisms (Branson & Diaz-Munoz 2006), followed by massive mortalities.

The aim of this work was to describe the relation between the infection with RLOs found

in sea cucumber *Holothuria polii*, pollution with heavy metals, and the antioxidant defense responses in the animal.

## Material and methods:

### 1-Sampling sites:

Samples of sea cucumber were collected from 5-9m depth at Abu-Qir and Miami stations at Alex sea water (Fig1). Samples were collected seasonally (winter and summer).

### 2-Sample collection:

Collected samples were kept alive in containers filled with water. Samples of sea cucumber *Holothuria polii* were taken to the laboratory and transferred to well aerate aquaria. 40 specimens were collected from each station. The water temperature was 27<sup>o</sup> for water sampling. Samples were collected by hand-picking through forceps.



Fig.(1).Map showing the sites of collection. Google Maps.

### 3-Metal analysis:

The body wall of sea cucumber was separated to remove all the internal organs before analysis and only the body wall was used for analysis after cleaning. The body walls were weighted (g). Ash samples of each specimen were weighed (g), dissolved in HCl (0.1 mol/L) and further treated with H<sub>2</sub>O<sub>2</sub> (30%) till lucid solutions were formed, and then diluted by water. The 0.45 µm Whatman filter papers were chosen for the filtration purposes. Cu, Cd, Zn, Mn and Pb levels were determined by Perkin Elmer Analyst 700 Atomic Absorption Spectrophotometer, and the detection limits. The levels of different metals in the sea water were also analyzed three times by Atomic absorption spectrophotometer as described. Heavy metals in tissue were expressed as (mg/kg tissue) and in the sea water as (mg/L water).

### 4- Determination of the antioxidant parameters:

The tissue of the animal was separated and homogenized in TRIS buffer and prepared in ice cold saline (0.9%), and the homogenized tissues were centrifuged at 3000 rpm at 4°C

for 30 min. The obtained supernatants were used for determination of malondialdehyde (MDA) as a measure of lipid peroxidation (Yoshioka *et al.*, 1979), also some antioxidant parameters were measured such as; superoxide dismutase (SOD) activity was measured according to the method of (Beauchamp and Fridovich, 1971), Tissue catalase (CAT) activity was assayed using the method described previously (Claiborn 1986), Glutathione Transferase (GST) activity was measured according to method described by (Habig *et al.*, 1974), and reduced glutathione (GSH) was measured according to the method of (Beutler *et al.*, 1963). The results were expressed as (Unit/100 g tissue).

### 5-Estimation of Total Protein:

Total protein content in the tissue of the animal was estimated according to method described by (Lowry *et al.*, 1951), data expressed as (mg /L).

### 6- Physicochemical analysis:

**Table (1):** The physicochemical parameters and the technique /tools that used to analyze the Seawater.

Parameter	Technique / Tool
Water temperature (c)	Dry thermometer
Salinity (S%)	Conductivity Salinometer (Backman; model RS.10)
pH	Digital pH meter model 209
Dissolved oxygen (mg/L)	Modified Winkler method (Grasshoff, 1976)

## 7- Histological examination:

Part of infected organs (respiratory tree and digestive gland) was fixed in 10% formalin solution. Fixed samples were dehydrated and embedded in paraffin wax (Bell and Lightner, 1988), then sectioned at 5-6 $\mu$ m in thickness using a microtome, mounted on glass slides and stained using haematoxylin and eosin (H & E) and toluidine blue.

## 8- Transmission electron microscopy:

Respiratory tree and digestive gland of sea cucumber were fixed in 2.5% glutaraldehyde solution (pH 7.2, buffered 0.1 M phosphate buffer) for 2-4 hrs at 4°C and rinsed in 0.1 M phosphate buffer and then post-fixed in 1% osmium tetroxide (OsO<sub>4</sub>) solution for 2 hrs at 4°C. After fixation, the specimens were washed with 0.1 M phosphate buffer 4 times for 2 hrs and dehydrated with ascending grades of ethanol, then were embedded in Epon 812, cut at ultrathin sections (70 nm in thickness) and placed on copper grids (200 meshes) in order to double stain with uranyl acetate and lead citrate. Specimens were examined using a TEM (Jem1200exII, Jeol, Japan).

## 9-Statistical analysis

Student t-test for comparing between the two seasons in each location was applied; the values are mean  $\pm$  S.D. for 6 animals in each group. P-Values < 0.05 were considered as significant.

## **RESULTS:**

### **Metal analysis:**

Measurements of heavy metals in water samples in the investigated areas; Abo-Qir and Miami showed significant increase in values of zinc (Zn), Manganese (Mn), lead (Pb), copper (Cu) and cadmium (Cd) in sea water collected from Abo-Qir during summer. The accumulation of these metals in tissue samples of sea cucumber collected from the same sites showed that the concentrations of metals in the order: Cu>Cd>Zn>Mn>Pb. The concentrations of Zn, Mn, Pb, Cd in tissue samples were significantly higher in Abo-Qir station during summer season, while the concentration of Cu showed non-significant increase (table 2).

Table (2): Comparison between the two locations in each season according to heavy metals.

	Winter			Summer		
	Abo-Qir (n = 6)	Miami (n = 6)	P	Abo-Qir (n = 6)	Miami (n = 6)	P
<b>Heavy metals in tissue (mg/kg tissue)</b>						
Zn	4.6±0.2	4.3 <sup>@</sup> ±0.2	0.008*	5.0±0.2	4.6 <sup>@</sup> ±0.2	0.003*
Mn	6.2±0.4	4.2 <sup>@</sup> ±0.3	<0.001*	6.4±0.3	5 <sup>@</sup> ±0.4	<0.001*
Pb	1.9±0.1	0.6 <sup>@</sup> ±0	<0.001*	1.9±0.1	1 <sup>@</sup> ±0.2	<0.001*
Cu	2.7±0.4	2.4±0.2	0.063	2.9±0.2	2.6±0.2	0.059
Cd	1.2±0.2	0.9 <sup>@</sup> ±0.1	0.018*	1.3±0.2	1 <sup>@</sup> ±0.1	0.005*
<b>Heavy metals in water (mg/L water)</b>						
Zn	5.2±0.2	4.8 <sup>@</sup> ±0.1	0.009*	5.4±0.3	5 <sup>@</sup> ±0.2	0.017*
Mn	4.4±0.2	2.6 <sup>@</sup> ±0.2	<0.001*	4.2±0.3	3 <sup>@</sup> ±0.2	<0.001*
Pb	2.2±0.2	1.1 <sup>@</sup> ±0.2	<0.001*	2.4±0.2	1.3 <sup>@</sup> ±0.2	<0.001*
Cu	1.9±0.1	1.2 <sup>@</sup> ±0.2	<0.001*	1.9±0.1	1.3 <sup>@</sup> ±0.2	<0.001*
Cd	1.2±0.2	0.9 <sup>@</sup> ±0.1	0.003*	1.3±0.2	1 <sup>@</sup> ±0.1	0.008*

p<sub>1</sub>: p value for **Student t-test** for comparing between the two locations in each seasons

@: Statistically significant between the two locations in each seasons

\*: Statistically significant at  $p \leq 0.05$

### Antioxidant parameters investigation:

The current results showed a significant decrease in the antioxidant parameters; catalase (CAT), superoxide dismutase (SOD), Glutathione reductase (GSH), and Glutathione transferase (GST) activities, while the Malondialdehyde (MDA) level was

significantly increased in the body wall tissue of sea cucumber collected from Abo-Qir station as compared with Miami station. The protein content in tissue samples recorded insignificant decrease in Abo-Qir compared with Miami location (table 3 and Figs 2).


**Table (3): Comparison between the two seasons in each location according to different parameters**

	Abo-Qir			Miami		
	Winter (n = 6)	Summer (n = 6)	P	Winter (n = 6)	Summer (n = 6)	P
MDA(Unit/100 g tissue)	47.5±2.4	53.3 <sup>#</sup> ±3.5	0.007*	35.7±2.8	41.7 <sup>#</sup> ±2.7	0.004*
CAT(Unit/100 g tissue)	80±3.9	70.8 <sup>#</sup> ±5.5	0.008*	136.3±7.8	122.7 <sup>#</sup> ±7.9	0.013*
SOD (Unit/100 g tissue)	19.7±2.1	17.5±1.9	0.086	34.5±4.5	28.5 <sup>#</sup> ±2.9	0.021*
GSH (Unit/100 g tissue)	25.5±3.1	21.5 <sup>#</sup> ±1.9	0.022*	52.8±4.9	41.5 <sup>#</sup> ±4	0.001*
GST(Unit/100 g tissue)	17.3±2.6	16.3±2.6	0.518	37.5±2.7	36.8±4.5	0.764
Total Protein(mg/L)	15.3±2.2	14.7±1.9	0.580	19.8±2.3	18.5±1.9	0.298

p<sub>1</sub>: p value for **Student t-test** for comparing between the two seasons in each location

#: Statistically significant between the two seasons in each location

\*: Statistically significant at  $p \leq 0.05$

**Figure (2): Comparison between the two seasons in each location according to different antioxidant parameters.**

### Physicochemical analysis:

The ecological investigation of water was done as (salinity, dissolved oxygen (DO) and PH). The results showed insignificant increase in

these parameters in sea water samples collected from the studied areas Abo-Qir and Miami during summer and winter (Table 4).

**Table (4): Comparison between the two locations in each season according to physicochemical parameters.**

	Winter		P	Summer		P
	Abo-Qir (n = 6)	Miami (n = 6)		Abo-Qir (n = 6)	Miami (n = 6)	
<b>Salinity(S%)</b>	35.1 ± 1.4	34.5 ± 2.1	0.573	36.4 ± 1.2	35.2 ± 1.8	0.204
<b>pH</b>	8.5 ± 1.2	8 ± 1.3	0.505	8.4 ± 1.1	8.2 ± 1.3	0.780
<b>Dissolved Oxygen (mg/L)</b>	5.6 ± 0.6	6 ± 0.7	0.313	5.8 ± 0.9	6.4 ± 1.1	0.326

p<sub>1</sub>: p value for **Student t-test** for comparing between the two locations in each season

@: Statistically significant between the two locations in each season


\*: Statistically significant at  $p \leq 0.05$

### Histology and Ultrastructure:


Histopathological study of respiratory tree and digestive tract of sea cucumber *Holothuria polii* showed some abnormal changes like external melanized focal lesions of varying size and severity tissues under light microscope revealed micro colonies of round, basophilic to purple intra-cytoplasmic inclusion bodies are

detected in the digestive tract (Fig.3) and respiratory tree (Fig.4) of Sea cucumber. On the other side, examination of tissue using transmission electron microscope revealed the presence of Rickettsia-like organism (RLOs) as round dense inclusion inside the respiratory tree and the connective tissue of digestive tract (Fig 5&6).







**Fig (3):** Photomicrograph showing Rickettsia-like organism (RLOs) in the digestive tract in sea cucumber *Holothuria polii*.



**Fig (4):** Photomicrograph showing Rickettsia-like organism (RLOs) in the epithelia of respiratory tree in sea cucumber *Holothuria polii*.



**Fig (5):** Transmission electron micrograph showing RLOs in the respiratory tree of sea cucumber *Holothuria polii*. Bar equal 500nm.



**Fig (6):** Transmission electron micrograph showing RLOs.in the digestive tract of sea cucumber *Holothuria polii*.Bar equal 500nm.

## **DISCUSSION:**

The toxicity of pollutants to the organisms is partially mediated by the production of reactive oxygen species (ROS). When production of ROS exceeds the inherent capacity of an organism's antioxidant defence systems, oxidative stress ensues (Prevodnik *et al.*,2007). Heavy metal accumulation, which represent one of the most widespread and serious forms of environmental contamination, are known to induce oxidative stress by way of enhancement of intracellular reactive oxygen species (ROS) production, which often preludes the onset of alterations, such as protein carboxylation and DNA damage (Nunes *et al.*,2017).

The results of the present work indicated that the increased concentration of heavy metals (Zn, Mn, Pb, Cu and Cd) in the tissue of sea cucumber, *Holothuria polii* collected from Abo-Qir station as an industrial area compared with Miami station as a reference location. The elevated bio-accumulation of the heavy metals evoked the antioxidant response in the tissue of the animal, but with increased intracellular

ROS and lipid peroxidation production (measured by MDA) the antioxidant defence system have been obstructed. According to the current results, the level of malondialdehyde (MDA) was significantly increased in the body wall tissue of sea cucumber collected from Abo-Qir station as compared with Miami station. In contrast to MDA, the activities of the antioxidant parameters catalase (CAT), superoxide dismutase (SOD), Glutathione reductase (GSH), and Glutathione transferase (GST), showed a significant decrease in the tissue of sea cucumber collected from Abo-Qir station as compared with Miami station. The present finding is in agreement with those of previous studies (Pourahmad., *et al.*2001; Collen *et al.*,2003; Lee and Shin 2003) who stated that, heavy metals pollution have been previously regarded as provoking factors of oxidative stress. Also, the current results agreed with Hu (2000) who concluded that the heavy metals, such as lead, cadmium, copper, zinc, mercury and vanadium, had the ability to produce reactive oxygen species (ROS), resulting in

lipid peroxidation and antioxidant enzymes alterations, leading to oxidative stress. The present pollutant biomarkers data indicated that, the degree of pollution increased in Abo-Qir station compared with Miami station, and increased in summer as compared with winter season in the two locations. This is may be ascribed to the dumping of toxic industrial waste filled with heavy metals in the area of Abu Qir Bay compared to the open sea water of Miami station. Also, the number of resorts, the movement of industrial ships, its cleaning, painting, and shedding their wastes increased in summer than winter.

The protein content of the sea cucumber was heavily influenced by environmental contaminants, where it non-significantly decreased in Abo-Qir station as an industrial area compared with Miami station. The current results were in accordance with Shacter *et al.*, (1994 a, b) who found that exposure to ROS is known to cause modifications to amino groups of proteins and to alter protein structure or function. One such modification is the formation of carbonyl moieties at amino acid side chains (Shacter *et al.*, 1994 a, b).

The physicochemical parameters (pH, temperature and salinity) in the present study recorded no significant results. Temperature, salinity, diet and individual variation are among other factors affecting accumulation of heavy metals and consequently made cellular oxidative stress (Steniford, 2006). El Moselhy *et al.*, (2005) reported that different animals in the same community at the same trophic levels could accumulate pollutants differently due to differences in habitat physical and chemical properties. The present study indicated that the decrease in pH-value of water at Abo-Qir as ( $8.4 \pm 1.1$ ) related with the drop in oxygen content ( $5.8 \pm 0.9$ ) and this agreed with Nessim *et al.*, (2005) that recorded pH-values of water at Abo-Qir site as (7.57-8.26). Salinity reflects the degree of contamination in aquatic environment

(Zyadah *et al.*, 2004). The present study showed that pollution reflect salinity variation in Abo Qir station and this agreed with (Abou-Taleb *et al.*, 2004) who stated that El-Max Bay area exhibited wide fluctuation in salinity affected by discharge of huge amounts of agricultural, sewage and industrial waste waters.

Rickettsia infections were also found to be significantly higher in deep-sea mussels exposed to petroleum seeps compared to shallow water individuals, although it has yet to be demonstrated that this difference is a direct reaction to contaminant exposure (Powell *et al.*, 1999). In the current study the histopathological examination of respiratory tree and digestive tract of *Holothuria polii* collected from Abo-Qir in summer showed an infection with rickettsia-like organisms (RLOs) and this with agreement with Romero *et al.*, (2000) who stated that the rickettsia-like organism (RLO) revealed the presence of different developmental stages and these included a rod-shaped and uniformly electron dense elementary body (EB) and an intermediate body (IB).

Rickettsia-like organism (RLO) colonies were observed inside small granular cells. Azevedo *et al.*, (2005) showed that the ultrastructural morphology of rickettsia-like organisms (RLOs) present in gill epithelial tissue of the oyster, *Crassostrea rizophorae*, from the estuarine region of the Parnaíba river, on the northeastern Atlantic coast of Brazil. Numerous rod-shaped RLOs formed micro colonies that were located in intracytoplasmic vacuoles up to 85 µm in diameter. These RLOs, which measured about 2 µm x 0.6 µm, had ultrastructural characteristics of prokaryotes that included a plasma membrane and a thin, Gram-negative type cell wall. Some non-dividing RLOs had a transversal constriction indicative of binary fission. Numerous free RLOs were seen following disruption of the vacuoles during

host cell necrosis and degeneration. The presence of intracellular prokaryotes has been described in several marine invertebrates (Harshbarger *et al.*, 1977 & Hine and Diggles 2002), being suggested in some occasions an association with mortalities (Gulka *et al.*, 1983). Some authors indicated a key role in the appearance of disease and mortality (Gulka *et al.*, 1983 & Villalba *et al.*, 1999), whereas other authors suggested that these microorganisms do not cause damages to the host cells or cause just a limited pathology.

The conclusion of this study could be summarized as, the decreased activity of the antioxidant defence system in the sea cucumber, *Holothuria polii* may be ascribed to both; the infection with rickettsia-like organisms (RLOs) and the bio-accumulation of the heavy metals in the animal tissues which may enhance the production of ROS causing disruption in the oxidative homeostasis in the animal tissues.

#### **REFERENCES:**

- **Abou-Taleb AEA., Akel MM., Nessim RB., and Ramadan MH. (2004).** Assessment of some heavy metals and their Accumulation in Marine Organisms in Alexandria coastal Environment, Master Thesis, High Institute of Public Health, Alexandria University 260 p.
- **Azevedo C., Mendonça I., and Matos E. (2005).** Ultrastructural analysis of Rickettsia-like organisms in the oyster *Crassostrea rizophorae* from the Northern Atlantic Coast of Brazil.
- **Beauchamp C., and Fridovich I. (1971).** Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal. Biochem.*, 44(1): 276-87.
- **Bell TA., and Lightner DV. (1988).** A handbook of normal penaeid shrimp histology. World Aquaculture Society, Baton Rouge, LA.
- **Beutler E., Duron O., and Kelly D.O. (1963).** Improved method of determination of blood glutathione. *J. Lab. Clin. Med.*, 61(5):882-888.
- **Bordbar S., Anwar F., and Saari N. (2011).** *Mar. Drugs*, 9, 1761-1805.
- **Branson EJ., and Diaz-Munoz DN. (2006).** Description of a new disease condition occurring in farmed coho salmon, *Oncorhynchus kisutch* (Walbaum), in South America. *Journal of Fish Diseases* 14: 147-156.
- **Claiborn A. (1986).** Catalase activity. In: Greenwald RA (ed) *CRC handbook of methods for oxygen radical research* (2nd edn). CRC press, Boca Raton, pp 283-284.
- **Collen J., Pinto E., Pedersen M., and Colepicolo P. (2003).** Induction of oxidative stress in the red macroalga *Gracilaria tenuistipitata* by pollutant metals. *Arch Environ Contam Toxicol* 45:337-342.
- **Delle Chiaje S.,** *Memorie dell'Accademia di Scienze e Lettere di Napoli*, 1823, 4.
- **Edgerton BF., (2002).** A review of international biosecurity policy development in relation to movements of freshwater crayfish. *Bull. Fr. Pêche Piscic.*, 367, 805-812.

- **Edgerton BF., Evans LH., Stephens FJ., and Overstreet RM., (2002).** Synopsis of freshwater crayfish diseases and commensal organisms. *Aquaculture*, 206, 57- 135.
- **El MoselhyKhM., and Yassien MH. (2005).** Accumulation patterns on heavy metals in Venus clams, *Paphia undulate* (Born, 1780) and *Gafrarium pectinatum* (Linnaeus, 1758), from Lake Timsah, Suez Canal, Egypt. *Egypt. J. Aquat. Res.*, Vol. 31, No.1.
- **Ferrantini F., Fokin SI., Modeo L., Andreoli I., Dini F., Gortz HD., et al. (2009).** “Candidatus Cryptoprodotis polytropus,” A novel Rickettsia-like organism in the ciliated protest *Pseudomicrotho raxdubius* (Ciliophora, Nassophorea). *Journal of Eukaryotic Microbiology* 56: 119–129.
- **Ferreira M., Moradas FP., and Reis HMA. (2005).** Oxidative stress biomarkers in two resident species, mullet (*Mugilcephalus*) and flounder (*Platichthys flesus*), from a polluted site in River Douro Estuary. *Portugal. Aquat. Toxicol.*, 71: 39–48.
- **Ferreira M., Moradas FP., and Reis HMA. (2007).** The effect of long-termdeuration on levels of oxidative stress biomarkers in mullets (*Mugilcephalus*) chronically exposed to contaminants. *Mar. Env. Res.*, 64: 181–90.
- **Filho DW. (1996).** Fish antioxidant defences—a comparative approach. *Braz. J. Med. Biol. Res.*, 29: 1735– 1742.
- **Grasshoff K. (1976).** Methods of sea water analysis. Verlag. Chemie. Chapter 4; Determination of oxygen.
- **Gulka G, Chang PW, and Marti KA (1983).** Prokaryotic infection associated with a mass mortality of the sea scallop *Platopecten magellanicus*. *Journal of Fish Diseases* 6: 355–364.
- **Habig WH., Pabst MJ., and Jakoby WB. (1974).** Glutathione S transferases: The first enzymatic step in mercapturic acid formation. *J. Biol. Chem.*, 249: 7130–7139.
- **HarshbargerJC., Chang SC., and Otto SV. (1977).** Chlamydiae (with phages), mycoplasmas, and rickettsiae in Chesapeake Bay bivalves. *Science*.196, 666-668.
- **Hine PM., and Diggles BK. (2002).** Prokaryote infections in the New Zealand scallops *Pecten novae zelandiae* and *Chlamys delicatula*. *Dis Aquat Org*;50: 137-144.
- **Hu CJ.,Xu YH., Wen J., Zhang LP., Fan SG., and Su T. (2010).** Larval development and juvenile growth of the sea cucumber *Stichopus sp.* (Curry fish). *Aquaculture* 300(1):73–79
- **Hu H. (2000).** Exposure to metals. *Occup. Env. Med.* 27: 983–996
- **JulshamnK., and Grahl-Nielsen O. (1996).** Distribution of trace elements from industrial discharges in the Hardangerfjord, Norway: a multivariate data analysis of saithe, flounder and blue mussel as sentinel organisms. *MarinePollutionBulletin*. 32: 546- 571.

- **Langston WJ., and Spence SK. (1995).** Biological factors involved in metal concentrations observed in aquatic organisms. In: Tessier A., Turner DR. (Eds.), Metal Speciation and Bioavailability. John Wiley and Sons Ltd., pp. 407-478.
- **Lauckner G. (1983).** Diseases of Mollusca: Bivalvia. In: Kinne O (ed) Diseases of marine animals, Vol 2. Biologische Anstalt Helgoland, Hamburg, p 447-520.
- **Lee MY., and Shin HW. (2003).** Cadmium-induced changes in antioxidant enzymes from the marine alga *Nannochloropsis oculata*. J Appl. Physiol 15:13-19
- **Lowry OH., Rosebrough NJ., Farr AL., and Randall RJ. (1951).** Protein Measurement with the Folin Phenol Reagent. J. Biol. Chem., 193: 265-275.
- **Luo W., Lu Y., Wang T., Hu W., Jiao W., Naile JE., and Giesy JP. (2010).** Ecological risk assessment of arsenic and metals in sediments of coastal areas of northern Bohai and Yellow Seas, China. Ambio 39(5-6):367-375
- **Muduli BP., Panda CR. (2010).** "Physico chemical properties of water collected from Dhamra estuary", Intern. J. Environ. Sci., 1(3), pp 334-342.
- **NessimRB., Masoud MS., and Maximous N. (2005).** Water Characteristics of Alexandria Hot Spots. Eyp. J. Aqu. Research, 31, 25- 37.
- **Nunes B., Nunes J., Soares AMVM., Figueira E., and Freitas R. (2017).** Toxicological effects of paracetamol on the clam *Ruditapes hilippinarum*: exposure vs recovery. Aquat Toxicol. 192:198-206.
- **Orbea A., Ortiz ZM., Sol'e MPC., and Cajaraville MP. (2002).** Antioxidant enzymes and peroxisome proliferation in relation to contaminant body burdens of PAHs and PCBs in bivalve molluscs, crabs and fish from the Urdaibai and Plentziaestuorios (Bay of Biscay). Aquat. Toxicol., 58: 75-98.
- **Pourahmad J., Ross S., O'Brien PJ. (2001).** Lysosomal involvement in hepatocyte cytotoxicity induced by Cu<sup>2+</sup> but not Cd<sup>2+</sup>. Free Radical Biol Med 30:89-97.
- **Powell EN., Barber RD., Kennicutt MC., and Ford II SE.(1999).** influence in controlling the health, reproduction and PAH body burden of petroleum seep mussels. Deep-Sea Res.PartI-Oceanogr. Res. Pap. 46, 2053-2078.
- **Prevodnik A., Gardestrom J., Lilja K., Elfwing T., McDonagh B., Petrovic N., et al (2007).** Oxidative stress in response to xenobiotics in the blue mussel *Mytilus edulis* L.: evidence for variation along a natural salinity gradient of the Baltic Sea. Aquat Toxicol. 82:63-71
- **Radwan EH., Hamed SS., and Saad GA. (2014).** Temporal and Spatial Effects on Some Physiological Parameters of the Bivalve *Lithophaga lithophaga* (Linnaeus, 1758) from Coastal Regions of Alexandria, Egypt. Open J. Ecolo.; 4, 732-743.
- **Romero X., Turnbull JF. and Jimenez R. (2000).** Ultrastructure and Cytopathology of a Rickettsia-like Organism Causing Systemic Infection in the Red claw Crayfish *Cherax quadricarinatus*

- carinatus* (Crustacea: Decapoda) in Ecuador. Jour.Inver.Patho.;76, 95–104.
- **Shacter E., Williams JA., Lim M., and Levine RL. (1994a).** Differential susceptibility of plasma proteins to oxidative modification: examination by western blot immunoassay. Free Radic Biol Med 17:429–437
  - **Shacter E., Williams JA., Stadtman ER., and Levine RL. (1994b).** Determination of carbonyl groups in oxidized proteins. In: Punchard NA, Kelley FJ (eds) Free radicals—a practical approach. IRL Press, Oxford, p 159
  - **Spanopoulos-Zarco P., Ruelas-Inzunza J., Meza-Montenegro M., Osuna-Sánchez K., and Amezcua-Martínez F. (2014).** Health risk assessment from mercury levels in Bycatch fish species from the Coasts of Guerrero, Mexico (Eastern Pacific). Bull Environ Contam Toxicol 93(3):334–338
  - **Steniford G. (2006).** Disease Interactions and Pathogen Exchange between Farmed and Wild Aquatic Animal Populations A European Network. Cefas Weymouth Laboratory, Weymouth.
  - **Villalba A., CarballalMJ.,López C., Cabada A., Corral L., et al. (1999).** Branchial rickettsia-like infection associated with clam *Venerupis rhomboides* mortality. Dis Aquat Org; 36: 53-60.
  - **Yoshioka T., Kawada K., Shimada T., and Mori M. (1979).** Lipid peroxidation in maternal and cord blood and protective mechanism against activated oxygen toxicity in the blood. Am. J. Obstet. Gynecol., 135:372-376.
  - **Zyadah M., Ibrahim M., and Madkour A. (2004).** Impacts of environmental parameters on benthic invertebrates and zooplankton biodiversity of the Eastern region of Delta coast at Damietta, Egypt. J. Aquat. Biol. And Fish. ; Vol.8, No.4: 37-52.