



BioBacta



Journal of Bioscience and Applied Research

www.jbaar.org



## Morphospecies diversity of soil invertebrates in Cultivated and Uncultivated fields

Abrar Maqtan<sup>1</sup>, Hishamuddin Omar<sup>1</sup>, Muskhazli Mustafa<sup>1</sup>, Nur Ain Izzati M Zainudin<sup>1</sup> and Daljit Singh Karam<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Science, Universiti Putra Malaysia 43400, Serdang, Selangor, Malaysia  
Postal Address of the First Author

<sup>2</sup>Department of Land Management, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

(Email: hishamspirulina@gmail.com)

### Abstract

The soil environment is teeming with living organisms full of biological life and is one of the most abundant and diverse ecosystems on earth. However, due to unsustainable agriculture practices, the fertile topsoil is often lost; the soil becomes useless for agriculture. Therefore, we cannot lose sight on the fundamental role of the soil biological entities in maintaining the sustainability of the soil, and the role of organism's biodiversity in the soil to counter against various stress. Therefore, the main objective of this work is detecting diversity and density of soil invertebrates between cultivated field (Ladang 2) and uncultivated field (CEFS). Results showed that the highest abundance to invertebrates was recorded at CEFS 834 with 3.7 m<sup>2</sup> density and the lowest was in Ladang 2: 225 with 1.2 m<sup>2</sup> density, belonging to 35 different morphospecies from 4 Phylum, 4 subphyla, 6 class, 2 subclass and 19 orders. A total of 15 morphospecies were collected in the cultivated field Ladang 2, while 25 morphospecies were collected in the uncultivated field CEFS. Out of 25 morphospecies found in this area, 10 were exclusive to this environment. Annelida, Amphipoda, Isopoda, Collembola and Hymenoptera were the dominant taxa in CEFS, comprising: 8.15%, 10.55%, 17.98%, 14.14% and 26.97% respectively. But Acarina, Coleoptera and Insect larva 39.68%, 5.95% and 3.97% respectively were the dominated taxa in Ladang 2. The diversity was highest in the CEFS ( $H = 2.29$ ) and lowest in the Ladang 2 field ( $H = 1.74$ ). Unfortunately, many aspects of the soil invertebrates have not been documented in Malaysia. For example, why they become dominant in certain location and what is the impact on that particular ecosystem. This study showed that it is essential to maintain natural habitat because agricultural practices have major influence on soil invertebrate diversity and hopefully through this study able to suggest sound agriculture management to safeguard the soil ecosystems. Due to the limited information pertaining to soil invertebrates in Malaysia and limited scope of this study, it is premature to make any conclusion and therefore warrant further studies.

**Keywords:** cultivated land; diversity; morphospecies; uncultivated land; soil Invertebrates.

### 1 Introduction

Although not seen by the naked eye, the soil environment teems with living organisms and is among the most diverse ecosystems on earth. Soil flora and fauna with their vast range of functions

actively contribute to the development, structure and productivity of the soil (McCauley, 2005). The wide range and the sheer number of soil invertebrates play critical roles in subsoil and topsoil operations, which are also affected by regional factors such as climatic conditions, soil parent matter, elevation, and ground

cover patterns (Grossi & Brun, 1997; Materna, 2004; Çakıra & Makinecib, 2018). One of the soil ecosystem processes is the recycling of nutrients by soil invertebrates which are responsible for the decomposition of organic matter (Mason, 1980; Haag et al., 1985; Lavelle et al., 1992). The role of invertebrates in this operation includes the breakdown and redistribution of plant residues (Correia & Oliveira, 2000), or indirect energizing of microbial activity and the regulation of decaying fungal groups (Moço et al., 2005). Additionally, soil invertebrates also contribute to soil fertility by the decomposition of organic matter. Disruption of the invertebrates community in the soil is the result of changes in the environment (Correia, 2002), very often caused by human activities. Differences in the composition of vegetation, soil management, quality of soil litter, food availability, degree of ecosystem degradation or restoration have significant impacts on the soil invertebrate community (Moço et al., 2005; Silva et al., 2006; Baretta et al., 2008; Dias et al., 2007; Azevedo et al., 2008; Ribas et al., 2011; Cunha Neto et al., 2012). Although some soil biota has the ability to adapt to harsh environmental conditions, total activity generally decreases when conditions degrade beyond the optimal range (McCauley, 2005). A well-managed site for native vegetation and other fauna using natural processes is also conducive for native invertebrates as they are the lowest common denominator of their societies. Biological indicators are genus or groups of species that reflect the influence of habitat alterations due to their susceptibility to environmental conditions (McGeoch, 1998). Since invertebrates are often specific to certain environmental conditions, are abundant and are associated with ecosystem processes, and they may serve as useful indicators. Some species respond more to environmental changes than others or have certain characteristics that make them useful to indicate changes in the ecosystem or to monitor environmental conditions (Hutcheson et al. 1999; Greg, and Alison, 2016). However, the major and most diverse community groups of native invertebrates are those found in relatively undisturbed or natural sites (Patrick, 1994). The interactions between soil environmental differences and soil invertebrates are not commonly used in soil quality assessment (Cunha Neto et al., 2012). It is well recognized that excessive human activities such as changes in reforestation and destructive land-use may have significant impacts on biodiversity (Vitousek et al., 1997; Sala et al., 2000; León-Gamboa et al., 2010; Çakıra & Makinecib, 2018). Therefore, human activities and degradation in environmental conditions resulting from global climate change can potentially influence the native communities of soil invertebrates (Nielsen & King, 2015). Management practice may affect the diversity

of invertebrates in the soil, acting through direct quality (litter quality) and indirect effects (microhabitats, environmental factors such as pH, soil moisture and soil fertility) (Kinasih et al., 2016). All these lead to disturbances in the ecosystem or ecological imbalance such as the breakdown of forest areas leading to the direct extinction of some species (Loyola et al., 2006). Malaysia is developing country in the tropics where local soil fauna is often unappreciated and suffers from human influence. In the quest for sustainable living, there is need for more scientific information that can facilitate political decisions to be made related to soil conservation. Past studies have shown that human activities often lead to soil deterioration and reduction in the abundance of soil fauna. In most cases, the stress-tolerant species prevail and scarce varieties fall in abundance or totally disappear as the result of intensification in agriculture (Fauziah et al., 1997; Eisenbeis, 2006; Jeffery et al., 2010; Menta, 2012). However, there is a lack of studies about soil diversity of invertebrate communities and aspects of soil biology in Malaysia. Thus; this paper offers new data about soil invertebrates communities of two locations in Malaysia that have different soil characteristics. The main objective was to make a comparison of the morphospecies composition, diversity and density of the soil fauna between a natural or untouched site of management practice (CEFS) and a cultivated site with some farming and human activities (Ladang 2).

## **2 Materials and Methods**

### **2.1. Study area**

This study was carried out in August, 2017 and January 2018, in Malaysia. Two study areas were chosen and compared – one untouched by human activity and soil impacted by human activities. The first is the Ladang 2, which is a research plot in the Universiti Putra Malaysia (UPM) campus, with some farming and other human activity (300°31.22'N, 101042°12.92'E). The second plot is known to be a grass area behind the Faculty of Environmental Studies, UPM, with no known human activities (300°22.38'N, 101042°29.57'E).

### **2.2. Procedures**

Two methods were used to collect samples: pitfall traps for invertebrate's surface soil (Maftu'ah et al, 2005), and Berlese funnels to separate micro invertebrates from soil samples (Southwood, 1994). The large invertebrates were then sorted manually in the laboratory. In each sampling plot, a quadrant (of 15 x 15m) was selected randomly and further divided into 9 sub plots (5x5m). The distance between traps

was 5 meters. The trap was made from plastic bottle and cut into 10.5 cm length and 7.5 cm width and carefully placed in holes at soil level. Each trap contained about 50ml detergent solution and left in the field for about 24 hours. Another 9 soil core samples (5cm depth) were taken at random from each plot to collect micro invertebrates, brought to the laboratory and placed in Berlese funnels (Çakır & Makineci, 2018). All samples collected were stored in a jar containing 70% ethanol. From each sample, with the aid of a dissecting microscope, the number of invertebrates was counted. The collected invertebrates were then identified to the possible taxonomic level (using dissecting microscope), applying the ecological pattern (morphologically-based classification) based on the dichotomous key classification and categorized as adult or immature (Wade et al., 1989; Thyssen, 2010). Record of abundance in each sample was done by counting the total number for each taxon or morphospecies, in the laboratory of plant physiology, Faculty of Science, UPM.

### 2.3. Data analysis

#### 2.3.1. Taxonomic groups

The results of the invertebrate community analyses were recorded for each soil sample, as density (i.e., individual number/m<sup>2</sup> soil) and relative abundance (i.e., individual number of each taxon/individual number) (Santorufu et al., 2012).

#### 2.3.2. Biological indices

The species diversity was calculated by the Shannon Diversity Index (Shannon, 1949). Data were entered into the spreadsheet that could be set up to automatically calculate the diversity index

$$\text{Shannon Index (H)} = -\sum_{i=1}^s p_i \ln p_i$$

p = proportion (n/N), n= one particular species found, N = total number of individuals found.

ln= natural log, Σ = sum of the calculations. s =number of species.

Dominance indicators were according to the Simpson index (Simpson, 1949): Simpson Index (D)

$$= \frac{1}{\sum_{i=1}^s p_i^2}$$

Similarity index: Sorenson's index (QS) =  $\frac{2 \cdot C}{A+B}$

C = number of species the two communities have in common. A = total number of species found in community 1, B= total number of species found in community 2.

#### 2.3.3. Statistical analysis

Statistical analysis was performed with IBM SPSS statistics version 23. Data of invertebrates after normality test were tested using Kruskal Wallis test and value of **P > 0.05** was considered to be the level of statistical significance.

## 3 Results

### 3.1. Taxonomic groups

A total of 1,086 individuals were collected during the sampling period. Among the two study areas, CEFS had the highest abundance of individuals totaling 834 (Table1), followed by Ladang 2 with 252 individuals. Table2 shows morphospecies classification into four Phylum: (Annelida, Nematoda, Arthropoda, and Mollusca), four subphyla (Crustacea, Myriapoda, Chelicerata and Hexapoda), six classes (Oligochaeta, Gastropoda, Malacostraca, Arachnida, Entognatha and Insecta), two subclasses (Acari and Collembola) and 19 orders.

### 3.2. Density of invertebrate categories:

Table 3 shows the density between invertebrates from different categories. Highest density was in CEFS with 3.70 individual/m<sup>2</sup>. Invertebrate categories with high densities included Hymenoptera, Amphipoda, Collembola, Isopoda and Annelida comprising 1.00, 0.66, 0.52, 0.39 and 0.30 individual/m<sup>2</sup> respectively. Overall, invertebrates density in Ladang 2 was 1.12 individual/m<sup>2</sup> represented by Acarina, Coleoptera and Insect larva comprising 0.44, 0.06 and 0.044 individual/m<sup>2</sup> respectively. Kruskal Wallis test indicated that mean abundance of soil invertebrates at CEFS was highly significant at (P= .000) than Ladang 2 (P=0.05) (Table 4).

Table 1: Composition and abundance of soil invertebrates that collected by traps and funnels from study sites

Taxon	CEFS			Ladang 2			Grand Total
	Traps	Funnels	Total	Traps	Funnels	Total	
Annelida	63	5	68	4	NF	4	72
Nematoda	19	5	24	4	4	8	32
Isopoda	88	*NF	88	NF	NF	NF	88
Amphipoda	150	NF	150	NF	NF	NF	150
Myriapoda	8	1	10	2	NF	2	12
Acarina	14	17	31	47	53	100	131
Araneae	16	1	17	6	4	10	27
Gastropoda	11	NF	11	NF	NF	NF	11
Collembola	112	6	118	5	2	7	125
Diplura	2	NF	2	NF	NF	NF	2
Diptera	31	1	32	2	NF	2	34
Psocoptera	6	NF	6	1	NF	1	7
Coleoptera	12	Nf	12	15	NF	15	27
Hymenoptera	224	1	225	59	21	80	305
Mecoptera	7	Nf	7	1	NF	1	8
Orthoptera	2	Nf	2	4	NF	4	6
Hemiptera	12	Nf	12	6	NF	6	18
Isoptera	1	Nf	1	2	NF	2	3
Ephemeroptera	1	Nf	1	NF	NF	Nf	1
Insect larvae	4	1	5	8	NF	10	15
Plecoptera	3	Nf	3	NF	NF	Nf	3
Dermaptera	2	Nf	2	NF	NF	Nf	2
Thysanoptera	2	1	3	NF	NF	Nf	3
Blattodea	2	Nf	2	NF	NF	Nf	2
Siphonaptera	1	1	2	NF	NF	NF	2
Total	793	41	834	166	86	252	1086

\*Nf=not found

Table2: Morphospecies classification and abundance of soil invertebrates in UPM

Phylum	Subphylum	Class	Subclass	Order	number	
Annelida		Oligochaeta			72	
Nematoda					32	
Mollusca		Gastropoda			11	
Arthropoda	Crustacea	Malacostraca		Isopoda	88	
				Amphipoda	150	
		Myriapoda				12
	Chelicerata	Arachnida		Acari	Acarina	131
					Araneae	27
	Hexapoda	Entognatha	Collembola			125
					Diplura	2
			Insecta		Diptera	34
					Psocoptera	7
					Coleoptera	27
				Hymenoptera	305	
				Mecoptera	8	
				Orthoptera	6	
				Hemiptera	18	
				Isoptera	3	
				Ephemeroptera	1	
				Plecoptera	3	
				Dermaptera	2	
				Thysanoptera	3	
				Blattodea	2	
				Siphonaptera	2	

### 3.3. Invertebrates diversity

Table 4 shows that the diversity index for CEFS with  $H = 2.3$  was more diverse than Ladang 2 with  $H = 1.7$ . Both sites have similar 15 morphospecies, therefore the similarity index between the n two sites was  $QS = 0.75$ . The prominent taxa

were Annelida, amphipoda, isopoda, collembola and hymenoptera were the dominant taxa in CEFS: 8.15%, 10.55%, 17.98%, 14.14% and 26.97% respectively. In Ladang 2, the prominent taxa were Acarina, Coleoptera and insect larva: 39.68%, 5.95% and 3.97% respectively (Table 4).

Table 3: The density of invertebrates categories in both sites

Taxon	CEFS	Density	Ladang 2	Density
Annelida	68	0.30	4	0.02
Nematoda	24	0.11	8	0.04
Isopoda	88	0.39	NF	0.00
Amphipoda	150	0.67	NF	0.00
Myriapoda	10	0.04	2	0.01
Acarina	31	0.14	100	0.44
Araneae	17	0.08	10	0.04
Gastropoda	11	0.05	NF	0.00
Collembola	118	0.52	7	0.03
Diplura	2	0.01	NF	0.00
Diptera	32	0.14	2	0.01
Psocoptera	6	0.03	1	0.00
Coleoptera	12	0.05	15	0.07
Hymenoptera	225	1.00	80	0.36
Mecoptera	7	0.03	1	0.00
Orthoptera	2	0.01	4	0.02
Hemiptera	12	0.05	6	0.03
Isoptera	1	0.00	2	0.01
Ephemeroptera	1	0.00	NF	0.00
Insect larvae	5	0.02	10	0.04
Plecoptera	3	0.01	NF	0.00
Dermaptera	2	0.01	NF	0.00
Thysanoptera	3	0.01	NF	0.00
Blattodea	2	0.01	NF	0.00
Siphonaptera	2	0.01	NF	0.00
Total	834	3.71	252	1.12

Table 4: The diversity index for CEFS and Ladang 2

Taxon	Sites			
	CEFS		Ladang 2	
	Mean	Percentage	Mean	Percentage
Annelida	7.55±0.72 <sup>a</sup>	8.15%	0.44±0.17 <sup>b</sup>	1.59%
Nematoda	2.66±0.23 <sup>a</sup>	2.90%	0.88±0.11 <sup>b</sup>	3.17%
Isopoda	9.77±0.87 <sup>a</sup>	10.55%		
Amphipoda	16.66±0.80 <sup>a</sup>	17.98%		
Myriapoda	2.50±1.00 <sup>a</sup>	1.19%	0.50±0.28 <sup>b</sup>	0.79%
Acarina	3.44±0.33 <sup>b</sup>	3.71%	11.11±1.0 <sup>a</sup>	39.68%
Areaneae	1.88±0.53 <sup>a</sup>	2.03%	1.11±0.11 <sup>b</sup>	3.97%
Gastropoda	3.66±1.20 <sup>a</sup>	1.31%		
Collembola	13.11±1.60 <sup>a</sup>	14.1%	0.77±0.27 <sup>b</sup>	2.77%
Diplura	1.00±000 <sup>a</sup>	0.24%		
Diptera	4.00±1.050 <sup>a</sup>	3.83%	0.25±0.16 <sup>b</sup>	0.79%
Psocoptera	3.00±000	0.71%	0.50±0.50 <sup>a</sup>	0.39%
Coleoptera	1.50±0.32 <sup>b</sup>	1.43%	1.87±0.22 <sup>a</sup>	5.95%
Hymenoptera	20.66±8.80 <sup>a</sup>	26.90%	8.88±0.73 <sup>b</sup>	31.7%
Mecoptera	2.33±0.88 <sup>a</sup>	0.83%	0.33±0.33 <sup>b</sup>	0.39%
Orthoptera	0.33±0.16 <sup>b</sup>	0.23%	0.44±0.24 <sup>a</sup>	1.59%
Hemiptera	1.28±0.18 <sup>b</sup>	8.15%	0.85±0.26 <sup>a</sup>	2.38%
Isoptera	4.00±00	2.90%	2.00±0.00	0.79%
Ephemeroptera	1.00±000	10.5%		0.43%
Insectlarva	0.57±0.29 <sup>b</sup>	0.59%	1.42±0.20 <sup>a</sup>	3.97%
Plecoptera	1.33±0.33	1.19%		
Dermaptera	1.00±000	3.71%		
Thysanoptera	1.00±000	2.03%		
Blattodea	1.00±000	1.31%		
Siphonaptera	0.33±0.500 <sup>a</sup>	14.14%		
Shannon Index	2.3		1.7	
Simpson index	6.6		3.7	
Sorenson's index		0.75		

All data are mean ±SE (n=140) means followed by a different letter are significantly at P<0.05

### 3.4. Function groups

To simplify the explanation, the soil invertebrates were divided into three size categories (Lavelle et al 1996) (Figure 1). The first category, also known as macrofauna with size of 10mm or more, forms 25.30% of soil invertebrates in CEFS and only 5.40% in Ladang 2 (Figure 2). The macrofauna (Group 1) comprises Annelida-like earth worms, Mollusca, Myriapoda, Hymenoptera like ants and Isoptera or termites, and another Insect like Orthoptera, Coleoptera and Diptera. Mesofauna (Group 2), the intermediate size between macrofauna and Microfauna (Group 3) comprise 6.70% in CEFS and 4.22% in Ladang 2. Members of Mesofauna were Small Annelida, Acarina, Collembola, small Myriapoda and diplura. The final group was Microfauna which consisted of only 0.91% in CEFS and 0.31% in Ladang 2. The three groups can be categorized based on how they interact with their habitat. The macrofauna, also known as the ecosystem engineer (Group A) comprises 11.60% in CEFS and 3.39% in Ladang 2. They consist mainly of earth worms, ants and termites. The second functional group was known as Litters transformer groups (Group B) which made up 15.15% in CEFS and 7.37% in Ladang 2. The Litter transformer consisted of Mesofauna (small Annelida and Myriapoda, Acarina and Collembola) and some smaller Macrofauna. The final functional group, also known as Predatory group (Group C) made up 12.11% in CEFS and 7.49% in Ladang 2, including one microfauna-like nematoda with some Mesofauna-like Acarina and many Macrofauna such as Myriapoda, Hymenoptera and Araneae.

### 4. Discussion

Choosing two contrasting sampling areas provided the opportunity to test hypotheses regarding the impact of land activities on invertebrate communities. Findings of this study provide ample evidence about the ecological significance of decomposer assemblages in uncultivated as well as cultivated areas. Interaction of invertebrates and soil is the key to the functioning of the ecosystem. Although soils serve as suitable habitat for invertebrates, some of these organisms also cause changes in soil texture and structure by integrating soil organic matter (Zech, 1993; Pappoe et al., 2009). Arthropoda formed the largest phylum (Tables 2, 3) indicating it is the most successful and forming the largest group of invertebrates [Postlethwait et al.,

1991). However, when the soil environment is disturbed, Mesofaunal (such as Acari, some smaller insect groups, and smaller Araneae) become more and more dominant. The Acari (mites) are an extremely diverse group of arachnids which can successfully adapt to a wide range of habitats (Blakely et al., 2002; Eisenbeis, 2006), in both habitats. However, findings suggest CEFS habitat is more hospitable to different soil fauna than Ladang 2. It is commonly observed that communities living in undisturbed or moderately disturbed habitat were higher and more diverse than those living in habitat frequently disturbed and causing changes in texture and structure, exposed to fertilizer and different pesticides and herbicides applications (Prashar & Shah, 2016). It is recommended not to make hasty comparison pertaining to soil communities as other factors beyond the scope of this paper might have some influence on the outcome of species composition and diversity. Turbe et al. (2010) explained that contrary to popular belief, disturbances do not necessarily lead to long-term loss of biodiversity, and in many cases, moderate stress can be a positive force, enabling species to coexist, thereby increasing biodiversity. Perhaps this is what happened to the uncultivated field CEFS. Although there were some disturbances in CEFS, because the soil has some grooves or indentations to slow down surface runoff and soil erosion, the soil was covered with wild grass that provided food and shelter for a variety of soil fauna. That is probably the reason why CEFS has higher fauna species composition and diversity than Ladang 2. To show that land use and soil type can influence fauna number and diversity. On the other hand, Ladang 2 is the agriculture experimental plot which has been ploughed and tilled from time to time, received fertilizer, herbicide and pesticide periodically whereas soil in CEFS is virtually undisturbed, free from agricultural activities such as soil tilling, herbicide and pesticide. Land disturbance and the use of herbicide and pesticide are the most probable agents affecting diversity (Menta, 2012). This paradox is resolved by the average nuisance hypothesis, which posits that biodiversity is highest when the stress is mediocre. The main idea is that, with low disturbance, competitive exclusion arises through dominant species, while with severe disturbance, only stress-tolerant species can survive. Also, in the cultivated field Ladang 2 the intensive tillage and agricultural activity in this site led to increased stress on the diversity of invertebrates, so, the diversity was low. The present findings are in contrast to Frainer and Duarte, (2009). There was a difference in the



Coleoptera between the two communities where the density was high in the disturbed fields compared to the undisturbed fields, where some coleopterans were resistant to the disturbance (Ronqui & Lopes 2006; Frainer & Duarte, 2009). Furthermore, there could be several explanations for differences like the physical properties of the soil, soil cover, soil moisture, soil organic matter that can be the determining factors. It has been argued that communities that are subjected to little disturbance or no nuisance in species compared to those experiencing severe conditions are often produced by pesticide and fertilizer applications and changes in structure and texture (Smith et al., 2008; Abdu et al., 2008; Poppoe et al., 2009). Macrofauna was the highest group in soil invertebrates in both site. furthermore, this high rate was related to hymenoptera, which have more influencing the composition of the faunal communities and they have high dispersal ability in different environments (Mateos et al., 2011; Manhães et al.,2013)

### Conclusion

The findings of this study suggest that cultivated fields are generally lower in the number of soil organisms and diversity compared to uncultivated land. Since there are not many similar studies available in the literature, the present study provides valuable baseline data on the diversity of soil surface invertebrates in Universiti Putra Malaysia specifically at the Centre of Environmental Forensic Studies and the Research Farm (Ladang 2). Knowing the species composition and diversity of soil fauna is the first step towards understanding the state of the soil and its contribution to soil health and crop production.

### Acknowledgements

This research was funded by Putra grant code project: GP/2017/9568000 of UPM Malaysia.

### References

**Abdul, G., Muhammad, H., & Alvi, Z. H.** (2008). Biodiversity of earthworm species from various habitats of District Narowal, Pakistan. *International Journal of Agriculture and Biology*, 10(6): 681-684.

**Azevedo, V.F, Pereira, M.G, Corrêa Neto, T.A, Schermack, V, & Machado, D.L.**(2008). Changes in the of the edaphic fauna community due to the burning in secondary forest in Flona Mário Xavier, Seropédica, RJ. *R. Ci. Vida*, 28: 09-17.

**Baretta D, Duarte C.R, Baretta M, & Cardoso**

**N.E.J.B.** (2008). Multivariate attributes analysis in a high Andean forest. *Rev. Biol. Trop*, 58:1031–1048.

**Blakely JK, Neher DA, Spongberg AL.** (2002). Soil invertebrate and microbial communities, and decomposition as indicators of polycyclic aromatic hydrocarbon contamination. *Applied Soil Ecology*, 21(1):71-88.

**Çakır, M., & Makineci, E.** (2018). Community structure and seasonal variations of soil microarthropods during environmental changes. *Applied Soil Ecology*, 123: 313-317.

**Correia, M.E, & Oliveira, L.C.M.** (2000). Soil Fauna: Aspects general and methodological. *Seropédica, Embrapa Agrobiologia*, ;112:46.

**Correia, M.E.F.** (2002). Relations between the diversity of wildlife and the process of decomposition and its reflections on the stability of ecosystems. *Seropédica, Embrapa Agrobiologia*:156 ,33.

**Cunha Neto, F. V. D., Correia, M. E. F., Pereira, G. H. A., Pereira, M. G., & Leles, P. S. D. S.** (2012). Soil fauna as an indicator of soil quality in forest stands, pasture and secondary forest. *Revista Brasileira de Ciência do Solo*, 36(5): 1407-1417.

**Dias, P. F., Souto, S. M., Correia, M. E. F., de Menezes Rodrigues, K., & Franco, A. A.** (2007). Effect of legume trees on soil macrofauna of a brachiaria brizantha cv. marandu pasture. *Pesquisa Agropecuária Tropical*, 37(1): 38.

**Eisenbeis, G.** (2006). Biology of soil invertebrates. In *Intestinal microorganisms of termites and other invertebrates* Springer, Berlin, Heidelberg: 3-53.

**Fauziah, C. I., Jamilah, I., & Syed Omar, S. R.** (1997). An evaluation of cation exchange capacity methods for acid tropical soils. *Pertanika journal of tropical agricultural science*, 20: 113-120.

**Frainer, A., & Duarte, M. M.** (2009). Soil invertebrates in southern Brazilian Araucaria forest-grassland mosaic: differences between disturbed and undisturbed areas. *Iheringia. Série Zoologia*, 99(3): 307-312.

**Greg S. and Alison E.** (2016). Invertebrates: search and extraction methods, Inventory and monitoring toolbox: invertebrates DOCCM-255606.

**Grossi, J. L., & Brun, J. J.** (1997). Effect of climate and plant succession on lumbricid populations in the French Alps. *Soil Biology and Biochemistry*, 29(3-4): 329-333.

**Haag H.P, Valera F.P, Chiaranda R, Kikuti P, Chávez J.M, Donald E.L.F, et al.** (1985). Nutrient cycling in tropical forests. Campinas, Cargill Foundation,: 144.

**Havlin, J. L., Beaton, J. D., Tisdale, S. L., & Nelson, W. L.** (2005). Soil fertility and fertilizers: An

- introduction to nutrient management (Vol. 515, pp. 97-141). Upper Saddle River, NJ: Pearson Prentice Hall.
- Hutcheson, J., Walsh, P., & Given, D. R.** (1999). Potential value of indicator species for conservation and management of New Zealand terrestrial communities Wellington: Department of Conservation: 90.
- Jeffery, S., Gardi, C., Jones, A., Montanarella, L., Marmo, L., Miko, L., ... & Van der Putten, W. H.** (2010). European atlas of soil biodiversity. Publications Office of the European Union, Commission européenne, Luxembourg.
- Kinasih, I., Cahyanto, T., widiana, A., kurnia, D. N. I., julita, U., & putra, R. E.** (2016). Soil invertebrate diversity in coffee-pine agroforestry system at Sumedang, West Java. *Biodiversitas Journal of Biological Diversity*, 17(2).
- Lavelle P.** (1996). Diversity of soil fauna and ecosystem function. *Biology International*. 33:3-16.
- Lavelle, P., Spain, A. V., Blanchart, E., Martin, A., & Martin, S. (1992). Impact of soil fauna on the properties of soils in the humid tropics. *SSSA Special Publication*, 29:157-157.
- León-Gamboa, A. L., Ramos, C., & García, M. R.** (2010). Effect of pine plantations on soil arthropods in a high Andean forest. *Revista de biologia tropical*, 58(3): 1031-1048.
- Loyola, R. D., Brito, S. L., & Ferreira, R. L.** (2006). Ecosystem disturbances and diversity increase: implications for invertebrate conservation. In *Arthropod Diversity and Conservation* Springer, Dordrecht. :25-42.
- Maftu'ah, E., Alwi, M., & Willis, M.** (2005). Potential of soil macrofauna as bioindicator of peat land quality. *Bioscientiae*, 2(1), 1-14.
- Manhães, C. M. C., Gama-Rodrigues, E. F., Moço, M. K. S., & Gama-Rodrigues, A. C.** (2013). Meso- and macrofauna in the soil and litter of leguminous trees in a degraded pasture in Brazil. *Agroforestry systems*, 87(5), 993-1004.
- Mason, C.F.** (1980). *Decomposition - Biology topics*. Sao Paulo, EPU / University of São Paulo,;18: 63.
- Mateos, E., Santos, X., & Pujade-Villar, J.** (2011). Taxonomic and functional responses to fire and post-fire management of a Mediterranean Hymenoptera community. *Environmental management*, 48(5), 1000.
- Materna, J.** (2004). Does forest type and vegetation patchiness influence horizontal distribution of soil Collembola in two neighbouring forest sites? *Pedobiologia*, 48(4): 339-347.
- McCauley A, Jones C, Jacobsen J.** (2005). Basic soil properties. *Soil and water management module*,1(1):1-2.
- Mcgeoch MA.** (1998). The selection, testing and application of terrestrial insects as bioindicators. *Biological reviews*,73(2):181-201.
- Menta, C.** (2012). Soil fauna diversity-function, soil degradation, biological indices, soil restoration. In *Biodiversity conservation and utilization in a diverse world*. InTech.
- Moço, M K S. Gama-Rodrigues, E.F. Gamarodrigues, A.C. & Correia, M.E.F.** (2005). Description edaphic fauna in different vegetation North Fluminense region. *R. Bras. Ci. Solo*, 29: 555-564.
- Nielsen UN, King CK.** (2015). Abundance and diversity of soil invertebrates in the Windmill Islands region, East Antarctica. *Polar Biology*,38(9):1391-400.
- Pappoe AN, Armah FA, Akotoye HK.** (2009). Diversity of soil surface invertebrates in cultivated and uncultivated fields in Coastal Savanna zone of Ghana. *Research Journal of Soil Biology*,1(2):77-83.
- Patrick B.** (1994). The importance of invertebrate biodiversity: An Otago Conservancy review. Department of Conservation.
- Postlethwait JH, Hopson JL, Veres RC.** (1991). *Biology! : Bringing Science to Life*. McGraw-Hill Companies.
- Prashar, P., & Shah, S.** (2016). Impact of fertilizers and pesticides on soil microflora in agriculture. In *Sustainable Agriculture Reviews* (pp. 331-361). Springer, Cham.
- Ribas CR, Schmidt FA, Solar RR, Campos RB, Valentim CL, Schoederer JH.** (2012). Ants as indicators of the success of rehabilitation efforts in deposits of gold mining tailings. *Restoration Ecology*, 20(6):712-20.
- Ronqui, D. C., & Lopes, J.** (2006). Composition and diversity of Scarabaeoidea (Coleoptera) attracted by light trap in the rural areas of Northern Paraná. *Iheringia. Série Zoologia*, 96(1): 103-108.
- Sala OE, Chapin FS, Armesto JJ, Berlow E, Bloomfield J, Dirzo R, et al.** (2000). Global biodiversity scenarios for the year 2100. *science*.;287(5459):1770-4.
- Santorufu L, Van Gestel CA, Rocco A, Maisto G.** (2012). Soil invertebrates as bio indicators of urban soil quality. *Environmental Pollution*, 161:57-63.
- Shannon CE, Weaver W.** (1949).The mathematical theory of information, *AT&T Tech. J*,27,359-423.
- Silva R.F, Aquino A.M, Mercante F.M. & Guimarães, M.F.** (2006). Macrofauna invertebrate soil under different production systems in Latosol of the Region of the Cerrado. *Pesq. Agropec. Bras*, 41: 697-704.

**Simpson E.H** (1949). Measurement of diversity. *Nature*, 163, 688, doi:10.1038/163688a0.

**Smith RG, McSwiney CP, Grandy AS, Suwanwaree P, Snider RM, Robertson GP.** (2008). Diversity and abundance of earthworms across an agricultural land-use intensity gradient. *Soil and Tillage Research*,100(1-2):83-8.

**Southwood T.R.E.** (1994). *Ecological methods*. Chapman & Hall.. ISBN 0-412-30710-3.

**Thyssen P. J.** (2010). Keys for Identification of Immature Insects. *Current Concepts in Forensic Entomology*. :25-42.

**Turbe A, Toni A. D, Benito P., Lavelle P., Lavelle P., Camacho N. R. et al.** (2010). Soil biodiversity: functions, threats and tools for policy makers Rapport de synthèse. <bioemco-00560420> HAL. <http://hal-bioemco.ccsd.cnrs.fr/bioemco-00560420>.

**Vitousek PM, Mooney HA, Lubchenco J, Melillo JM.** (1997). Human domination of Earth's ecosystems. *Science*, 277(5325):494-9.

**Wade KR, Ormerod SJ, Gee AS.** (1989). Classification and ordination of macroinvertebrate assemblages to predict stream acidity in upland Wales. *Hydrobiologia*, 171(1):59-78.

**Zech, W.**( 1993). *Geology and Soils*. In: *Tropical Forestry Handbook*, Pancel, L. (Ed.) Springer-Verlag, New York:515-567. References in the text should be inserted in parentheses in full for single and dual authored papers, but using the first author and et al. for multiple authored papers.

Figure1: Divided soil invertebrates according to body size and function group.




Figure 2: The deferent percentage between invertebrates groups among 2 sites

